

Coyinema gen. n., a new genus of nematode (Thelastomatoidea, Hystrignathidae) parasites of Passalidae (Coleoptera) from Cuba

Jans Morffe Rodríguez†, Nayla García Rodríguez‡

Instituto de Ecología y Sistemática, Carretera de Varona km 31/2, Capdevila, Boyeros, A.P. 8029, C.P. 10800, Ciudad de La Habana, Cuba

† [urn:lsid:zoobank.org:author:6285C0EA-922E-467F-BE19-D50BB7601360](https://zoobank.org/urn:lsid:zoobank.org:author:6285C0EA-922E-467F-BE19-D50BB7601360)

‡ [urn:lsid:zoobank.org:author:B74CF649-3FBC-4862-8B6E-801437F87FEB](https://zoobank.org/urn:lsid:zoobank.org:author:B74CF649-3FBC-4862-8B6E-801437F87FEB)

Corresponding authors: *Jans Morffe Rodríguez* (jans@ecologia.cu), *Nayla García Rodríguez* (nayla@ecologia.cu)

Academic editor: *H.-P. Fagerholm* | Received 12 June 2010 | Accepted 15 December 2010 | Published 12 January 2011

[urn:lsid:zoobank.org:pub:14BAF70D-0A9E-427A-852F-2E37B3465656](https://zoobank.org/urn:lsid:zoobank.org:pub:14BAF70D-0A9E-427A-852F-2E37B3465656)

Citation: Rodríguez JM, Rodríguez NG (2011) *Coyinema* gen. n., a new genus of nematode (Thelastomatoidea, Hystrignathidae) parasites of Passalidae (Coleoptera) from Cuba. ZooKeys 75: 9–19. doi: 10.3897/zookeys.75.809

Abstract

The new genus *Coyinema* **gen. n.** is described as parasite of the two passalid beetles from Cuba: *Passalus interstitialis* Escholtz, 1829 (type host) and *P. pertyi* Kaup, 1869. Females are characterized by the shape of their cephalic end, cervical cuticle unarmed, a sub-cylindrical procorpus with its base abruptly dilated, fore region of intestine dilated as a sac-like structure, genital system didelphic-amphidelphic and eggs markedly ovoid and smooth-shelled. Males have a digestive system similar to females, tail sharply pointed, bearing a Y-like thickening of the dorsal cuticle. They also present a big, median, mammiform pre-cloacal papillae and a pair of small, sub-dorsal pre-cloacal papillae anterior to the cuticular thickening of the tail.

Keywords

Nematoda, Hystrignathidae, *Coyinema*, Passalidae, *Passalus*, new genus

Introduction

The family Hystrignathidae Travassos, 1920 (Oxyurida, Thelastomatoidea) includes about 100 species of parasitic nematodes specific of the gut caeca of passalid beetles. At present, 26 genera are recognized, most of them on the basis of females (Adamson and Van Waerebeke 1992). The shortage of male descriptions is mainly due to the difficulty

in assigning specimens to their correct species in co-infections which are common for passalid beetles. The morphological homogeneity of males further contributes to this difficulty.

The Hystrignathidae have a mostly Gondwanian distribution with species in North, Central and South America, West Indies, Africa, Madagascar, Australia and New Guinea. The majority of the genera have been recorded in Brazil (21), most of them being endemic (Travassos and Kloss 1957a, 1957b, 1958; Kloss 1962, Cordeira 1981, Cordeira and Artigas 1983). Some of these genera are also recorded from Mexico (1), Venezuela (2), Saint Lucia (3) and Trinidad (3) (Guerrero 1980, Hunt 1981, 1982, Coy and García 1995, García and Coy 1997).

Australia shows a characteristic fauna of hystrignathids, with the endemic and monotypic genera *Anuronema* Clark, 1978; *Phalacronema* Clark, 1978 and *Sprentia* Clark, 1978 (Clark 1978). Also, *Lepidonema* Cobb, 1898 and *Xyo* Cobb, 1898 have been recorded (Cobb 1898).

In Cuba, the study of this group is recent (Coy 1990). Currently, members of the genera *Artigasia* Christie, 1934; *Glaber* Travassos & Kloss, 1958; *Hystrignathus* Leidy, 1850; *Lepidonema*, *Longior* Travassos & Kloss, 1958 and *Salesia* Travassos & Kloss, 1958 are recorded (Coy et al. 1993, García and Coy 1994, 1995a, b, García et al. 2009a, b, c, Morffe and García 2009, Morffe and García 2010a, b).

Coy et al. (1993) described *Glaber poeyi* Coy, García & Alvarez, 1993 from *Passalus interstitialis*. Subsequent examination of type material and specimens collected from other localities demonstrate that these belong to a new genus described in this paper.

Materials and methods

Five specimens of *Passalus pertyi* Kaup, 1869 were collected by hand from rotting logs between May, 2007 and March, 2008. Two specimens are from La Melba, Nipe-Sagua-Baracoa, Holguín Province, Cuba, one from El Mulo, Sierra del Rosario, Pinar del Río Province, Cuba and two from La Jaula, San José de las Lajas, La Habana Province, Cuba. Specimens from La Melba and El Mulo were immediately killed and fixed in 70% ethanol. Beetles were dissected by practicing longitudinal incisions in the abdominal pleural membranes and the intestines were extracted and excised in water in Petri dishes under a stereo microscope. The parasites were collected and stored in 70% ethanol.

Beetles from La Jaula were kept alive in plastic jars with moistened wood chips as food and a humidity source. They were killed as soon as possible with ethyl ether vapours and dissected as described above. The intestines were dissected in normal saline instead of water. Nematodes removed from guts were killed with hot water (60–70°C) and fixed in 70% ethanol.

In this study the type material of *Glaber poeyi* deposited in the Colección Helminológica of the Colecciones Zoológicas (CZACC), Instituto de Ecología y Sistemática, Havana, Cuba was included.

Nematodes were clear-mounted on slides in glycerine and coverslips were sealed around the edges with nail polish. Measurements were made after Morffe et al. (2009) and are given in millimetres. Variables are showed as range followed by median plus standard deviation and the number of measurements in parentheses. De Man's ratios a, b, c and V% were calculated.

Micrographs were taken with an AxioCam digital camera attached to a Carl Zeiss AxiosKop 2 Plus compound microscope. Line drawings were made with the softwares CorelDRAW X3 and Adobe Photoshop CS2 using micrographs as templates. Scales of all the plates are given in millimetres.

The materials examined are deposited in the CZACC and the Coleção Helminologica do Instituto Oswaldo Cruz (CHIOC), Rio de Janeiro, Brazil.

Systematics

Family Hystriognathidae Travassos, 1920

Genus *Coynema* gen. n.

urn:lsid:zoobank.org:act:3883651B-C4DC-4876-A7D2-1DBAA849E54C

Generic diagnosis. Female. Body robust, markedly fusiform. Cervical cuticle unarmed. Lateral alae present, from the esophageal region to a distance before the level of the vulva. Head bearing eight small, paired papillae. First cephalic annule slightly expanded, set-off from head by a deep single groove, its fore half with concave margins when totally relaxed. Esophagus consisting of a muscular sub-cylindrical procorpus, its base abruptly expanded in its joint with the short isthmus. Intestine simple, its fore region very expanded, forming a sac-like structure. The end of this region of the intestine is abruptly set-off and next to it the gut continues as a simple, rectilinear tube. Nerve ring encircling procorpus at its posterior half. Excretory pore post-bulbar. Reproductive system didelphic-amphidelphic. Eggs large, ovoid, smooth-shelled. Tail long, subulate and filiform.

Male. Body shorter and more slender than female. Cervical cuticle without spines. Head similar to female. First cephalic annule inconspicuous. Digestive system similar to female. Nerve ring encircling procorpus at its posterior half. Excretory pore post-bulbar. One testis present. Spicules absent. Tail conical, very short, sharply pointed, ventrally curved, its dorsal cuticle thickened forming a Y-like structure. A single, median, mammiform, big pre-cloacal papilla present. A pair of small, sub-dorsal, pre-cloacal papillae situated before the dorsal thickening.

Type species. *Coynema poeyi* (Coy, García & Alvarez, 1993) Morffe & García, comb. n. (monotypic genus).

Distribution. Cuba.

Etymology. The generic name is a combination of Coy, after Alberto Coy Otero, eminent Cuban parasitologist and the suffix -nema. The name is in neuter gender.

***Coyinema poeyi* (Coy, García & Alvarez, 1993) Morffe & García, comb. n.**

Figure 1 A–H, Figure 2 A–C, Figure 3 A–H

Glaber poeyi Coy, García & Alvarez, 1993: 57–59, fig. 3 A–E

Type material. ♂ holotype of *Glaber poeyi*, Cuba, Pinar del Río Province, Sierra del Rosario, El Salón; in *Passalus interstitialis*; IV.1990; A. Coy & N. García coll.; CZACC 11.4168. ♀ allotype of *Glaber poeyi*, same data as holotype, CZACC 11.4169.

Other material examined. 5 ♀♀, Cuba, Pinar del Río Province, Sierra del Rosario, El Mulo; in *Passalus pertyi*; X.2007; R. Núñez & O. Madruga coll.; CZACC 11.4468–11.4472. 9 ♀♀, Cuba, La Habana Province, San José de las Lajas, La Jaula; in *Passalus pertyi*; 15.III.2008; E. Fonseca, J. Morffe, G. León & F. Alvarez coll.; CZACC 11.4611–11.4619. 2 ♀♀, same data as anterior, deposited in the CHIOC. 6 ♀♀, Cuba, Holguín Province, Nipe-Sagua-Baracoa, La Melba; in *Passalus pertyi*; V.2007; R. Barba & D. Ortiz coll.; CZACC 11.4460–11.4465. 4 ♂♂, same data as anterior; CZACC 11.4473–11.4476.

Type host. *Passalus interstitialis* Escholtz, 1829 (Coleoptera, Passalidae).

Other host. *Passalus pertyi* Kaup, 1869 (Coleoptera, Passalidae).

Site. Gut caeca.

Type locality. El Salón, Sierra del Rosario, Pinar del Río Province, Cuba.

Other records. El Mulo, Sierra del Rosario, Pinar del Río Province, Cuba; La Jaula, San José de las Lajas, La Habana Province, Cuba; La Melba, Nipe-Sagua-Baracoa, Holguín Province, Cuba.

Measurements. Holotype (male) a = 14.40, b = 5.74, c = 83.08, total length = 1.080, maximum body width = 0.075, stoma length = 0.018, procorpus length = 0.130, isthmus length = 0.023, diameter of basal bulb = 0.038, total length of esophagus = 0.188, nerve ring to anterior end = 0.098, excretory pore to anterior end = 0.260, cloaca to posterior end = 0.013.

Allotype (female) a = 16.69, b = 9.15, c = 4.41, V% = 52.63, total length = 2.470, maximum body width = 0.148, first cephalic annule (length×width) = 0.013×0.035, stoma length = 0.025, procorpus length = 0.200, isthmus length = 0.025, diameter of basal bulb = 0.065, total length of esophagus = 0.270, nerve ring to anterior end = 0.130, excretory pore to anterior end = 0.370, vulva to posterior end = 1.170, anus to posterior end = 0.560, eggs = 0.113–0.118×0.048–0.050 (0.116 ± 0.004×0.049 ± 0.001 n = 2).

Population from El Mulo, Pinar del Río Province. Females (n = 5) a = 17.00–19.72 (18.25 ± 1.32 n = 4), b = 7.73–8.65 (8.18 ± 0.38 n = 4), c = 3.65–3.99 (3.82 ± 0.17 n = 4), V% = 46.99–52.94 (49.50 ± 2.84 n = 4), total length = 1.700–2.075 (1.863 ± 0.164 n = 4), maximum body width = 0.090–0.120 (0.102 ± 0.011 n = 5), first cephalic annule (length×width) = 0.008–0.013×0.023–0.030 (0.010 ± 0.002×0.026 ± 0.004), stoma length = 0.018–0.020 (0.019 ± 0.001 n = 5), procorpus length = 0.160–0.175 (0.166 ± 0.006 n = 5), isthmus length = 0.023–0.025 (0.023 ± 0.001 n = 5), diameter of basal bulb = 0.040–0.045 (0.043 ± 0.002 n = 5), total




Figure 1. *Coyinema poeyi* (Coy, García & Alvarez, 1993) Morffe & García comb. n. Female. **A** Esophageal region, lateral view **B** Cephalic end, internal view **C** Cephalic end, external view **D** Tail, lateral view **E** Vulva, lateral view **F** Egg **G** Reproductive system **H** Entire nematode, lateral view.

length of esophagus = 0.220–0.240 (0.226 ± 0.009 $n = 5$), nerve ring to anterior end = 0.108–0.128 (0.119 ± 0.009 $n = 5$), excretory pore to anterior end = 0.340 (0.340 $n = 1$), vulva to posterior end = 0.800–1.100 (0.944 ± 0.133 $n = 4$), anus to posterior end = 0.430–0.520 (0.484 ± 0.038 $n = 5$), eggs = 0.105–0.113 \times 0.035–0.048 ($0.110 \pm 0.003 \times 0.040 \pm 0.003$ $n = 11$).




Figure 2. *Coynema poeyi* (Coy, García & Alvarez, 1993) Morffe & García comb. n. Male. **A** Esophageal region, lateral view **B** Tail end, lateral view **C** Entire nematode, lateral view.

Population from La Jaula, La Habana Province. Females ($n = 11$) $a = 13.13$ – 20.00 (16.26 ± 1.95 $n = 11$), $b = 7.65$ – 9.33 (8.35 ± 0.53 $n = 11$), $c = 3.93$ – 4.49 (4.22 ± 0.19 $n = 10$), $V\% = 47.72$ – 51.92 (50.44 ± 1.07 $n = 11$), total length = 1.740 – 2.425 (2.010 ± 0.211 $n = 11$), maximum body width = 0.108 – 0.150 (0.125 ± 0.014 $n = 11$), first cephalic annule (length \times width) = 0.005 – 0.008×0.028 – 0.033 ($0.007 \pm 0.001 \times 0.029 \pm 0.002$ $n = 7$), stoma length = 0.020 (0.020 $n = 11$), procarpus length = 0.163 – 0.185 (0.174 ± 0.008 $n = 11$), isthmus length = 0.020 – 0.028 (0.023 ± 0.003 $n = 11$), diameter of basal bulb = 0.043 – 0.058 (0.049 ± 0.005 $n = 11$), total length of esophagus = 0.228 – 0.260 (0.240 ± 0.010 $n = 11$), nerve ring to anterior end = 0.120 – 0.140 (0.130 ± 0.006 $n = 11$), excretory pore to anterior end = 0.280 – 0.360 (0.320 ± 0.029 $n = 11$), vulva to posterior end = 0.860 – 1.200 (0.996 ± 0.107 $n = 11$), anus




Figure 3. *Coynema poeyi* (Coy, García & Alvarez, 1993) Morffe & García, comb. n. Female. **A** Cephalic end, relaxed first cephalic annule (specimen killed inside the host) **B** Cephalic end, first cephalic annule not relaxed (heat killed specimen) **C** Sac-like structure of the intestine **D** Esophagus **E** Egg **F** Vulva, lateral view. Male **G** Tail end showing the dorsal cuticular Y-like thickening **H** Esophageal region. Scale bars: **A, B, G** 0.025 mm. **D, E, F** 0.05 mm. **C, H** 0.1 mm.

to posterior end = 0.400–0.560 (0.478 ± 0.061 n = 10), eggs = 0.108–0.125×0.035–0.048 ($0.116 \pm 0.004 \times 0.042 \pm 0.004$ n = 20).

Population from La Melba, Holguín Province. Females (n = 6) a = 13.27–16.00 (14.80 ± 0.95 n = 6), b = 6.77–8.38 (7.53 ± 0.57 n = 6), c = 3.35–3.84 (3.60 ± 0.20 n = 6), V% = 43.18–48.24 (44.92 ± 2.27 n = 4), total length = 1.460–1.800 (1.653 ± 0.126 n = 6), maximum body width = 0.110–0.120 (0.112 ± 0.004 n = 6), first cephalic annule (length×width) = 0.010×0.025–0.028 ($0.010 \times 0.027 \pm 0.001$ n = 5), stoma length = 0.018–0.020 (0.018 ± 0.001 n = 6), procorpus length = 0.153–0.173 (0.160 ± 0.008 n = 6), isthmus length = 0.018–0.200 (0.019 ± 0.001 n = 6), diameter of basal bulb = 0.043–0.065 (0.048 ± 0.009 n = 6), total length of esophagus = 0.200–0.240 (0.220 ± 0.014 n = 6), nerve ring to anterior end = 0.113–0.125 (0.119 ± 0.006 n = 6), excretory pore to anterior end = 0.270–0.360 (0.307 ± 0.047 n = 3), vulva to posterior end = 0.820–1.000 (0.894 ± 0.076 n = 4), anus to posterior end = 0.380–0.530 (0.462 ± 0.052 n = 6), eggs = 0.105–0.125×0.035–0.048 ($0.113 \pm 0.006 \times 0.043 \pm 0.004$ n = 11).

Males (n = 4) a = 10.84–13.18 (11.87 ± 1.05 n = 4), b = 5.62–6.03 (5.88 ± 0.18 n = 4), c = 55.00–74.67 (66.92 ± 8.39 n = 4), total length = 1.030–1.120 (1.073 ± 0.044 n = 4), maximum body width = 0.085–0.095 (0.091 ± 0.004 n = 4), stoma length = 0.015–0.020 (0.017 ± 0.002 n = 4), procorpus length = 0.125–0.300 (0.128 ± 0.002 n = 4), isthmus length = 0.015–0.023 (0.018 ± 0.004 n = 4), diameter of basal bulb = 0.040–0.048 (0.044 ± 0.003 n = 4), total length of esophagus = 0.175–0.188 (0.183 ± 0.005 n = 4), nerve ring to anterior end = 0.093–0.100 (0.096 ± 0.003 n = 4), excretory pore to anterior end = 0.240–0.250 (0.248 ± 0.005 n = 4), cloaca to posterior end = 0.015–0.020 (0.016 ± 0.003 n = 4).

Description. Female. Body comparatively robust, markedly fusiform, maximum width at level of the anterior part of intestine. Cervical cuticle unarmed, finely annulated (annule more conspicuous toward post-esophageal region). Sub-cuticular longitudinal striae present. Lateral alae from the end of procorpus or the beginning of its basal dilation to about half of a body width before the level of vulva. Head well developed, set-off from body by a single, deep groove and bearing eight small paired papillae. First cephalic annule slightly expanded, when totally stretched (in dead nematodes inside the hosts) consisting of an anterior half with concave margins (its diameter initially inferior to that of head and increasing gradually in the posterior direction) and a posterior half wider and with convex margins. In heat-relaxed specimens the cephalic annule appears to be less stretched and only the posterior part is visible. Stoma short, about two first cephalic annule lengths long, surrounded by an esophageal collar. Esophagus consists of a muscular, sub-cylindrical procorpus, base abruptly dilated in its joint with the short isthmus. Basal bulb pyriform, valve plate well developed. Intestine simple, sub-rectilinear, anterior portion notably dilated, forming a sac-like structure slightly longer than esophagus. In this part the external surface of intestine almost touching the body wall. The continuation of intestine is about one third of the diameter of the sac-like structure. Rectum short, anus not prominent. Nerve ring encircling procorpus at about 60% of its length. Excretory pore situated at about half of a body width pos-

terior to basal bulb. Genital tract didelphic-amphidelphic, both ovaries thin, reflexed. Anterior ovary commencing just behind the saccular structure and posterior ovary arising slightly more than a body width anterior to the level of anus. Vulva a median transverse slit near midbody or slightly displaced forward, lips more or less prominent. Vagina muscular, thin-walled, forwardly directed. Conduct next to the vagina forming a loop. Eggs comparatively large, markedly ovoid in shape, smooth-shelled. Tail comparatively long, filiform and subulate.

Male. Body shorter and slender than female. Cervical cuticle unarmed. Sub-cuticular longitudinal striae present. Cephalic end similar to female, except by the cephalic annule inconspicuous. Digestive system similar to female. Sac-like region of the intestine slightly larger than the esophagus. Rectum short and cloaca inconspicuous, not prominent. Nerve ring encircling procorpus in posterior half, at about 60% of its length. Excretory pore situated at about less than a body-width posterior to the basal bulb. Testis single, commencing just posterior to the sac-like structure of intestine. Tail conical, sharply pointed, very short and ventrally curved. Dorsal cuticle near the tail tip bearing a Y-like thickening, its inferior part posteriorly directed. A single, median, large, mammiform pre-cloacal papilla present. A pair of small, sub-dorsal, pre-cloacal papillae situated before the dorsal cuticular thickening. Spicules absent.

Discussion. *Coynema* gen. n. can be placed in Hystrignathidae by having males with the single median pre-cloacal papilla characteristic of the family, oesophagus with its anterior portion supported by cuticularized rods and elongated eggs (Adamson and Van Waerebeke 1992). The new genus has affinities with the Brazilian genus *Glaber* by having a similar arrangement and form of the first cephalic annule and the characteristic basal dilation of the procorpus. A similar procorpus is present in *Vulcanonema* Travassos & Kloss, 1958, that however differs in its first cephalic annule separated from head by a conical region. These last two genera can be differentiated from *Coynema* gen. n. in having a monodelphic-prodelphic reproductive system. The marked fusiform shape of *Coynema* gen. n. only appears in the monotypic Malagasian genus *Pas-salidophila* Van Waerebeke, 1973, which is also monodelphic.

There are eight hystrignathid genera that are digonant and lack spines in the cervical cuticle: *Anomalostoma* Cordeira, 1981; *Anuronema*, *Klossnema* Cordeira & Artigas, 1983; *Papillabrum* Cordeira, 1981; *Phalacronema*, *Sprentia*, *Triumphalisenema* Kloss, 1962 and *Ventelia* Kloss, 1962. *Coynema* gen. n. differs from all of them in the form of cephalic end, the esophagus and the sac-like structure of the intestine. At present, the latter feature appears to be unique in the family.

The male of *Coynema* gen. n. presents the head and digestive system very similar to female. This is unusual in the few genera of Hystrignathidae having males described, which have a cylindrical procorpus. Due to this, the shape of the digestive system differentiates the males of *Coynema* gen. n. from the other males of Hystrignathidae.

Comments. There are small metric differences in the average values among females from some of the populations studied. In spite of this measurements tend to overlap, and there were no evident morphological differences observed. Males from El Salón and La Melba did not show marked morphometric variation. In addition, no male

specimens were found in the populations from El Mulo and La Jaula, when looking for more features that would support the existence of other species of *Coynema* gen. n. These are the reasons why all populations are considered conspecific until larger series of specimens are at hand for further study.

Acknowledgements

We are indebted to our colleagues Elier Fonseca, Gunnary León (Universidad de La Habana), Ormaily Madruga (Museo Nacional de Historia Natural), Rayner Núñez, René Barba (Instituto de Ecología y Sistemática), the undergraduate student Facundo Alvarez and David Ortiz for their help during the field work. To the inhabitants of La Jaula community (La Habana Province) for their hospitality and support. To MSc. Yamir Torres (Instituto de Ecología y Sistemática) and Dr. Alejandro Barro (Universidad de La Habana) for their help with the micrographs. We also thank Dr. Luis F. de Armas from the Instituto de Ecología y Sistemática for his critical review of the manuscript and Dr. Pedro Herrera for the revision of the English manuscript. This work was financed by IDEAWILD and the project “Colecciones Zoológicas, su Conservación y Manejo” (DB-06) of the Ministerio de Ciencia, Tecnología y Medio Ambiente, Cuba.

References

- Adamson M, Van Waerebeke D (1992) Revision of the Thelastomatoidea, Oxyurida of invertebrate hosts III. Hystrignathidae. *Systematic Parasitology* 22: 111–130.
- Clark WC (1978) Nematoda (Oxyuroidea: Thelastomatidae) from Australian passalid beetle (Coleoptera: Passalidae) with description of new species. *Australian Journal of Parasitology* 26: 603–615.
- Cordeira N (1981) Nematóides intestinais de Passalidae; revisao de sua sistemática e catalogação das espécies conhecidas. PhD thesis, Sao Paulo, Brazil: Institute of biomedical Sciences, University of Sao Paulo.
- Cordeira N, Artigas P (1983) Nematóides de Passalidae (Coleoptera). Descrição de *Klossnema repentina* n. gen., n. sp. (Nematoda: Hystrignathidae). *Memorias do Instituto Butantan* 47/48: 107–111.
- Coy A (1990) Nemátodos de la familia Hystrignathidae parásitos de coleópteros (Passalidae) de Cuba. *Poeyana* 402: 1–7.
- Coy A, García N (1995) Nuevas especies de nemátodos parásitos de insectos mexicanos. *Avicent* 12: 10–15.
- Coy A, García N, Alvarez M (1993) Nemátodos parásitos de insectos cubanos, Orthoptera (Blattidae y Blaberidae) y Coleoptera (Passalidae y Scarabaeidae). *Acta Biológica Venezolana* 14(3): 53–67.
- García N, Coy A (1994) Descripción de dos nuevas especies y registro de nuevos hospederos de nemátodos (Nematoda) de la región oriental de Cuba. *Avicennia* 1: 13–17.

- García N, Coy A (1995a) Nuevas especies de nemátodos (Nematoda) parásitos de artrópodos cubanos. *Avicennia* 3: 87–96.
- García N, Coy A (1995b) Nemátodos parásitos de la Sierra de los Organos, Cuba. *AvaCient* 14: 26–30.
- García N, Coy A (1997) Nueva especie y nuevo registro de nemátodos (Nematoda) parásitos de artrópodos mexicanos. *AvaCient* 20: 27–31.
- García N, Ventosa L, Morffe J (2009a) Dos especies nuevas de los géneros *Lepidonema* y *Longior* (Thelastomatoidea: Hystrignathidae) de la Isla de la Juventud, Cuba. *Solenodon* 8: 1–7.
- García N, Ventosa L, Morffe J (2009b) Especie nueva de *Artigasias* (Thelastomatoidea: Hystrignathidae) de Pico La Melba, Holguín, Cuba. *Solenodon* 8: 8–11.
- García N, Ventosa L, Morffe J (2009c) Nuevas especies de hystriagnátidos (Thelastomatoidea: Hystrignathidae) de la Sierra del Rosario, Pinar del Río, Cuba. *Novitates Caribaea* 2: 17–22.
- Guerrero R (1980) Descripción de cuatro especies nuevas de Thelastomatidae (Nematoda) endoparásitos de *Popilius* sp. (Coleoptera: Passalidae) de Los Andes venezolanos. *Memorias de la Sociedad de Ciencias Naturales (La Salle)* 40(114): 63–78.
- Hunt DJ (1981) On *Artigasias horridospina* n. sp., *Longior semialata* n. sp., *Paraxyo ensicrinatus* n. sp. (Oxyurida: Hystrignathidae) and *Pulchrocephala pulchrocephala* Travassos, 1925 (Oxyurida: Pulchrocephalidae). *Systematic Parasitology* 3: 33–52.
- Hunt DJ (1982) *Hystrignathus ferox* n. sp. and *Xyo xiphacanthus* n. sp. (Oxyurida: Hystrignathidae) with additional data on *Carlosia tijucana* Travassos and Kloss, 1957. *Systematic Parasitology* 4: 59–68.
- Kloss GR (1962) Alguns parasitos do intestino de Coleopteros da familia Passalidae. *Papeis Avulsos do Departamento de Zoologia de Sao Paulo* 15: 163–171.
- Morffe J, García N, Ventosa ML (2009) *Longior similis* sp. nov. (Thelastomatoidea: Hystrignathidae) parasite of *Passalus interstitialis* from Western Cuba and new records of *Longior zayasi*. *Solenodon* 8: 12–19.
- Morffe J, García N (2010a) A new species of *Artigasias* Christie, 1934 (Thelastomatoidea: Hystrignathidae) from Cuba, and redescription of *Artigasias simplicitas* García et Coy, 1995. *Novitates Caribaea* 3: 53–61.
- Morffe J, García N (2010b) Two new species of nematode (Oxyurida, Hystrignathidae) parasites of *Passalus interstitialis* Escholtz, 1829 (Coleoptera, Passalidae) from Cuba and a new locality for *Longior similis* Morffe, García et Coy, 2009. *ZooKeys* 49: 1–11.
- Travassos L, Kloss GR (1957a) Nématodeos de invertebrados. 1.^a nota. *Revista Brasileira de Biologia* 17(3): 295–302.
- Travassos L, Kloss GR (1957b) Nématodeos de invertebrados. 2.^a e 3.^a notas. *Revista Brasileira de Biologia* 17(4): 467–477.
- Travassos L, Kloss GR (1958) Sobre a fauna de Nematodeos dos Coleopteros-Passalidae da Estacao biologica de Boraceia. *Arquivos de Zoologia do Estado de Sao Paulo* 11: 23–57.