

Review

Critical Impact of Different Conserved Endoplasmic Retention Motifs and Dopamine Receptor Interacting Proteins (DRIPs) on Intracellular Localization and Trafficking of the D₂ Dopamine Receptor (D₂-R) Isoforms

Kaja Blagotinšek Cokan ^{1,†}, Maša Mavri ¹, Catrin Sian Rutland ², Sanja Glišić ³, Milan Senčanski ³, Milka Vrecl ¹ and Valentina Kubale ^{1,*}

¹ Department of Anatomy, Histology with Embryology and Cytology, Institute of Preclinical Sciences, Veterinary Faculty, University of Ljubljana, Gerbičeva 60, 1000 Ljubljana, Slovenia; kaja.blagotinsek@mf.uni-lj.si (K.B.C.); masa.mavri@vf.uni-lj.si (M.M.); milka.vrecl@vf.uni-lj.si (M.V.)

² School of Veterinary Medicine and Science, Medical Faculty, University of Nottingham, Sutton, Bonington Campus, Loughborough LE12 5RD, UK; Catrin.Rutland@nottingham.ac.uk

³ Center for Multidisciplinary Research, Institute of Nuclear Sciences VINCA, University of Belgrade, Mike Petrovića Alasa 12-14, 11351 Vinča, Belgrade, Serbia; sanja@vin.bg.ac.rs (S.G.); sencanski@vin.bg.ac.rs (M.S.)

* Correspondence: valentina.kubale@vf.uni-lj.si

† Centre for Functional Genomics and Bio-Chips, Institute of Biochemistry, Faculty of Medicine, University of Ljubljana, 1000 Ljubljana, Slovenia.

Received: 3 July 2020; Accepted: 18 September 2020; Published: 23 September 2020



Abstract: The type 2 dopamine receptor D₂ (D₂-R), member of the G protein-coupled receptor (GPCR) superfamily, exists in two isoforms, short (D_{2S}-R) and long (D_{2L}-R). They differ by an additional 29 amino acids (AA) in the third cytoplasmic loop (ICL3) of the D_{2L}-R. These isoforms differ in their intracellular localization and trafficking functionality, as D_{2L}-R possesses a larger intracellular pool, mostly in the endoplasmic reticulum (ER). This review focuses on the evolutionarily conserved motifs in the ICL3 of the D₂-R and proteins interacting with the ICL3 of both isoforms, specifically with the 29 AA insert. These motifs might be involved in D₂-R exit from the ER and have an impact on cell-surface and intracellular localization and, therefore, also play a role in the function of dopamine receptor signaling, ligand binding and possible homo/heterodimerization. Our recent bioinformatic data on potential new interaction partners for the ICL3 of D₂-Rs are also presented. Both are highly relevant, and have clinical impacts on the pathophysiology of several diseases such as Parkinson's disease, schizophrenia, Tourette's syndrome, Huntington's disease, manic depression, and others, as they are connected to a variety of essential motifs and differences in communication with interaction partners.

Keywords: D₂ dopamine receptor; intracellular trafficking; endoplasmic reticulum; retention motifs; ICL3; interacting partners

1. Dopamine Receptors

G-protein-coupled receptors (GPCRs), also termed seven-transmembrane receptors (7TMRs) are by far the largest family of membrane-bound receptors, which are involved in the regulation of the neurotransmitter dopamine effects as one of their targets [1]. Based on functional, structural, and pharmacological properties, five types of dopamine receptors have been described, that belong to

the D₁- or D₂-like subfamily of receptors (D₁-R and D₂-R respectively), with differing abilities of stimulation or inhibition of adenylyl cyclase (AC), respectively.

The D₁-R subfamily is comprised of D₁ and D₅ receptors (D₁-R and D₅-R), and the D₂ subfamily includes D₂, D₃ and D₄ receptors (D₂-R, D₃-R, and D₄-R). Members of the D₁-R subfamily have a short third cytoplasmic loop (ICL3) and a very long C-terminal cytoplasmic end. In contrast, D₂-Rs have a very long ICL3 and a short C-terminal end and include the receptor variants generated by alternative splicing (D₂ and D₃) or polymorphic variation (D₄) (reviewed by Beaulieu et al.) [2]. The D₂-R subgroup has a long ICL3 whose structure is common to the receptor interaction with the heterotrimeric protein G α_i [1]. In D₁-R, with its characteristically short ICL3, coupling with G α_s proteins occurs [1,3]. The C-terminal end is approximately seven times longer in D₁-Rs than in D₂-Rs. Both the ICL3 and the C-terminal end are thought to serve as possible communication points for interaction with intracellular proteins. The N-terminal tail has a similar number of amino acids in all receptor subtypes and contains sites for N-glycosylation. D₁-R and D₅-R have two glycosylation sites, located at the N-terminal end and in extracellular loop 2 (ECL2). D₂-R has three potential N-linked glycosylation sites, all in the N-terminus: N5, N17, and N23, D₃-R has four potential glycosylation sites: N12 and N19 in the N-terminus, N97 in the first extracellular loop (ECL1), and N173 in the second extracellular loop (ECL2) [4] and D₄-R only one in the N-terminus [5]. Cysteines located in the first and second extracellular loops (ECL1 and ECL2) are linked by a disulfide bond that stabilizes the receptor structure [6]. The endogenous ligand for the dopamine receptors is the neurotransmitter dopamine. After dopamine binds to the D₁-R, the signaling pathway is canonically activated via the heterotrimeric protein G α_s and G α_{olf} G-proteins, leading to adenylyl cyclase (AC) activation and cyclic adenosine monophosphate (cAMP) formation in the cell. Diversity in functional outcomes may also be achieved via selective binding to G α_i and G α_o proteins. Previous work has shown that D₂-R can be stabilized by an agonist, which affect the selectivity and amount of coupling with G α_i and G α_o [7,8]. Although previous work had indicated that G α_{i2} was selective for D_{2L}-R [9,10], experimental data has indicated that selectivity regulation of G α_i is driven by the agonist-activated conformation of D₂-R. R(+)-3-PPP hydrochloride stimulation of D₂-R resulted in reduced coupling with G α_{i1} or G α_{i2} and preferential coupling with G α_{i3} [11]. The movement magnitude of the sixth transmembrane helix of the activated receptor was predicted to be the primary modulator of the selectivity of the G-protein subtypes [12].

Using cryo-electron microscopy, the structure of an agonist-bound activated D₂-G α_i complex reconstituted into a phospholipid membrane has been demonstrated recently [13], both as the first experimental model of a GPCR complex embedded in a phospholipid bilayer, as well as the first model of activated D₂-R. The models revealed interactions that are unique to the membrane-embedded complex, such as conformational changes in ECL2, TM5, TM6 and TM7, propagating to the opening of the intracellular G α_i -binding site and helix 8 burial in the inner leaflet, ordered lysine and arginine side chains in the membrane interfacial regions, and lipid anchoring of the G-protein in the membrane [13].

Although all D-Rs recognize the same ligand, they have a differential tissue distribution and are involved in different functions in vivo [3,14]. By binding to various types of D-Rs, dopamine controls locomotor system functions, cognition, emotion, hunger, satiety, and endocrine secretion [3,5]. Impaired D₂-R signaling is associated with the pathophysiology of many psychiatric and neurological diseases or states, including Parkinson's disease, schizophrenia, Tourette's syndrome, Huntington's disease, bipolar disorder, depression, dementia, as well as others, such as restless leg syndrome and sexual dysfunction. D-Rs are an essential target for currently available modern drugs, including the dopamine precursor levodopa [3,5] for Parkinson's disease, where dopaminergic neurons are damaged and a dopamine deficiency leads to a combination of movement and psychiatric pathologies. Thus, D-Rs are targets for motor deficits, cognitive, and motivational deficits in neuropsychiatric disorders [15]. In schizophrenia and psychosis inhibitors of D₂-R are used to reduce increased dopaminergic signaling [16].

2. Dopamine Receptor Type 2 (D₂-R)

The D₂-R is a key component of the dopamine system that is present in two alternatively spliced transcripts of the *Drd2* gene and classified as short (D_{2S}-R) and long (D_{2L}-R) receptor isoforms. The long isoform differs from the short one only by the presence of an additional 29 amino acids (AA) encoded by exon 5 in the ICL3 of the D_{2L}-R [17–19]. The inclusion is interspersed between the AA lysine (K241) and glutamic acid (E271). D_{2S}-R in mice and rats are made up of 415 AA and D_{2L}-R is made up of 444 AAs. Human D_{2S}-R and D_{2L}-R are shorter than murine and rat equivalents by one AA, consisting of 414 and 443 AAs, respectively. The isoleucine is missing between lysine (K331) and aspartic acid (D332). This region might have an essential role in the functional differences between both D₂-R isoforms such as interactions related to G-proteins [20–22], post-translation modification and cell localization [11,23]. D₂-R isoforms also indicate different *in vivo* functions, whereby D_{2L}-R primarily acts at postsynaptic and D_{2S}-R in presynaptic dopaminergic transmissions [24,25]. Data acquired on genetically engineered D₂-R mouse model indicates additional evidence for different roles of two isoforms in cognitive and motor functions [24], responsiveness to cocaine exposure [26], and therapeutic effects of antipsychotic drugs [27]. Furthermore, they are expressed in the same cell types with more abundant expression of the D_{2L}-R isoform over D_{2S}-R, but with differences in their intracellular localization. While D_{2S}-R is primarily localized on the plasma membrane (PM), a substantial fraction of D_{2L}-R is located intracellularly, especially in the perinuclear compartments around the Golgi apparatus (GA) [14] and endoplasmic reticulum (ER) [23].

The D₂-R is the most commonly studied dopamine receptor subtype since the majority of antipsychotic drugs act as D₂-R antagonists in the mesolimbic dopaminergic system [28]. As a primary target for atypical and typical antipsychotic drugs and treatment of the Parkinson's disease, many of those agents can cause potentially life-threatening and severe side effects due to the promiscuous activities against related D₂-Rs [29]. Precisely because of this reason, it is necessary to be familiar with the details of the dopamine receptor's complex structure and functions.

3. Localization Differences Between D_{2S}-R and D_{2L}-R

D₂-R isoforms localization is neither species nor tissue specific. They were found in different tissues, but in highly variable ratios [25,30,31]. D_{2S}-R is predominantly localized in the PM [14,22,32], whereas an intracellular D_{2L}-R reservoir has been reported in the primates brain [33] and several cell lines [14,22,23].

The primary intracellular localization of the D_{2L}-R in transiently transfected HEK-293, COS-7, and HeLa is the ER [23], whereas in transfected NG108-15 cells intracellularly localized D_{2L}-R is predominantly co-localized with the GA matrix protein marker GM130 [14]. At the level of confocal microscopy, higher proportions of D_{2L}-R than D_{2S}-R were retained intracellularly in heterologous cell lines [22,23]. This finding could be due to the retention of overexpressed or incorrectly folded tagged receptors in the ER. However, immunoelectron microscopy also revealed the predominant intracellular localization of the D_{2L}-R [34] in monkey dopaminergic neurons, and that these sites are cisterns of GA and ER. Ligand-promoted recruitment of the D_{2L}-R [35], D₄-R [36] and other GPCRs, such as thrombin receptors (PAR1 and PAR2), D₁-Rs, and opioid receptors on the PM presented additional evidence for the existence of functional, pre-existing intracellular stores (reviewed by Achour, 2008) [37].

N-terminal glycosylation of different GPCR's has a vital role in cell surface receptor expression. Mutations of potential N-terminal glycosylation sites found for D₂-R (N5, N17, and N23) lead to decreased surface expression for D₂-R, showing their important role in receptor distribution [38]. Fishburn et al. performed studies regarding post-translational processing of the D_{2L}-R and D_{2S}-R isoforms [39]. Three post-translational states were observed in both receptor isoforms: a newly synthesized protein (35 kDa), a partially glycosylated product (45 kDa) and a fully glycosylated receptor (70 kDa) [39].

A difference in the processing of the mature receptor was observed. The initial N-glycosylation of the newly synthesized receptor protein occurred shortly after synthesis in both D₂-R isoforms,

suggesting that a rapid and efficient maturation towards partially glycosylated product occurs in D₂-R. However, a marked difference was observed in subsequent N-linked glycosylation, where D_{2L}-R showed slower production of fully glycosylated proteins in comparison to the D_{2S}-R. Additionally, they showed that 20% of the D_{2L}-R remains in the partially processed form and never undergoes the second stage of N-linked glycosylation.

4. Functional Differences between D_{2S}-R and D_{2L}-R

Identifying probable functional differences between D_{2S}-R and D_{2L}-R has been the subject of numerous studies. Depending on the site of action and the effect on D₂-R-mediated responses, isoforms have different and likely antagonistic functions *in vivo* [24]. D_{2S}-R is mainly a presynaptic receptor, but at the postsynaptic level, it negatively modulates D₁-R-dependent responses. In contrast, D_{2L}-R is found predominantly at postsynaptic sites where it acts synergistically with D₁-Rs [24].

The location of the inclusion in ICL3 also led to the assumption that it may affect the specificity of the interaction with G-proteins and the sequential activation of specific effector proteins.

Several studies have shown that structural differences between isoforms can determine the specificity of interactions with G-proteins [10,40]. A recent study showed the preferential coupling of both D₂-R isoforms with G-proteins (G_{i1} and G_{i2}), due to the differences in ICL3, which affect receptor behavior [11,41]. Results obtained with the messenger gene construct controlled by the cAMP response promoter suggest constitutive, i.e., agonist-independent D_{2L}-R activity [42]. Studies with D₂-R knockout mice provided additional evidence for their diverse roles in motor and cognitive functions [43], sensitivity to cocaine [44], and therapeutic/side effects of antipsychotic agents [45].

5. The ER Retention Motifs in GPCRs and Both D₂-R Isoforms

The synthesis and transfer of newly formed proteins via secretory pathways from the ER to the PM is a complex process involving different mechanisms and many additional proteins and motifs that are important in protein interaction and enable their proper formation, quality control, selective retention, and transport [46]. Mechanisms that regulate the secretory transport of GPCRs or their transfer to the cell surface are poorly elucidated [47,48]. It is known that the interconnection of the same or different GPCRs, homo- and heterodimerization, is essential in the transport of GPCRs in families A and C, but not for representatives of family B on the PM [49]. Transfer of the protein to the PM requires control of transport from the ER via the GA to the PM. This process is regulated by COPI and COPII vesicles [50]. COPII-coated transfer vesicles serve anterograde transport from the ER to GA, whilst transport between GA cisternae and retrograde transport from GA to ER takes place with COPI-coated vesicles. Improperly synthesized proteins or those having exposed sequences encoding motives for retention in the ER are transported retrogradely into the ER by COPI-coated vesicles [51]. Moreover, correctly folded proteins might be retained in the ER because they hold ER retention motifs, which prevent their export from the ER. Three types of ER retention motifs have been identified in the intracellular domains of various proteins: KDEL, KKXX, and RXR type motifs [46]. The presence of specific conserved sequences, so-called ER retention signals, could be responsible for preventing D₂-R proteins from leaving the ER. Since the level of GPCR expression dictates the magnitude of cellular responses elicited by a signal at the PM, which is the balance of elaborately regulated endocytic and exocytic trafficking, it is crucial to know the motifs as well as the proteins involved in this interplay.

6. KDEL and KKXX Motifs

The KDEL motif is a short C-terminal retrieval signal (Lys-Asp-Glu-Leu) identified in ER luminal chaperone proteins, such as immunoglobulin heavy chain-binding protein (BiP) and other soluble ER resident proteins [20,52,53]. The KDEL receptor recognizes this motif in the post-ER compartments, which mediates retrograde transport to the ER by COPI coatomer structures [54]. For the proper sorting of cargo into COPI vesicles, a Ras-like small GTPase ADP-ribosylation factor 1 (ARF1) activation is required [55]. ARFGAP1 activates ARF1 by hydrolysis of GTP to GDP [56]. Ligand binding on the

luminal side of the KDEL receptor induces interaction with ARFGTP1 on the cytoplasmic side of the receptor, resulting in the recruitment of ARFGTP1 from the cytosol to the PM leading to ARF1 activation [57].

In zebrafish D₂-Rs and D₃-Rs, the expression of gene *Hsp47* was identified, which is an ER-resident collagen-specific chaperone with a C-terminal KDEL retention motif and plays a fundamental role in the folding, stability, and intracellular transport of procollagen triple helices [58].

The KDEL receptor cycles between the ER and the GA and its affinity for KDEL containing proteins changes between these two compartments. Retrieval of proteins mediated by the KDEL receptor can occur from different sites, ranging from early Golgi complex locations to trans Golgi networks [30]. In the GA, the KDEL receptor could associate with G α_o , one of the abundant G α subunit and regulate receptor trafficking through G-proteins [59].

In addition to KDEL, the di-lysine motif (KKXX), have been identified as retrieval signal, important for recycling proteins from the GA back to the ER [60]. These signals are necessary for determining the localization of modified secretory and PM proteins in the ER [30].

Type I integral membrane proteins, ERGIC53, and p24 family proteins contain di-lysine KKXX motifs [46]. This carboxyl-terminal retrieval signal usually consists of two lysine residues on positions -3 and -4 relative to the C-terminus, followed by any amino acid [61]. The KKXX signal is evolutionarily conserved as it also appears in yeast [20]. Like KDEL, the KKXX motif also serves as a retrieval signal for the transport of proteins from GA by COPI vesicles, although it binds directly with coatamer structures and does not require a receptor [62].

7. RXR and RSRR Motifs

The RXR motif, and in some proteins, RSRR, has been found on different proteins where it disables the exit of proteins from the ER. Initially, they were found in ion channels and also in several GPCRs [46,63]. The first discovered of two arginine retention signals (the RXR and RSRR) are known to be located at the C-terminal end of the Kir 6.2 potassium channel (RXR signal) and a GPCR representative, the gamma-aminobutyric acid type B1 receptor (GABA_{B1}; RSRR signal) [27]. The GABA_{B1} receptor contains the C-terminal RXR type ER retention motif RSRR, which prevents protein release from the ER. GABA_{B1} is functionally impaired in terms of ligand binding when expressed alone, whereas GABA_{B2} is nonfunctional in its signaling properties. Only when co-expressed with GABA_{B2}, GABA_{B1} receptor releases from the ER and translocates to the cell surface [27,64]. Upon co-expression, the RSRR retention signal of GABA_{B1} is proposed to be masked by interaction of the C-terminus of both subunits due to highly stable α coil-coil interactions. Therefore, GABA_{B1} and GABA_B are functionally combined of distinct subunits as obligatory constitutive heterodimers [27,64–66].

There are more examples of the RXR-type retention motif in GPCR receptors. A published study showed that in the C-terminal end of the type 2c α -adrenergic receptor (α 2C-AR), there is a set of five arginine residues (RRRRR), which represent a possible retention signal of the ER type RXR [67]. The RXR-type retention motif has also been described in the ICL3 of the kainate receptor, which is a ligand-dependent ion channel [68]. Disease-causing vasopressin type 2 receptors (V2R) mutations are retained in different compartments of the early secretory pathway [69]. V2R mutants connected to nephrogenic diabetes insipidus in the contrast with the wild-type V2R are less expressed on the cell surface. Additionally, the D₂-R RXR motifs have been revealed. In our study [70], we showed that the evolutionarily conserved arginine cluster in the insert of ICL3 of D_{2L}-R (R267-R269) acts as an ER retention signal and is potentially crucial for anterograde trafficking of the D₂-R and receptor PM availability. However, we must take into account that ER exit is a highly regulated process and that one motif is not responsible solely for it. Other proposed mechanisms involving interaction with the ER-resident gatekeeper prenylated Rab acceptor 1 domain family member 3 (PRAF3) or other D_{2L}-R binding and interaction proteins, such as fatty acid-binding protein 3 (FABP3), which possibly bind to 29 AA in ICL3 in D_{2L}-R may be included (Figure 1) [71].

Additionally, the mutational analysis revealed that heteromers between the dopamine D₂-R, adenosine A_{2A} and cannabinoid CB₁ receptors are stabilized by electrostatic interactions between arginine-rich motifs in the ICL3 of D₂-R and A_{2A} receptors and phosphorylated casein kinase 1/2 sites in ICL3 and C-tail of the CB1 receptor, and the C-terminus of the A_{2A} receptor [72]. The RXR-rich motif in ICL3 of the D₂-R is also involved in stabilizing electrostatic interactions with a di-glutamate motif in the C-terminus of the serotonin 5-HT_{2A} and D₁-R [73,74].




Figure 1. Motif and interaction partners' differences between D_{2L}-R and D_{2S}-R. CaM—Ca²⁺-binding protein calmodulin; CGAT—Chromaffin granule amine transporter; DAT—dopamine transporter; FABP3—Fatty acid binding protein 3; FLN-A—filamin A; NMDA (NR2B)—NR2B subunit of the NMDA glutamate (N-methyl-D-aspartate); NSF—N-ethylmaleimide-sensitive factor; Par-4—Prostate apoptosis response-4; PM—plasma membrane; PREB—prolactin regulatory element-binding protein; Rabex-5—Rabaptin-5 interacting protein; TRAPPC9—Trafficking protein particle complex subunit 9.

Likhite, N. et al. [75] used bioinformatics analysis to identify 583 RGG and RXR-type motifs in GPCRs. Approximately 34% of those were conserved in human GPCRs within the ICL3 and could serve as arginine methylation motifs. They showed that R217 and R219 within the ICL3 N-terminal end common to both D₂-R isoforms serve as an arginine N-methyltransferase 5 (PRMT5) methylation motif important for modulating receptor signaling [75]. This corroborates with R217 and R219 location of within the G α interaction domain (reviewed in [75]).

Furthermore, Table 1 summarizes selected motifs from the eukaryotic linear motif (ELM) [76] resources, which are potentially involved in ER-GA trafficking of the D₂-R. The experimentally validated short linear motifs (SLiM) were manually curated after globular domain filtering, structural filtering, and context filtering.

Table 1. Motifs obtained from eukaryotic linear motif (ELM) bioinformatics analysis of type 2 dopamine receptor D₂ (D₂-R) sequences.

ELM Name	Instances (Matched Sequence)	AA Position	ELM Description	Cell Compartment	Pattern	Probability (×10 ³)
TRG_ER_diArg_1	RRRR RRR RRKR RKRK RRR RRV	217–220 218–220 219–222 220–223 267–269 268–270	The di-Arg ER retention motif is defined by two consecutive RR residues or with a single residue insertion (RXR). The motif is completed by an adjacent hydrophobic/arginine residue, which may be on either side of the R pair.	ER membrane, integral protein, ER-GA transport vesicle membrane, ER membrane, GA-ER transport vesicle membrane, rough ER, ER	([LIVMFYWPR]R[^Y FWDE]{0,1}R)(R[^Y FWDE]{0,1}R[LIVM FYWPR])	5.37
CLV_NRD_NRD_1	RRK RRV RRA RRK FRK	219–221 268–270 274–276 360–362 433–435	N-Arg dibasic convertase (NRD/Nardilysin) cleavage site (X- -R-K or R- -R-X).	extracellular, GA, cell surface	(.RK) (RR[^Y KR])	7.47
CLV_PCSK_FUR_1	RRRK RRKRV	217–221 219–223	Furin (PACE) cleavage site (R-X-[RK]-R- -X).	extracellular, GA, GA membrane	R.[RK]R.	50.09
CLV_PCSK_KEX2_1	KRR RRR RRR RRK KRV KRS RRR RRV RRA RRK	149–151 217–219 218–220 219–221 221–223 226–228 267–269 268–270 274–276 360–362	Yeast kexin 2 cleavage site (K-R- -X or R-R- -X).	extracellular, GA	[KR]R.	7.97
CLV_PCSK_PC1ET2_1	KRR KRV KRS	149–151 221–223 226–228	NEC1/NEC2 cleavage site (K-R- -X).	extracellular, GA, GA membrane	KR.	3.90
CLV_PCSK_PC7_1	RYSSKRR RRRRKRV RVNTRKS	145–151 217–223 222–228	Proprotein convertase 7 (PC7, PCSK7) cleavage site (R-X-X-X-[RK]-R- -X).	extracellular, GA, GA membrane	R...[KR]R.	50.09

Table 1. Cont.

ELM Name	Instances (Matched Sequence)	AA Position	ELM Description	Cell Compartment	Pattern	Probability ($\times 10^3$)
CLV_PCSK_SKI1_1	KIAKI KKATQ RKAFL KAFLK	336–340 369–373 434–438 435–439	Subtilisin/kexin isozyme-1 (SKI1) cleavage site ([RK]-X-[hydrophobic]-[LTKF]-X).	ER lumen, ER GA, extracellular	[RK].[AILMFV][LTKF].	6.82
LIG_deltaCOP1_diTrp_1	EWKF	99–105	Tryptophan-based motifs enable targeting of the tethering and (dis)assembly factors to the C-terminal mu homology domain (MHD) of the coatmer subunit delta, delta-COP.	ER membrane GA membrane, COPI coated vesicle membrane, cytosol, COPI vesicle coat, transport vesicle	[DE]{1,3}.{0,2}W.{1,6}[WF]	50.10
LIG_LIR_Gen_1	EWKFSRI	99–105	Canonical LIR motif that binds to Atg8 protein family members to mediate processes involved in autophagy.	cytosol, cytoplasmic side of late endosome membrane	[EDST].{0,2}[WFY] [^RKPG][^PG][ILV]	3.06
LIG_LIR_Nem_3	EWKFSRI TRYSPI	99–105 293–298	Nematode-specific variant of the canonical LIR motif that binds to Atg8 protein family members to mediate processes involved in autophagy.	cytosol, cytoplasmic side of late endosome membrane	[EDST].{0,2}[WFY].[ILVfy]	6.36
MOD_N-GLC_1	EWKFSRI TRYSPI	99–105 293–298	Generic motif for N-glycosylation. It was shown that Trp, Asp, and Glu are uncommon before the Ser/Thr position. Efficient glycosylation usually occurs when ~60 residues or more separate the glycosylation acceptor site from the C-terminus.	extracellular, GA, ER	[EDST].{0,2}[WFY].[ILVfy]	5.02
MOD_N-GLC_2	NEC	180–182	Atypical motif for N-glycosylation site. Examples are Human CD69, which is uniquely glycosylated at typical (Asn-X-Ser/Thr) and atypical (Asn-X-Cys) motifs, beta protein C.	extracellular, GA, ER	(N)[^P]C	29.7
TRG_ENDOCYTTIC_2	YTAV YSPI	133–136 295–298	Tyrosine-based sorting signal responsible for the interaction with mu subunit of AP (Adaptor Protein) complex.	PM, clathrin-coated endocytic vesicle, cytosol	Y.[LMVIF]	2.59

Legend: GA—Golgi apparatus; ER—endoplasmic reticulum; PM—plasma membrane.

8. ER Export Motifs in GPCRs and D₂-R

GPCRs originate in the ER, where they are synthesized, folded, and assembled. Properly folded receptors are recruited and packaged into ER-derived COPII-coated vesicles. Transport vesicles carrying cargo receptors then migrate from the ER to the ER-Golgi intermediate complex (ERGIC), the GA, and the trans Golgi network (TGN). During their transport, receptors undergo post-translational modifications (e.g., glycosylation). Mature receptors then move from the TGN to their destination at the PM [77]. For this process, conserved sequences and motifs essential for the exit of GPCRs from the ER are critical. Export from the ER is the first step in the intracellular trafficking of GPCRs. These motifs are found on the C- or N-terminal tail of the receptors and are therefore common to both D₂-R isoforms. A triple phenylalanine motif [F(X)₃F(X)₃F] [14] has been identified in the membrane-proximal C-terminus of the D₁-R that is required for receptor cell-surface expression [78]. However, no motifs have yet been identified on the D₂-R. A newly identified ER-membrane-associated protein, DRIP78, binds to this motif, as described later.

9. D₂-R Interaction Proteins (DRIPs)

More than 20 dopamine receptor-interacting membrane-associated or cytoplasmic D₂-R interaction proteins (DRIPs) are known and several of them bind the ICL3 of the D₂-R [79]. Using the informational spectrum method (ISM), a virtual spectroscopy method for investigating protein-protein interactions, the analysis of known interaction partners of ICL3 of D₂-R [79] was performed as previously described [80,81] and obtained the results presented in Table 2. ISM analysis of the ICL3 D₂-R interaction with protein partners corroborates with published data (reviewed in [79]) (Table 2, Figure 1). However, in addition to previously identified protein partners it has also been suggested that there are some new potential interaction partners.

Among previously described interaction partners, the highest affinity for the interaction with the D₂-R was ascribed to *N*-methyl-D-aspartate (NMDA) receptor NR2B subunits. It was shown that a distinct region within the first 32 AA of the D₂-R ICL3 interacts with the NR2B and disrupts the association of Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) with NR2B, reduces NR2B phosphorylation at a CaMKII-sensitive site (Ser1303), and inhibits NMDA receptor-mediated currents in medium-sized striatal neurons. The D₂-R-NR2B interaction is therefore critical for modulating NMDA receptor-mediated currents and behavioral responsiveness to cocaine [82]. The second highest propensity for interaction with the D₂-R was observed for prostate apoptosis response-4 (Par-4). Par-4 is a protein expressed in the nervous system, where it is known to be a regulatory component in dopaminergic signaling. It is a mediator of neuronal degeneration, and is associated with the pathogenesis of Alzheimer's disease [83]. Par-4 directly interacts with the D₂-R via the calmodulin-binding motif in the ICL3. Furthermore, Par-4 constitutes a molecular link between impaired dopaminergic signaling and depression [84]. The N-terminal segment of the D₂-Rs and D₃-R was also shown to interact with neuronally enriched 4.1N protein; an interaction that contributes to the localization and stability of D₂-Rs at the neuronal PM [85]. Similarly, filamin-A (FLN-A) also interacts with the N-terminal segment of the ICL3 of the D₂-R and D₃-R, and connects D-Rs with some other GPCRs, such as rhodopsin and, metabotropic glutamate receptors to the cytoskeleton, and therefore participate in their final subcellular localization [86]. The dopamine transporter (DAT) is a membrane-spanning protein that facilitates the reuptake of extracellular dopamine to the cytosol and is therefore, an essential target for cocaine, amphetamine, and some other drugs of abuse. One study showed a direct interaction between the DAT and the ICL3 (I340-Q373) of both D₂-R isoforms. However, D_{2L}-R is more capable of physically interacting with the DAT [87].

The Ca²⁺-binding protein calmodulin (CaM) binds to the N-terminal portion of the ICL3 of the D_{2L}-R, within an Arg-rich epitope (VLRRRRKRVN) that is also involved in the binding to G_{i/o} proteins and the adenosine A_{2A} receptor, with the formation of A_{2A}-D₂-R heteromers [88,89]. N-ethylmaleimide-sensitive factor (NSF) is an ATPase and an essential part of the protein network responsible for different membrane fusion events, including transport through the GA and exocytosis [90]. Using immunoprecipitation and

in vitro binding assays, it has been shown that NSF binds to the ICL3 of D₂-R (F341-Q373) and has a putative role in the interaction of D₂-R and the Glu2 AMPA receptor [91]. Agonist stimulation of D₂-R promotes the formation of direct protein-protein interactions between the ICL3 of the D₂-R and the ATPase N-ethylmaleimide-sensitive factor (NSF). Spinophilin is F-actin and protein phosphatase-1-binding protein with a single PDZ domain that was identified as a protein associated with the ICL3 region of the D₂-R. It is hypothesized to be necessary for establishing signaling complexes for dopaminergic neurotransmission through D₂-Rs by linking receptors to downstream signaling molecules and the actin cytoskeleton [92].

Three additional hypothetical ICL3 D₂-R interaction partners were suggested by ISM: prolactin regulatory element-binding protein (PREB), chromaffin granule amine transporter (CGAT) and trafficking protein particle complex subunit 9 (TRAPPC9). Among prospective partners, CGAT displayed the highest affinity for interacting with the ICL3 D₂-R, followed by TRAPPC9 and PREB. For all three prospective interaction partners we were unable to find experimental evidence for the direct interaction with the ICL3 of the D₂-R but only some indirect indication for their involvement in dopamine synthesis, transport, or D₂-R binding. PREB is an ubiquitously expressed protein and, a member of the WD-repeat protein family, that acts as a transcriptional regulator and suppresses the expression of the adiponectin gene [93], regulates prolactin (PRL) gene expression [94] and functions as a transcriptional regulator of PRL promoter activity, and therefore might be involved in thyrotropin-releasing hormone (TRH)-induced PRL gene transcription [95]. PRL gene expression and secretion are regulated by various hormones and growth factors, including dopamine, epidermal growth factor, and thyrotropin-releasing hormone (TRH) [95]. PREB is highly expressed in the anterior pituitary. Prolactinomas are the most common pituitary tumors and are treated with the selective dopamine D₂-R agonist cabergoline [96]. Mutation of the PREB-binding site within the promoter abrogated the ability of cabergoline to inhibit PRL promoter activity. The chromaffin granule amine transporter (CGAT), also named the vesicular monoamine transporter 1 (VMAT1), is involved in the transport of biogenic monoamines, such as serotonin, from the cytoplasm into the secretory vesicles of neuroendocrine and endocrine cells. It has a positive impact on dopamine synthesis, secretion, and transport to storage vesicles, which releases neurotransmitters into synapses as chemical messages to postsynaptic neurons [97]. The pharmaceutical industry also targets VMATs for treating hypertension, drug addiction, psychiatric disorders, Parkinson's disease, and other neurological disorders. The trafficking protein particle complex subunit 9 (TRAPPC9), also known as NIBP, belongs to the TRAPP II multiprotein complex. TRAPPC9 is involved in vesicular trafficking from the ER to the GA and promotes the activation of NFκB signaling. It is highly expressed in the postmitotic neurons of the cerebral cortex [98].

To the best of our knowledge, only two proteins have been identified that specifically interact only with the D_{2L}-R i.e., 29 AA within its ICL3. These proteins are fatty acid-binding protein 3 (FABP3) [99] and Rabaptin-5 interacting protein (Rabex-5) [23]. Fatty acid-binding protein 3 (FABP3), also named the heart-type FABP (H-FAB), is one of the novel 29 AA insert binding protein on the position (G242-V270), which also alters D_{2L}-R function [99]. D_{2L}-R, when activated with a ligand, is known to activate the mitogen-activated protein kinase/extracellular signal-regulated kinase (ERK) pathways, which are enhanced by FABP3 in FABP3-overexpressed cells, showing that FABP3 enhances D_{2L}-R signaling [14]. A co-expression study of D_{2L}-R and D_{2S}-R with this protein in NG108-15 cells shows overexpression and colocalization of endogenous FABP only with the D_{2L}-R in the GA and ER but not in the PM [100]. Dysfunction of FABP3 protein binding to D_{2L}-R was shown in FABP3 KO mice [100], which affects emotional behavior, and is characteristic of neurodegenerative diseases such as schizophrenia and Alzheimer's disorder. These KO mice, which showed altered sensory, motor, and emotional behaviors, also exhibited decreased methamphetamine-induced sensitization and enhanced haloperidol-induced catalepsy due to D₂-R dysfunction. Impaired FABP brain function was observed as an essential factor in the perturbation of D₂-R signaling [101]. Rabaptin-5 interacting protein (Rabex-5) was identified in mouse brain lysates as another protein binding the 29 AA of D_{2L}-R and has been shown to promote the

early-endosome formation and Rab5 activation [71]. Both proteins are essential for prolonged D_{2L}-R mediated ERK signaling.

DRIPs have the propensity to bind to conserved motifs in receptors. For D₁-R it was shown that the ER-membrane-associated protein DRiP78 binds to a FXXXXXXF motif in the C-terminus of D₁-R and other GPCRs. Overexpression or down-modulation of this putative two-TM domain protein leads to ER retention of D₁-Rs, reduced ligand binding, and impaired kinetics of receptor glycosylation [48]. This mechanism acts as a chaperone and may control PM receptor targeting without traveling to the cell surface.

Some of the DRIPs are also possible “private” chaperones with other functions, escorting proteins for D_{2L}-R or proteins of the quality-control machinery involved in its retention within intracellular compartments [37] and facilitating receptor cell surface expression by enabling their trafficking to the PM. Pools of intracellular D₁-R exist in renal tubular cells, and receptor recruitment to the PM is independent of agonist activation elicited by the activation of cell surface receptors and via atrial natriuretic peptide-dependent heterologous activation [102,103].

Table 2. The bioinformatics approach—informational spectrum method (ISM) analysis of interaction partners of the third cytoplasmic loop (ICL3) of the D₂-R. A lower signal to noise S/N ratio suggests a lower interaction affinity between tested protein partners.

Interaction Partner	S/N Ratio	Function	Reference
Glutamate, NMDA (NR2B)	62.39	ionotropic glutamate receptor	Liu, X.Y. et al. (2006) [82]
Par-4	48.63	regulatory component in dopamine signaling	Guo, Q. et al. (1998) [83] Park, S.K. et al. (2005) [84]
Protein 4.1N	38.61	membrane-cytoskeleton adaptor	Binda, A.V. et al. (2002) [85]
FLN-A	26.65	actin binding protein	Lin, R. et al. (2001) [86]
DAT	20.29	facilitating reuptake of extracellular dopamine back in the cytosol	Lee, F.J. et al. (2007) [87]
Gα i/z/o	17.85	binding GPCRs	
CaM	13.36	intermediate calcium-binding messenger	Navarro, G. et al. (2009) [88]
NSF	13.03	ATPase	Hanson, P.I. et al. (1995) [90] Zou S. et al. (2005) [91]
Spinophilin	12.14	F-actin and protein phosphatase-1-binding protein	Smith, F.D. et al. (1999) [92]
Predicted Interaction Partner	12.14		
CGAT	19.90	involved in the transport of biogenic monoamines	
TRAPPC9	19.73	involved in vesicular trafficking from ER to GA	
PREB	18.78	transcriptional regulator	

Legend: Glutamate, NMDA (NR2B)—NR2B subunit of the NMDA glutamate receptor (N-methyl-D-aspartate); FLN-A—filamin-A; Par-4—prostate apoptosis response-4; DAT—dopamine transporter; CGAT—chromaffin granule amine transporter; TRAPPC9—trafficking protein particle complex subunit 9; PREB—prolactin regulatory element-binding protein; NSF—N-ethylmaleimide-sensitive factor; CaM—Ca²⁺-binding protein calmodulin.

10. Interaction with ER-Resident Gatekeeper Proteins

ER gatekeeper proteins tightly control receptor cell-surface export. GTRAP3-18, an integral ER membrane protein, was introduced as a protein both in vivo and in vitro, and is dynamically induced by retinoic acid and inhibits the activity of EAAC1 in a dose dependent manner. GTRAP3-18 forms

an oligomeric complex with D₂-R before exiting the ER, increasing the population of high-mannose oligosaccharide state proteins, and restricting its subcellular localization to the ER [104]. There is evidence that the specific gatekeeper protein PRAF2 binds to subunit GABA₁ of the GABA_B receptor and prevents its progression in the biosynthetic pathway [47]. Dupre, J.D. et al. showed that one of the ER-resident proteins, which is known to regulate trafficking via a FXXXXXXX motif of D-Rs and interact mostly with the G γ subunit and not G α or G β subunits in HEK-293 cells, is dopamine-receptor interaction protein 78 (DRIP78) [105]. Another ER-gatekeeper candidate of the D₂-D₃ heterodimer is an activator of G protein signaling 3 (AGS3), which binds to G_{i α} GDP and inhibits GDP dissociation in the prefrontal cortex during late withdrawal from repeated cocaine administration. However, this actual mechanism in D₂-R signaling is still unknown [106]. In familiar, as well as in sporadic Parkinson's disease, a mutation in the leucine-rich repeat kinase 2 gene (LRRK2) represents the most frequent genetic cause of disease. LRRK2 is a member of the Roco superfamily of proteins, a novel multi-domain family of Ras-like G-proteins, involved in vesicle-mediated transport to the cell membrane. Because LRRK2 could affect D₂-R turnover by decreasing this rate of trafficking from the GA to the CM, the LRRK2 could have an essential function in one of the possible retention mechanisms [107].

11. Clinical Relevance

This review examines the role of different types of conserved retention motifs and DRIPs on D₂-R and roles in routing regulation. The impact of both is of high importance for the physiological functions of D_{2L}-R and its export trafficking and precise localization in the cell. Defective transport of D_{2L}-R, as well as many other GPCRs from the ER to the cell surface, is a highly regulated, dynamic process and is associated with the pathogenesis of a variety of human diseases, therefore advances in our understanding of GPCR export. Thus, the secretory pathway and its role in proper cell function are of high importance.

So far, we are not aware of any known mutation in the 29 AA region that would be associated with a disease. No variant of D_{2L}-R has been linked or associated with schizophrenia, substance abuse, or alcoholism, including the most extensively investigated Ser311Cys polymorphisms of the D₂-R gene. In vitro studies showed that the Cys311-type D₂-R impairs dopamine-induced sequestration, which appears to be consistent with the dopamine hypothesis [108]. A naturally occurring synonymous mutation of the human D₂-R gene (C957T, P319P) is postulated to correlate with the schizophrenia phenotype, and was shown to markedly change mRNA stability via changes in mRNA secondary structure and reduced dopamine-induced up-regulation of D₂-R expression [109]. We also conducted a GPCR database (GPCRdb) search to find additional mutants of the D₂-R and presented them in Table 3.

Table 3. A summary of mutations within the ICL3 of the D₂-R (source: G protein-coupled receptor (GPCR) database (GPCRdb); <http://gpcrdb.org/mutations/render>).

Receptor	AA Residue	Location	Mutation	Reference
drd2_human	249	ICL3	D => V	Guiramand, J. et al. (1995) [110]
drd2_human	264	ICL3	P => G	Guiramand, J. et al. (1995) [110]
drd2_human	310	ICL3	P => S	Kaiser, R. et al. (2003) [111]
drd2_human	311	ICL3	S => C	Kaiser, R. et al. (2003) [111]
drd2_human	311	ICL3	S => C	Goldman D et al. (1997) [112]
drd2_mouse	251	ICL3	K => V	Guiramand, J. et al. (1995) [110]
drd2_mouse	271	ICL3	D => V	Guiramand, J. et al. (1995) [110]
drd2_rat	233	ICL3	R => G	Senogles, SE. et al. (2004) [113]
drd2_rat	234	ICL3	A => T	Senogles, SE. et al. (2004) [113]

We have not found additional mutations located within the insert (AA 241–270). Further elucidation of the regulatory mechanism underlying GPCR export trafficking may provide an essential foundation for developing new therapeutic strategies in treating diseases.

The described differences in the evolutionarily preserved region within the 29 AA insertion in the ICL3 of D_{2L}-R influences the regulation of D_{2L}-R cellular trafficking. A thorough search through motifs showed a conserved arginine cluster within the 29-AA insert of ICL3 of the D_{2L}-R, which appears to be the ER retention signal. Identifying possible candidates for DRIPs may also reveal “private” chaperones, which often display different functions or escort proteins for D_{2L}-R or proteins of the quality-control machinery that play a role in GPCR retention within intracellular compartments. We speculate that other specific retention mechanisms for D_{2L}-R exist. Thus, improving our knowledge of the routing regulation of these critical receptors will probably elicit the development of new therapeutic approaches in controlling the targeting of D_{2L}-R at the PM.

Author Contributions: Conceptualization, V.K. and M.V.; methodology, K.B.C., M.M., S.G., M.S., and M.M.; Investigation, K.B.C. and M.M.; resources, K.B.C. and M.M.; writing—review and editing, V.K., K.B.C., M.M., M.V., and C.S.R.; visualization, M.M.; project administration, V.K. and K.B.C.; funding acquisition, M.V. and M.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Slovenian Research Agency Programme (P4-0053), Slovenian-Serbian bilateral project (BI-RS/20-21-045), Slovenian Research Agency for PhD funding for M. Mavri and grant (No. 173001) from the Ministry of Education, Science and Technological Development of the Republic of Serbia.

Acknowledgments: The authors would acknowledge above mentioned funding. V. Kubale, M. Vrecl, M. Mavri, and M. Senčanski participate in the European COST Action CA 18133 (ERNEST).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Jackson, D.M.; Westlind-Danielsson, A. Dopamine receptors: Molecular biology, biochemistry and behavioural aspects. *Pharmacol. Ther.* **1994**, *64*, 291–370. [[CrossRef](#)]
2. Beaulieu, J.M.; Espinoza, S.; Gainetdinov, R.R. Dopamine receptors—IUPHAR Review 13. *Br. J. Pharmacol.* **2015**, *172*, 1–23. [[CrossRef](#)] [[PubMed](#)]
3. Missale, C.; Nash, S.R.; Robinson, S.W.; Jaber, M.; Caron, M.G. Dopamine receptors: From structure to function. *Physiol. Rev.* **1998**, *78*, 189–225. [[CrossRef](#)] [[PubMed](#)]
4. Min, C.; Zheng, M.; Zhang, X.; Guo, S.; Kwon, K.J.; Shin, C.Y.; Kim, H.S.; Cheon, S.H.; Kim, K.M. N-linked Glycosylation on the N-terminus of the dopamine D2 and D3 receptors determines receptor association with specific microdomains in the plasma membrane. *Biochim. Biophys. Acta* **2015**, *1853*, 41–51. [[CrossRef](#)] [[PubMed](#)]
5. Pivonello, R.; Ferone, D.; Lombardi, G.; Colao, A.; Lamberts, S.W.J.; Hofland, L.J. Novel insights in dopamine receptor physiology. *Eur. J. Endocrinol.* **2007**, *156* (Suppl. 1), S13–S21. [[CrossRef](#)]
6. Dohlman, H.G.; Caron, M.G.; DeBlasi, A.; Frielle, T.; Lefkowitz, R.J. Role of extracellular disulfide-bonded cysteines in the ligand binding function of the beta 2-adrenergic receptor. *Biochemistry* **1990**, *29*, 2335–2342. [[CrossRef](#)]
7. Gazi, L.; Nickolls, S.A.; Strange, P.G. Functional coupling of the human dopamine D2 receptor with G alpha i1, G alpha i2, G alpha i3 and G alpha o G proteins: Evidence for agonist regulation of G protein selectivity. *Br. J. Pharmacol.* **2003**, *138*, 775–786. [[CrossRef](#)]
8. Jiang, M.; Spicher, K.; Boulay, G.; Wang, Y.; Birnbaumer, L. Most central nervous system D2 dopamine receptors are coupled to their effectors by Go. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 3577–3582. [[CrossRef](#)]
9. Senogles, S.E.; Spiegel, A.M.; Padrell, E.; Iyengar, R.; Caron, M.G. Specificity of receptor-G protein interactions. Discrimination of Gi subtypes by the D2 dopamine receptor in a reconstituted system. *J. Biol. Chem.* **1990**, *265*, 4507–4514.
10. Montmayeur, J.P.; Guiramand, J.; Borrelli, E. Preferential coupling between dopamine D2 receptors and G-proteins. *Mol. Endocrinol.* **1993**, *7*, 161–170.
11. Zuk, J.; Bartuzi, D.; Matosiuk, D.; Kaczor, A.A. Preferential Coupling of Dopamine D2S and D2L Receptor Isoforms with Gi1 and Gi2 Proteins-In Silico Study. *Int. J. Mol. Sci.* **2020**, *21*, 436. [[CrossRef](#)]
12. Zhou, Q.; Yang, D.; Wu, M.; Guo, Y.; Guo, W.; Zhong, L.; Cai, X.; Dai, A.; Jang, W.; Shakhnovich, E.I.; et al. Common activation mechanism of class A GPCRs. *eLife* **2019**, *8*, e50279. [[CrossRef](#)] [[PubMed](#)]
13. Yin, J.; Chen, K.M.; Clark, M.J.; Hijazi, M.; Kumari, P.; Bai, X.C.; Sunahara, R.K.; Barth, P.; Rosenbaum, D.M. Structure of a D2 dopamine receptor-G-protein complex in a lipid membrane. *Nature* **2020**, *584*, 125–129. [[CrossRef](#)]

14. Takeuchi, Y.; Fukunaga, K. Differential subcellular localization of two dopamine D2 receptor isoforms in transfected NG108-15 cells. *J. Neurochem.* **2003**, *85*, 1064–1074. [[CrossRef](#)]
15. Kaczor, A.A.; Jorg, M.; Capuano, B. The dopamine D2 receptor dimer and its interaction with homobivalent antagonists: Homology modeling, docking and molecular dynamics. *J. Mol. Model* **2016**, *22*, 203. [[CrossRef](#)] [[PubMed](#)]
16. Madras, B.K. History of the discovery of the antipsychotic dopamine D2 receptor: A basis for the dopamine hypothesis of schizophrenia. *J. Hist. Neurosci.* **2013**, *22*, 62–78. [[CrossRef](#)] [[PubMed](#)]
17. Giros, B.; Sokoloff, P.; Martres, M.P.; Riou, J.F.; Emorine, L.J.; Schwartz, J.C. Alternative splicing directs the expression of two D2 dopamine receptor isoforms. *Nature* **1989**, *342*, 923–926. [[CrossRef](#)] [[PubMed](#)]
18. Monsma, F.J., Jr.; McVittie, L.D.; Gerfen, C.R.; Mahan, L.C.; Sibley, D.R. Multiple D2 dopamine receptors produced by alternative RNA splicing. *Nature* **1989**, *342*, 926–929. [[CrossRef](#)]
19. Dal Toso, R.; Sommer, B.; Ewert, M.; Herb, A.; Pritchett, D.B.; Bach, A.; Shivers, B.D.; Seeburg, P.H. The dopamine D2 receptor: Two molecular forms generated by alternative splicing. *EMBO J.* **1989**, *8*, 4025–4034. [[CrossRef](#)]
20. Pelham, H.R. Sorting and retrieval between the endoplasmic reticulum and Golgi apparatus. *Curr. Opin. Cell Biol.* **1995**, *7*, 530–535. [[CrossRef](#)]
21. Montmayeur, J.P.; Bausero, P.; Amlaiky, N.; Maroteaux, L.; Hen, R.; Borrelli, E. Differential expression of the mouse D2 dopamine receptor isoforms. *FEBS Lett.* **1991**, *278*, 239–243. [[CrossRef](#)]
22. Sedaghat, K.; Nantel, M.F.; Ginsberg, S.; Lalonde, V.; Tiberi, M. Molecular characterization of dopamine D2 receptor isoforms tagged with green fluorescent protein. *Mol. Biotechnol.* **2006**, *34*, 1–14. [[CrossRef](#)]
23. Prou, D.; Gu, W.J.; Le Crom, S.; Vincent, J.D.; Salamero, J.; Vernier, P. Intracellular retention of the two isoforms of the D(2) dopamine receptor promotes endoplasmic reticulum disruption. *J. Cell Sci.* **2001**, *114 Pt 19*, 3517–3527.
24. Usiello, A.; Baik, J.H.; Rouge-Pont, F.; Picetti, R.; Dierich, A.; LeMeur, M.; Piazza, P.V.; Borrelli, E. Distinct functions of the two isoforms of dopamine D2 receptors. *Nature* **2000**, *408*, 199–203. [[CrossRef](#)]
25. Radl, D.; Chiacchiaretta, M.; Lewis, R.G.; Brami-Cherrier, K.; Arcuri, L.; Borrelli, E. Differential regulation of striatal motor behavior and related cellular responses by dopamine D2L and D2S isoforms. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 198–203. [[CrossRef](#)] [[PubMed](#)]
26. Robinson, B.G.; Condon, A.F.; Radl, D.; Borrelli, E.; Williams, J.T.; Neve, K.A. Cocaine-induced adaptation of dopamine D2S, but not D2L autoreceptors. *eLife* **2017**, *6*, e31924. [[CrossRef](#)]
27. Margeta-Mitrovic, M.; Jan, Y.N.; Jan, L.Y. A trafficking checkpoint controls GABA(B) receptor heterodimerization. *Neuron* **2000**, *27*, 97–106. [[CrossRef](#)]
28. Li, P.; Snyder, G.L.; Vanover, K.E. Dopamine Targeting Drugs for the Treatment of Schizophrenia: Past, Present and Future. *Curr. Top. Med. Chem.* **2016**, *16*, 3385–3403. [[CrossRef](#)]
29. Wang, S.; Che, T.; Levit, A.; Shoichet, B.K.; Wacker, D.; Roth, B.L. Structure of the D2 dopamine receptor bound to the atypical antipsychotic drug risperidone. *Nature* **2018**, *555*, 269–273. [[CrossRef](#)]
30. Martire, G.; Mottola, G.; Pascale, M.C.; Malagolini, N.; Turrini, I.; Serafini-Cessi, F.; Jackson, M.R.; Bonatti, S. Different fate of a single reporter protein containing KDEL or KKXX targeting signals stably expressed in mammalian cells. *J. Biol. Chem.* **1996**, *271*, 3541–3547. [[CrossRef](#)]
31. Gomez-Navarro, N.; Miller, E. Protein sorting at the ER-Golgi interface. *J. Cell Biol.* **2016**, *215*, 769–778. [[CrossRef](#)] [[PubMed](#)]
32. Khan, Z.U.; Mrzljak, L.; Gutierrez, A.; de la Calle, A.; Goldman-Rakic, P.S. Prominence of the dopamine D2 short isoform in dopaminergic pathways. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 7731–7736. [[CrossRef](#)]
33. Lopez-Aranda, M.F.; Acevedo, M.J.; Gutierrez, A.; Koulen, P.; Khan, Z.U. Role of a Galphai2 protein splice variant in the formation of an intracellular dopamine D2 receptor pool. *J. Cell Sci.* **2007**, *120 Pt 13*, 2171–2178. [[CrossRef](#)]
34. Macey, T.A.; Gurevich, V.V.; Neve, K.A. Preferential Interaction between the dopamine D2 receptor and Arrestin2 in neostriatal neurons. *Mol. Pharmacol.* **2004**, *66*, 1635–1642. [[CrossRef](#)] [[PubMed](#)]
35. Ng, G.Y.; Varghese, G.; Chung, H.T.; Trogadis, J.; Seeman, P.; O'Dowd, B.F.; George, S.R. Resistance of the dopamine D2L receptor to desensitization accompanies the up-regulation of receptors on to the surface of Sf9 cells. *Endocrinology* **1997**, *138*, 4199–4206. [[CrossRef](#)] [[PubMed](#)]
36. Van Craenenbroeck, K.; Clark, S.D.; Cox, M.J.; Oak, J.N.; Liu, F.; Van Tol, H.H. Folding efficiency is rate-limiting in dopamine D4 receptor biogenesis. *J. Biol. Chem.* **2005**, *280*, 19350–19357. [[CrossRef](#)] [[PubMed](#)]

37. Achour, L.; Labbe-Jullie, C.; Scott, M.G.; Marullo, S. An escort for GPCRs: Implications for regulation of receptor density at the cell surface. *Trends Pharmacol. Sci.* **2008**, *29*, 528–535. [[CrossRef](#)]
38. Cho, D.I.; Min, C.; Jung, K.S.; Cheong, S.Y.; Zheng, M.; Cheong, S.J.; Oak, M.H.; Cheong, J.H.; Lee, B.K.; Kim, K.M. The N-terminal region of the dopamine D2 receptor, a rhodopsin-like GPCR, regulates correct integration into the plasma membrane and endocytic routes. *Br. J. Pharmacol.* **2012**, *166*, 659–675. [[CrossRef](#)]
39. Fishburn, C.S.; Elazar, Z.; Fuchs, S. Differential glycosylation and intracellular trafficking for the long and short isoforms of the D2 dopamine receptor. *J. Biol. Chem.* **1995**, *270*, 29819–29824.
40. Grunewald, S.; Reilander, H.; Michel, H. In vivo reconstitution of dopamine D2S receptor-mediated G protein activation in baculovirus-infected insect cells: Preferred coupling to Gi1 versus Gi2. *Biochemistry* **1996**, *35*, 15162–15173. [[CrossRef](#)]
41. Senogles, S.E. The D2 dopamine receptor isoforms signal through distinct Gi alpha proteins to inhibit adenylyl cyclase. A study with site-directed mutant Gi alpha proteins. *J. Biol. Chem.* **1994**, *269*, 23120–23127.
42. Choi, D.S.; Wang, D.; Tolbert, L.; Sadee, W. Basal signaling activity of human dopamine D2L receptor demonstrated with an ecdysone-inducible mammalian expression system. *J. Neurosci. Methods* **2000**, *94*, 217–225. [[CrossRef](#)]
43. Wang, Y.; Xu, R.; Sasaoka, T.; Tonegawa, S.; Kung, M.P.; Sankoorikal, E.B. Dopamine D2 long receptor-deficient mice display alterations in striatum-dependent functions. *J. Neurosci. Off. J. Soc. Neurosci.* **2000**, *20*, 8305–8314. [[CrossRef](#)]
44. Welter, M.; Vallone, D.; Samad, T.A.; Meziane, H.; Usiello, A.; Borrelli, E. Absence of dopamine D2 receptors unmasks an inhibitory control over the brain circuitries activated by cocaine. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 6840–6845. [[CrossRef](#)] [[PubMed](#)]
45. Xu, R.; Hranilovic, D.; Fetsko, L.A.; Bucan, M.; Wang, Y. Dopamine D2S and D2L receptors may differentially contribute to the actions of antipsychotic and psychotic agents in mice. *Mol. Psychiatry* **2002**, *7*, 1075–1082. [[CrossRef](#)]
46. Dong, C.; Filipeanu, C.M.; Duvernay, M.T.; Wu, G. Regulation of G protein-coupled receptor export trafficking. *Biochim. Biophys. Acta* **2007**, *1768*, 853–870. [[CrossRef](#)] [[PubMed](#)]
47. Doly, S.; Shirvani, H.; Gata, G.; Meye, F.J.; Emerit, M.B.; Enslin, H.; Achour, L.; Pardo-Lopez, L.; Yang, S.K.; Armand, V.; et al. GABAB receptor cell-surface export is controlled by an endoplasmic reticulum gatekeeper. *Mol. Psychiatry* **2016**, *21*, 480–490. [[CrossRef](#)]
48. Zhang, M.; Wu, G. Mechanisms of the anterograde trafficking of GPCRs: Regulation of AT1R transport by interacting proteins and motifs. *Traffic* **2019**, *20*, 110–120. [[CrossRef](#)]
49. Milligan, G. The role of dimerisation in the cellular trafficking of G-protein-coupled receptors. *Curr. Opin. Pharmacol.* **2010**, *10*, 23–29. [[CrossRef](#)]
50. Wu, G. Regulation of post-Golgi traffic of G protein-coupled receptors. *Sub-Cell. Biochem.* **2012**, *63*, 83–95.
51. Spang, A. Retrograde traffic from the Golgi to the endoplasmic reticulum. *Cold Spring Harb. Perspect. Biol.* **2013**, *5*, a013391. [[CrossRef](#)] [[PubMed](#)]
52. Jackson, M.R.; Nilsson, T.; Peterson, P.A. Identification of a consensus motif for retention of transmembrane proteins in the endoplasmic reticulum. *EMBO J.* **1990**, *9*, 3153–3162. [[CrossRef](#)] [[PubMed](#)]
53. Munro, S.; Pelham, H.R. A C-terminal signal prevents secretion of luminal ER proteins. *Cell* **1987**, *48*, 899–907. [[CrossRef](#)]
54. Jin, H.; Komita, M.; Aoe, T. The Role of BiP Retrieval by the KDEL Receptor in the Early Secretory Pathway and its Effect on Protein Quality Control and Neurodegeneration. *Front. Mol. Neurosci.* **2017**, *10*, 222. [[CrossRef](#)]
55. Rothman, J.E.; Wieland, F.T. Protein sorting by transport vesicles. *Science* **1996**, *272*, 227–234. [[CrossRef](#)] [[PubMed](#)]
56. Cukierman, E.; Huber, I.; Rotman, M.; Cassel, D. The ARF1 GTPase-activating protein: Zinc finger motif and Golgi complex localization. *Science* **1995**, *270*, 1999–2002. [[CrossRef](#)]
57. Aoe, T.; Huber, I.; Vasudevan, C.; Watkins, S.C.; Romero, G.; Cassel, D.; Hsu, V.W. The KDEL receptor regulates a GTPase-activating protein for ADP-ribosylation factor 1 by interacting with its non-catalytic domain. *J. Biol. Chem.* **1999**, *274*, 20545–20549. [[CrossRef](#)]
58. Boehmler, W.; Obrecht-Pflumio, S.; Canfield, V.; Thisse, C.; Thisse, B.; Levenson, R. Evolution and expression of D2 and D3 dopamine receptor genes in zebrafish. *Dev. Dyn. Off. Publ. Am. Assoc. Anat.* **2004**, *230*, 481–493. [[CrossRef](#)]

59. Kokubun, H.; Jin, H.; Aoe, T. Pathogenic Effects of Impaired Retrieval between the Endoplasmic Reticulum and Golgi Complex. *Int. J. Mol. Sci.* **2019**, *20*, 5614. [[CrossRef](#)]
60. Cabrera, M.; Muniz, M.; Hidalgo, J.; Vega, L.; Martin, M.E.; Velasco, A. The retrieval function of the KDEL receptor requires PKA phosphorylation of its C-terminus. *Mol. Biol. Cell* **2003**, *14*, 4114–4125. [[CrossRef](#)]
61. Jackson, L.P.; Lewis, M.; Kent, H.M.; Edeling, M.A.; Evans, P.R.; Duden, R.; Owen, D.J. Molecular basis for recognition of dilysine trafficking motifs by COPI. *Dev. Cell* **2012**, *23*, 1255–1262. [[CrossRef](#)] [[PubMed](#)]
62. Letourneur, F.; Gaynor, E.C.; Hennecke, S.; Demolliere, C.; Duden, R.; Emr, S.D.; Riezman, H.; Cosson, P. Coatamer is essential for retrieval of dilysine-tagged proteins to the endoplasmic reticulum. *Cell* **1994**, *79*, 1199–1207. [[CrossRef](#)]
63. Vrecl, M.; Anderson, L.; Hanyaloglu, A.; McGregor, A.M.; Groarke, A.D.; Milligan, G.; Taylor, P.L.; Eidne, K.A. Agonist-induced endocytosis and recycling of the gonadotropin-releasing hormone receptor: Effect of beta-arrestin on internalization kinetics. *Mol. Endocrinol.* **1998**, *12*, 1818–1829. [[PubMed](#)]
64. Pagano, A.; Rovelli, G.; Mosbacher, J.; Lohmann, T.; Duthey, B.; Stauffer, D.; Ristig, D.; Schuler, V.; Meigel, I.; Lampert, C.; et al. C-terminal interaction is essential for surface trafficking but not for heteromeric assembly of GABA(b) receptors. *J. Neurosci. Off. J. Soc. Neurosci.* **2001**, *21*, 1189–1202. [[CrossRef](#)]
65. Kubale, V.; Pogacnik, A.; Schwartz, W.T.; Vrecl, M. Seven transmembrane receptors (7TM) in the view of dimerization and experimental methods to study their dimerization and cross-talk. *Slov. Vet. Res.* **2008**, *45*, 89–102.
66. Kuner, R.; Kohr, G.; Grunewald, S.; Eisenhardt, G.; Bach, A.; Kornau, H.C. Role of heteromer formation in GABAB receptor function. *Science* **1999**, *283*, 74–77. [[CrossRef](#)]
67. Filipeanu, C.M.; Pullikuth, A.K.; Guidry, J.J. Molecular determinants of the human alpha2C-adrenergic receptor temperature-sensitive intracellular traffic. *Mol. Pharmacol.* **2015**, *87*, 792–802. [[CrossRef](#)]
68. Nasu-Nishimura, Y.; Hurtado, D.; Braud, S.; Tang, T.T.; Isaac, J.T.; Roche, K.W. Identification of an endoplasmic reticulum-retention motif in an intracellular loop of the kainate receptor subunit KA2. *J. Neurosci. Off. J. Soc. Neurosci.* **2006**, *26*, 7014–7021. [[CrossRef](#)]
69. Hermosilla, R.; Schulein, R. Sorting functions of the individual cytoplasmic domains of the G protein-coupled vasopressin V(2) receptor in Madin Darby canine kidney epithelial cells. *Mol. Pharmacol.* **2001**, *60*, 1031–1039. [[CrossRef](#)]
70. Kubale, V.; Blagotinsek, K.; Nohr, J.; Eidne, K.A.; Vrecl, M. The Conserved Arginine Cluster in the Insert of the Third Cytoplasmic Loop of the Long Form of the D(2) Dopamine Receptor (D2L-R) Acts as an Intracellular Retention Signal. *Int. J. Mol. Sci.* **2016**, *17*, 1152. [[CrossRef](#)]
71. Shioda, N.; Takeuchi, Y.; Fukunaga, K. Advanced research on dopamine signaling to develop drugs for the treatment of mental disorders: Proteins interacting with the third cytoplasmic loop of dopamine D2 and D3 receptors. *J. Pharmacol. Sci.* **2010**, *114*, 25–31. [[CrossRef](#)] [[PubMed](#)]
72. Navarro, G.; Ferre, S.; Cordini, A.; Moreno, E.; Mallol, J.; Casado, V.; Cortes, A.; Hoffmann, H.; Ortiz, J.; Canela, E.I.; et al. Interactions between intracellular domains as key determinants of the quaternary structure and function of receptor heteromers. *J. Biol. Chem.* **2010**, *285*, 27346–27359. [[CrossRef](#)] [[PubMed](#)]
73. Lukasiewicz, S.; Faron-Gorecka, A.; Dobrucki, J.; Polit, A.; Dziedzicka-Wasylewska, M. Studies on the role of the receptor protein motifs possibly involved in electrostatic interactions on the dopamine D1 and D2 receptor oligomerization. *FEBS J.* **2009**, *276*, 760–775. [[CrossRef](#)] [[PubMed](#)]
74. Lukasiewicz, S.; Polit, A.; Kedracka-Krok, S.; Wedzony, K.; Mackowiak, M.; Dziedzicka-Wasylewska, M. Hetero-dimerization of serotonin 5-HT(2A) and dopamine D(2) receptors. *Biochim. Biophys. Acta* **2010**, *1803*, 1347–1358. [[CrossRef](#)] [[PubMed](#)]
75. Likhite, N.; Jackson, C.A.; Liang, M.S.; Krzyzanowski, M.C.; Lei, P.; Wood, J.F.; Birkaya, B.; Michaels, K.L.; Andreadis, S.T.; Clark, S.D.; et al. The protein arginine methyltransferase PRMT5 promotes D2-like dopamine receptor signaling. *Sci. Signal.* **2015**, *8*, ra115. [[CrossRef](#)] [[PubMed](#)]
76. Kumar, M.; Gouw, M.; Michael, S.; Diakogianni, A.; Valverde, J.A.; Bukirova, D.; Calyseva, J.; Palopoli, N.; Davey, N.E.; Chemes, L.B.; et al. ELM—the eukaryotic linear motif resource in 2020. *Nucleic Acids Res.* **2020**, *48*, D296–D306. [[CrossRef](#)]
77. Duvernay, M.T.; Filipeanu, C.M.; Wu, G. The regulatory mechanisms of export trafficking of G protein-coupled receptors. *Cell. Signal.* **2005**, *17*, 1457–1465. [[CrossRef](#)]
78. Bermak, J.C.; Li, M.; Bullock, C.; Zhou, Q.Y. Regulation of transport of the dopamine D1 receptor by a new membrane-associated ER protein. *Nat. Cell Biol.* **2001**, *3*, 492–498. [[CrossRef](#)]

79. Kabbani, N.; Woll, M.P.; Nordman, J.C.; Levenson, R. Dopamine receptor interacting proteins: Targeting neuronal calcium sensor-1/D2 dopamine receptor interaction for antipsychotic drug development. *Curr. Drug Targets* **2012**, *13*, 72–79. [[CrossRef](#)]
80. Sencanski, M.; Glisic, S.; Snajder, M.; Veljkovic, N.; Poklar Ulrih, N.; Mavri, J.; Vrecl, M. Computational design and characterization of nanobody-derived peptides that stabilize the active conformation of the beta2-adrenergic receptor (beta2-AR). *Sci. Rep.* **2019**, *9*, 16555. [[CrossRef](#)]
81. Mandic, M.; Drinovec, L.; Glisic, S.; Veljkovic, N.; Nohr, J.; Vrecl, M. Demonstration of a direct interaction between beta2-adrenergic receptor and insulin receptor by BRET and bioinformatics. *PLoS ONE* **2014**, *9*, e112664. [[CrossRef](#)] [[PubMed](#)]
82. Liu, X.Y.; Chu, X.P.; Mao, L.M.; Wang, M.; Lan, H.X.; Li, M.H.; Zhang, G.C.; Parelkar, N.K.; Fibuch, E.E.; Haines, M.; et al. Modulation of D2R-NR2B interactions in response to cocaine. *Neuron* **2006**, *52*, 897–909. [[CrossRef](#)] [[PubMed](#)]
83. Guo, Q.; Fu, W.; Xie, J.; Luo, H.; Sells, S.F.; Geddes, J.W.; Bondada, V.; Rangnekar, V.M.; Mattson, M.P. Par-4 is a mediator of neuronal degeneration associated with the pathogenesis of Alzheimer disease. *Nat. Med.* **1998**, *4*, 957–962. [[CrossRef](#)]
84. Park, S.K.; Nguyen, M.D.; Fischer, A.; Luke, M.P.; Affar, E.B.; Dieffenbach, P.B.; Tseng, H.C.; Shi, Y.; Tsai, L.H. Par-4 links dopamine signaling and depression. *Cell* **2005**, *122*, 275–287. [[CrossRef](#)] [[PubMed](#)]
85. Binda, A.V.; Kabbani, N.; Lin, R.; Levenson, R. D2 and D3 dopamine receptor cell surface localization mediated by interaction with protein 4.1N. *Mol. Pharmacol.* **2002**, *62*, 507–513. [[CrossRef](#)] [[PubMed](#)]
86. Lin, R.; Karpa, K.; Kabbani, N.; Goldman-Rakic, P.; Levenson, R. Dopamine D2 and D3 receptors are linked to the actin cytoskeleton via interaction with filamin A. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 5258–5263. [[CrossRef](#)]
87. Lee, F.J.; Pei, L.; Moszczynska, A.; Vukusic, B.; Fletcher, P.J.; Liu, F. Dopamine transporter cell surface localization facilitated by a direct interaction with the dopamine D2 receptor. *EMBO J.* **2007**, *26*, 2127–2136. [[CrossRef](#)]
88. Navarro, G.; Aymerich, M.S.; Marcellino, D.; Cortes, A.; Casado, V.; Mallo, J.; Canela, E.I.; Agnati, L.; Woods, A.S.; Fuxe, K.; et al. Interactions between calmodulin, adenosine A2A, and dopamine D2 receptors. *J. Biol. Chem.* **2009**, *284*, 28058–28068. [[CrossRef](#)]
89. Woods, A.S.; Marcellino, D.; Jackson, S.N.; Franco, R.; Ferre, S.; Agnati, L.F.; Fuxe, K. How calmodulin interacts with the adenosine A(2A) and the dopamine D(2) receptors. *J. Proteome Res.* **2008**, *7*, 3428–3434. [[CrossRef](#)]
90. Hanson, P.I.; Otto, H.; Barton, N.; Jahn, R. The N-ethylmaleimide-sensitive fusion protein and alpha-SNAP induce a conformational change in syntaxin. *J. Biol. Chem.* **1995**, *270*, 16955–16961. [[CrossRef](#)]
91. Zou, S.; Li, L.; Pei, L.; Vukusic, B.; Van Tol, H.H.; Lee, F.J.; Wan, Q.; Liu, F. Protein-protein coupling/uncoupling enables dopamine D2 receptor regulation of AMPA receptor-mediated excitotoxicity. *J. Neurosci. Off. J. Soc. Neurosci.* **2005**, *25*, 4385–4395. [[CrossRef](#)] [[PubMed](#)]
92. Smith, F.D.; Oxford, G.S.; Milgram, S.L. Association of the D2 dopamine receptor third cytoplasmic loop with spinophilin, a protein phosphatase-1-interacting protein. *J. Biol. Chem.* **1999**, *274*, 19894–19900. [[CrossRef](#)] [[PubMed](#)]
93. Zhang, X.Z.; Imachi, H.; Lyu, J.Y.; Fukunaga, K.; Sato, S.; Ibata, T.; Kobayashi, T.; Yoshimoto, T.; Kikuchi, F.; Dong, T.; et al. Prolactin regulatory element-binding protein is involved in suppression of the adiponectin gene in vivo. *J. Endocrinol. Investig.* **2017**, *40*, 437–445. [[CrossRef](#)] [[PubMed](#)]
94. Park, J.M.; Kim, M.Y.; Kim, T.H.; Min, D.K.; Yang, G.E.; Ahn, Y.H. Prolactin regulatory element-binding (PREB) protein regulates hepatic glucose homeostasis. *Biochim. Biophys. Acta Mol. Basis Dis.* **2018**, *1864*, 2097–2107. [[CrossRef](#)] [[PubMed](#)]
95. Yu, X.; Murao, K.; Imachi, H.; Li, J.; Nishiuchi, T.; Dobashi, H.; Hosomi, N.; Masugata, H.; Zhang, G.X.; Iwama, H.; et al. The transcription factor prolactin regulatory element-binding protein mediates prolactin transcription induced by thyrotropin-releasing hormone in GH3 cells. *Endocrine* **2010**, *38*, 53–59. [[CrossRef](#)]
96. Zhang, W.; Murao, K.; Imachi, H.; Iwama, H.; Chen, K.; Fei, Z.; Zhang, X.; Ishida, T.; Tamiya, T. Suppression of prolactin expression by cabergoline requires prolactin regulatory element-binding protein (PREB) in GH3 cells. *Horm. Metab. Res.* **2010**, *42*, 557–561. [[CrossRef](#)]
97. Wimalasena, D.S.; Wimalasena, K. Kinetic evidence for channeling of dopamine between monoamine transporter and membranous dopamine-beta-monoxygenase in chromaffin granule ghosts. *J. Biol. Chem.* **2004**, *279*, 15298–15304. [[CrossRef](#)]

98. Mochida, G.H.; Mahajnah, M.; Hill, A.D.; Basel-Vanagaite, L.; Gleason, D.; Hill, R.S.; Bodell, A.; Crosier, M.; Straussberg, R.; Walsh, C.A. A truncating mutation of TRAPPC9 is associated with autosomal-recessive intellectual disability and postnatal microcephaly. *Am. J. Hum. Genet.* **2009**, *85*, 897–902. [[CrossRef](#)]
99. Shioda, N. Dopamine D2L receptor-interacting proteins regulate dopaminergic signaling. *J. Pharmacol. Sci.* **2017**, *135*, 51–54. [[CrossRef](#)]
100. Shioda, N.; Yamamoto, Y.; Watanabe, M.; Binas, B.; Owada, Y.; Fukunaga, K. Heart-type fatty acid binding protein regulates dopamine D2 receptor function in mouse brain. *J. Neurosci. Off. J. Soc. Neurosci.* **2010**, *30*, 3146–3155. [[CrossRef](#)]
101. Yabuki, Y.; Takahata, I.; Matsuo, K.; Owada, Y.; Fukunaga, K. Ramelteon Improves Post-traumatic Stress Disorder-Like Behaviors Exhibited by Fatty Acid-Binding Protein 3 Null Mice. *Mol. Neurobiol.* **2018**, *55*, 3577–3591. [[CrossRef](#)]
102. Brismar, H.; Asghar, M.; Carey, R.M.; Greengard, P.; Aperia, A. Dopamine-induced recruitment of dopamine D1 receptors to the plasma membrane. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 5573–5578. [[CrossRef](#)]
103. Holtback, U.; Brismar, H.; DiBona, G.F.; Fu, M.; Greengard, P.; Aperia, A. Receptor recruitment: A mechanism for interactions between G protein-coupled receptors. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 7271–7275. [[CrossRef](#)] [[PubMed](#)]
104. Ruggiero, A.M.; Liu, Y.; Vidensky, S.; Maier, S.; Jung, E.; Farhan, H.; Robinson, M.B.; Sitte, H.H.; Rothstein, J.D. The endoplasmic reticulum exit of glutamate transporter is regulated by the inducible mammalian Yip6b/GTRAP3-18 protein. *J. Biol. Chem.* **2008**, *283*, 6175–6183. [[CrossRef](#)] [[PubMed](#)]
105. Dupre, D.J.; Robitaille, M.; Richer, M.; Ethier, N.; Mamarbachi, A.M.; Hebert, T.E. Dopamine receptor-interacting protein 78 acts as a molecular chaperone for Ggamma subunits before assembly with Gbeta. *J. Biol. Chem.* **2007**, *282*, 13703–13715. [[CrossRef](#)]
106. Bowers, M.S.; McFarland, K.; Lake, R.W.; Peterson, Y.K.; Lapish, C.C.; Gregory, M.L.; Lanier, S.M.; Kalivas, P.W. Activator of G protein signaling 3: A gatekeeper of cocaine sensitization and drug seeking. *Neuron* **2004**, *42*, 269–281. [[CrossRef](#)]
107. Rassu, M.; Del Giudice, M.G.; Sanna, S.; Taymans, J.M.; Morari, M.; Brugnoli, A.; Frassinetti, M.; Masala, A.; Esposito, S.; Galioto, M.; et al. Role of LRRK2 in the regulation of dopamine receptor trafficking. *PLoS ONE* **2017**, *12*, e0179082. [[CrossRef](#)] [[PubMed](#)]
108. Itokawa, M.; Arinami, T.; Toru, M. Advanced research on dopamine signaling to develop drugs for the treatment of mental disorders: Ser311Cys polymorphisms of the dopamine D2-receptor gene and schizophrenia. *J. Pharmacol. Sci.* **2010**, *114*, 1–5. [[CrossRef](#)] [[PubMed](#)]
109. Duan, J.; Wainwright, M.S.; Comeron, J.M.; Saitou, N.; Sanders, A.R.; Gelernter, J.; Gejman, P.V. Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. *Hum. Mol. Genet.* **2003**, *12*, 205–216. [[CrossRef](#)]
110. Guiramand, J.; Montmayeur, J.P.; Ceraline, J.; Bhatia, M.; Borrelli, E. Alternative splicing of the dopamine D2 receptor directs specificity of coupling to G-proteins. *J. Biol. Chem.* **1995**, *270*, 7354–7358. [[CrossRef](#)]
111. Kaiser, R.; Hofer, A.; Grapengiesser, A.; Gasser, T.; Kupsch, A.; Roots, I.; Brockmoller, J. L-dopa-induced adverse effects in PD and dopamine transporter gene polymorphism. *Neurology* **2003**, *60*, 1750–1755. [[CrossRef](#)] [[PubMed](#)]
112. Goldman, D.; Urbanek, M.; Guenther, D.; Robin, R.; Long, J.C. Linkage and association of a functional DRD2 variant [Ser311Cys] and DRD2 markers to alcoholism, substance abuse and schizophrenia in Southwestern American Indians. *Am. J. Med. Genet.* **1997**, *74*, 386–394. [[CrossRef](#)]
113. Senogles, S.E.; Heimert, T.L.; Odife, E.R.; Quasney, M.W. A region of the third intracellular loop of the short form of the D2 dopamine receptor dictates Gi coupling specificity. *J. Biol. Chem.* **2004**, *279*, 1601–1606. [[CrossRef](#)] [[PubMed](#)]

