

Prevalence and extension of the anterior loop of the mental nerve in different populations and CBCT imaging settings: A systematic review and meta-analysis

Mahdi Hadilou¹, Leila Gholami², Morteza Ghojazadeh³, Naghmeh Emadi^{4,*}

¹Student Research Committee, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran

²Department of Oral Biological and Medical Sciences, Faculty of Dentistry, University of British Columbia, Vancouver, Canada

³Research Center for Evidence Based Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Dental Research Center, Research Institute of Dental Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ABSTRACT

Purpose: This study aimed to identify the prevalence and extension of the anterior loop (AL) of the mental nerve in different populations and according to different cone-beam computed tomography (CBCT) imaging settings.

Materials and Methods: Medline/PubMed, Embase, Scopus, Web of Science, and ProQuest were searched. The main inclusion criterion was ALs evaluated in CBCT images. The quality of studies was assessed with the Joanna Briggs Institute risk of bias checklist. Subgroup analyses were conducted for sex, side, continent, voxel size, field of view, and type of CBCT-reconstruction images with a random-effects model.

Results: Sixty-three studies with 13,743 participants (27,075 hemimandibles) were included. An AL was found in 40.6% (95% CI: 32.8%-48.9%, $P < 0.05$) of participants and 36.0% (95% CI: 27.5%-45.5%, $P < 0.05$) of hemimandibles, in 34.9% (95% CI: 25.1%-46.2%, $P < 0.05$) of males and 34.5% (95% CI: 23.5%-47.4%, $P < 0.05$) of females. The average length of ALs was 2.39 mm (95% CI: 2.07-2.70 mm, $P < 0.05$). Their extension was 2.13 mm (95% CI: 1.54-2.73 mm, $P < 0.05$) in males and 1.85 mm (95% CI: 1.35-2.36 mm, $P < 0.05$) in females. Significant differences were observed regarding the prevalence and length of ALs among continents and for its measured length on different CBCT-reconstruction images, but not between other subgroups.

Conclusion: AL was a relatively common finding. The voxel size and fields of view of CBCT devices were adequate for assessing AL; however, a 2-mm safety margin from anatomical structures (such as the AL) could be recommended to be considered when using CBCT imaging. (*Imaging Sci Dent* 2022; 52: 141-53)

KEY WORDS: Inferior Alveolar Canal; Mandibular Nerve; Cone-Beam Computed Tomography; Systematic Review

Introduction

Intra-oral surgery commonly results in postoperative side effects and complications, one of the most serious of which is unintentional injury to the inferior alveolar nerve (IAN).¹ The mental portion of the IAN is categorized into straight (linear) and perpendicular portions, as well as the anterior loop (AL).² Predicting the position of an AL

is essential in treatment planning for surgical procedures involving flap reflection, osteotomy, bone harvesting, and implant fixture placement in the anterior mandible.^{3,4} Neurosensory disturbances and hemorrhage are the most common complications of these surgical procedures.^{5,6} A recent systematic review on neurosensory disturbances after mandibular implant treatment procedures showed that the incidence of sensory disturbances ranged from 6.5% to 40%, and implants placed in the anterior region had a greater risk for nerve damage.⁷

Various approaches have been suggested for avoiding complications, such as direct visualization during surgery; identification of its location on periapical or panoramic radiography, magnetic resonance imaging, computed tomography, or cone-beam computed tomography (CBCT);

The research protocol was approved and supported by the Student Research Committee, Tabriz University of Medical Sciences [grant number: 68824].

Received January 2, 2022; Revised March 24, 2022; Accepted March 26, 2022

Published online April 28, 2022

*Correspondence to : Dr. Naghmeh Emadi

Dental Research Center, Research Institute of Dental Sciences, Shahid Beheshti University of Medical Sciences, Daneshjoo Blvd, Evin, Tehran, Iran

Tel) 98-2122413897, E-mail) Emadi.naghmeh@gmail.com

Copyright © 2022 by Korean Academy of Oral and Maxillofacial Radiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Imaging Science in Dentistry · pISSN 2233-7822 eISSN 2233-7830

Table 1. Search strategy sample in PubMed.

```

((((((((((((("Mandibular Nerve"[Mesh]) OR ('mandibular nerve'[Text Word])) OR ('mandible branch'[Text Word])) OR ('mandible
nerve'[Text Word])) OR ('mandible ramus'[Text Word])) OR ('mandibular branch'[Text Word])) OR ('mandibular branch, trigeminal
nerve'[Text Word])) OR ('mandibular nerve'[Text Word])) OR ('mandibular ramus'[Text Word])) OR ('nerve, mandibular'[Text Word]))
OR ('nervus mandibularis'[Text Word])) OR ('ramus mandibulae'[Text Word])) OR ('ramus mandibularis nervi trigemini'[Text Word]))
OR ('trigeminal nerve mandibular branch'[Text Word])) OR ('trigeminus nerve mandibular branch'[Text Word])) OR ('anterior loop'[Text
Word])) OR ('inferior alveolar nerve'[Text Word])) OR ('nervus alveolaris inferior'[Text Word])) AND (((((((((((CBCT[Text Word]) OR
('cone beam computed tomography'[Text Word])) OR ('cone beam ct'[Text Word])) OR ('cone beam computed tomography'[Text Word]))
OR ('cone beam computerized tomography'[Text Word])) OR ('cone-beam computed tomography'[Text Word])) OR ('spiral cone-beam
computed tomography'[Text Word])) OR ('volume ct'[Text Word])) OR ('volume computed tomography'[Text Word])) OR ('volumetric
ct'[Text Word])) OR ('volumetric computed tomography'[Text Word])) OR ('three dimensional imaging'[Text Word])) OR ("Cone-Beam
Computed Tomography"[Mesh]))
    
```

or maintaining an average safety distance, which may vary among different individuals or ethnicities.^{8,9} Currently, the most accurate available modality for dentists to quantitatively determine the presence of AL is high-resolution CBCT.¹⁰ According to the literature, major variations have been reported in the prevalence and length of the AL; its prevalence is reported to range from 7.7%¹¹ to 95.2%¹² and its length from 0.25 mm¹³ to 19 mm.¹¹ It has been suggested that sex, ethnicity, and age differences may exist;^{8,9,14} therefore, evaluating the prevalence and length of AL and determining a precise and safe distance from it in different population subgroups can be of clinical significance.

To the authors’ best knowledge, a considerable number of publications exist on this topic and a meta-analysis has previously been conducted on the overall prevalence of AL.¹⁵ However, no meta-analysis has assessed the prevalence and length of the AL according to sex, continent, and side of the mandible, or while taking into account the effect of imaging-related factors such as various types of CBCT-reconstruction images, voxel sizes, and fields of view at the side- and patient-based levels. The current systematic review and meta-analysis aimed to assess the prevalence and extension of the AL of the mental nerve in terms of different population-based characteristics and imaging settings in CBCT images.

Materials and Methods

The present study adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.¹⁶ The study protocol was registered and published in the International Prospective Register of Systematic Reviews (PROSPERO) (registration ID: CRD42020195984). The study question was: “What is the prevalence and extension of AL of the mental nerve in terms of different population characteristics and CBCT imaging settings?”

Article screening was performed using the following

eligibility criteria. The inclusion criteria were defined in the CoCoPop frame:¹⁷ 1) condition: prevalence and extension of the AL of the mental nerve in terms of different population characteristics and CBCT imaging settings; 2) context: CBCT-reconstruction radiographs; and 3) population: patients from whom CBCT images were taken for different therapeutic purposes.

All animal studies, abstracts, unpublished articles, reviews, and studies conducted on cadavers or patients with systemic diseases that could affect bone metabolism, anatomic abnormalities, or a history of trauma in the mandible were excluded. Articles reporting opposing and contradictory results in different parts of their texts were also excluded. No language restrictions were imposed.

A thorough search was executed (by MH) in Medline/PubMed, Web of Science, Scopus, Embase, and ProQuest Dissertations & Theses for articles published up to November 23, 2020. The search strategies were designed using modifications or the combinations of the free keywords and those obtained from the MeSH database of PubMed and the Emtree database of Embase through Boolean operators (AND, OR). The search strategy in the PubMed database is presented in Table 1. Google Scholar and the Open Gray database were also searched for existing gray literature relevant to the study’s topic. Studies included in similar published systematic reviews and the references of the included articles in the present study were also searched to find relevant articles.

Two reviewers (MH and LG) independently carried out the screening process in 2 steps. First, some of the articles were excluded based on titles and abstracts. Next, the full texts of articles were thoroughly reviewed, and the articles that did not meet the eligibility criteria were ruled out. In case of disagreement, the third author (NE) was consulted to reach a consensus.

An extraction table was designed by assessing 15 studies included in the pilot phase of the extraction stage. After an

Table 2. Characteristics of studies reporting the prevalence of the anterior loop

Author	Country	Study population (patient/side)	Total population		Side	Sex (M/F)	
			Patient-based	Side-based	Right/left/bilateral	Patient-based	Side-based
Velasco-Torres et al. ¹	Spain	348/696	202	403*	209/194/-	-	-
Vujanovic-Eskenazi et al. ¹⁰	Spain	82/-	40	-	-	-	-
de Brito et al. ¹¹	Brazil	90/180	7	26	-	-	-
Haghanifar et al. ¹²	Iran	207/-	197	-	-	-	-
do Nascimento et al. ¹³	Brazil	250/500	135	208	29/33/73	-	92/116
Apostolakis and Brown ¹⁴	UK	93/186	-	91	-	-	-
Xie et al. ²¹	China	1008/2016	147	212	41/41/65	-	-
Yang et al. ²²	China	412/824	-	771	-	-	312/459
Sinha et al. ²³	India	1000/-	97	131	36/27/34	58/39	-
do Carmo Oliveira et al. ²⁴	Brazil	202/404	48	96*	50*/46*/-	-	40*/56*
Phraisukwisarn et al. ²⁵	Thailand	219/250	38*	161	-/-/31	-	64/97
Lu et al. ²⁶	US	366/732	-	624	314/310/-	-	314/310
Chen et al. ²⁷	US/Taiwan	200/-	-	-	-	-	-
von Arx et al. ²⁸	Switzerland	142/167	100	117	-	-	-
Krishnan et al. ²⁹	Australia	109/-	52	-	-	-	-
Vieira et al. ³⁰	Brazil	240/480	-	49	20/29/-	-	12/37
Shalash et al. ³¹	Egypt	120/-	66	-	-	-	-
Abidullah et al. ³²	India	120/240	24	24	11/13/-	10/14	-
Genú et al. ³³	Brazil	143/286	27	37*	8/9/10	13/14	-
Siddiqui et al. ³⁴	India	193/386	72	72	34/38/-	42/30	42/30
Panjnoush et al. ³⁵	Iran	200/400	119	197	22/19/78	59/60	-
Karnasuta et al. ³⁶	Thailand	248/441	184	368*	198/170/-	74/110	162/206
Shokry et al. ³⁷	Saudi Arabia	67/-	59	-	-	-	-
Sakhdari et al. ³⁸	Iran	200/400	38	45*	16/15/7	-	-
do Couto-Filho et al. ³⁹	Chile	47/94	14	17*	7/4/3	-	-
Chen et al. ⁴⁰	China	60/-	23	-	-	-	-
Ritter et al. ⁴¹	Germany	1010/-	313	-	-	174/137	-
Raju et al. ⁴²	United States	124/248	31	59	-/-/28	13/18	-
Prakash et al. ⁴³	India	90/-	45	-	-	15/30	-
Puri et al. ⁴⁴	India	80/-	42	-	-	-	-
Kastala et al. ⁴⁵	India	90/180	-	102	52/50/-	-	59*/43*
Chibber et al. ⁴⁶	India	60/-	32	-	-	-	-
Eren et al. ⁴⁷	Turkey	141/282	-	242	-	44/66	-
Dhumad and Saliem ⁴⁸	Iraq	90/180	27	65*	31/34/-	-	31/34
Demir et al. ⁴⁹	Turkey	279/558	-	332	168/164/-	167/165	-
Filo et al. ⁵⁰	Switzerland	694/1384	-	965	494/471/-	-	476/489
Wei et al. ⁵¹	China	306/612	251*	415	214/201/-	-	195/220
Rodricks et al. ⁵²	India	200/400	115	128	102/74/115	102/74	-
Tofiño-Medina et al. ⁵³	Peru	80/-	-	-	-	-	-
Lee et al. ⁵⁴	South Korea	20/-	-	-	-	-	-
Dos Santos Oliveira et al. ⁵⁵	Brazil	174/348	-	240	120/120/-	-	-
Chappidi et al. ⁵⁶	India	250/500	-	100	38/62/24	-	59/41
Sahman and Sisman ⁵⁷	Turkey	494/988	141	217	40/25/76	-	124/93
Koivisto et al. ⁵⁸	United States	106/212	11	15	-	2/9	-
de Oliveira-Santos et al. ⁵⁹	Brazil	100/-	-	-	-	-	-
Kung et al. ⁶⁰	Taiwan	215/-	-	-	-	-	-

Table 2. Continued

Author	Country	Study population (patient/side)	Total population		Side	Sex (M/F)	
			Patient-based	Side-based	Right/left/bilateral	Patient-based	Side-based
Alyami et al. ⁶¹	Saudi Arabia	149/298	-	37	20/17/-	-	-
Sridhar et al. ⁶²	India	146/292	26	28	11*/13*/2*	15/11	11*/17*
Goller Bulut and Köse ⁶³	Turkey	48/-	-	-	-	-	-
Kheir and Sheikhi ⁶⁴	Iran	180/360	59	122	17*/21*/42	-	63*/59*
Al-Mahalawy et al. ⁶⁵	Saudi Arabia	302/604	-	92	50/42/-	-	-
J et al. ⁶⁶	India	85/140	10	14	3/3/4	-	-
Roshene and Kumar ⁶⁷	India	30/-	-	-	-	-	-
Rosa et al. ⁶⁹	Brazil	326/-	-	-	-	-	-
Kajan and Salari ⁷⁸	Iran	84/168	-	62	33/29/-	21/19	-
Parnia et al. ⁷⁹	Iran	96/192	81	140	80/60/-	-	-
Marieiro et al. ⁸⁰	Brazil	82/-	11	-	-	-	-
Eachempati et al. ⁸¹	Malaysia	99/198	-	60*	37/23/-	-	-
Moghddam et al. ⁸²	Iran	234/451	95	106	-/-/11	46/47	-
Gala et al. ⁸³	India	65/130	-	104	-	-	-
Wong and Patil ⁸⁴	Malaysia	100/-	94	-	-	-	-
Bosykh et al. ⁸⁵	Russia	400/800	37	74	37/37/-	-	60/14
Gupta et al. ⁸⁶	India	149/298	52	122	57/65/-	-	-

M: male, F: female

agreement was reached on the appropriateness of the extraction table, 2 authors (MH and LG) started the extraction process independently, and in cases of disagreement, the third author (NE) was consulted to reach a consensus. The extraction table reported AL prevalence based on total population, sex (male/female), side (right/left), and country, as reported in Table 2.

The Joanna Briggs Institute Critical Appraisal Checklist for Studies Reporting Prevalence Data¹⁸ was used to assess the risk of bias independently by 2 reviewers (NE and MG). Both reviewers discussed any possible disagreement and reached a consensus. After completing the risk of bias assessment form for each study, the percentage of “yes” answers was obtained and used to quantify the existing bias. Bias was categorized into “high risk” (<50%), “moderate risk” (50% to 99%), and “low risk” (100%).

Statistical analysis

The fields of view were categorized into 3 subgroups according to Barbosa et al.¹⁹ Meta-analyses were conducted (by MG) with Comprehensive Meta-Analysis (CMA) software, version 2. The prevalence of AL was considered a dichotomous variable and reported as relative risk. The length of AL was considered a continuous variable and reported as means and standard deviations. The significance

level was considered as 5%. Heterogeneity was reported through the chi-square test (Cochrane Q), the related *P*-value (significance level: $P < 0.05$), and the I^2 value. An I^2 value above 50% was considered to indicate high heterogeneity,²⁰ and if it was accompanied by design and methodological heterogeneity among the studies, a random-effects model was implemented. The heterogeneity Q-test was used to compare the subgroups through the amount of overlap existing among point estimates and their 95% confidence intervals. To decrease heterogeneity, according to the information in the literature, the prevalence and length data were subgrouped based on sex (male/female), side (right/left), continent, the type of CBCT-reconstruction images, voxel size, and field of view.

Results

Study selection

The search process resulted in 2,645 articles. After excluding duplications, 1,699 studies remained. No study was found from other sources. Following the first phase of screening, 1,599 studies were ruled out due to irrelevant titles or abstracts. A number of 70 studies went through a complete full-text evaluation, in which 7 were excluded and 63 articles met the eligibility criteria. The flow chart




Fig. 1. PRISMA flow chart.

in Figure 1 presents the screening process according to the PRISMA guideline.

Study characteristics

An extraction table containing the characteristics of the included articles is presented in Table 2. Sixty-three stud-

ies with 13,743 participants (27,075 sides) satisfied the eligibility criteria and were ultimately included. Fifty-eight studies reported their sample sizes (12,256 participants) in terms of sex (5,829 males and 6,427 females). The distribution of countries in which the studies were conducted was as follows: 14 studies in India, 9 studies in Brazil, 7 studies




Fig. 2. Risk of bias assessment summary.

Table 3. Anterior loop length based on CBCT-reconstruction image types analyzed using a random-effects model

Subgroup	Study name	Mean	Standard error	Variance	95% confidence interval	P-value
Cross-sectional	do Nascimento et al. ¹³	1.060	0.055	0.003	0.951-1.169	<0.05
	Apostolakis and Brown ¹⁴	0.890	0.123	0.015	0.650-1.130	<0.05
	Yang et al. ²²	2.530	0.046	0.002	2.440-2.620	<0.05
	Phraisukwisarn et al. ²⁵	2.160	0.095	0.009	1.975-2.345	<0.05
	Siddiqui et al. ³⁴	3.660	0.235	0.055	3.200-4.120	<0.05
	Siddiqui et al. ³⁴	1.160	0.033	0.001	1.096-1.224	<0.05
	Filo et al. ⁵⁰	0.440	0.044	0.002	0.353-0.527	<0.05
	Rodricks et al. ⁵²	2.790	0.158	0.025	2.481-3.099	<0.05
	Christopher et al. ⁶⁶	2.780	0.103	0.011	2.578-2.982	<0.05
Eachempati et al. ⁸¹						
Total	1.927	0.300	0.090	1.339-2.515	<0.05	
Panoramic	do Carmo Oliveira et al. ²⁴	2.200	0.105	0.011	1.994-2.406	<0.05
	Genu et al. ³³	3.140	0.205	0.042	2.737-3.543	<0.05
	Eren et al. ⁴⁷	3.150	0.110	0.012	2.935-3.365	<0.05
	Dhumad and Saliem ⁴⁸	3.140	0.114	0.013	2.916-3.364	<0.05
	Total	2.901	0.265	0.070	2.382-3.420	<0.05
Other	Velasco-Torres et al. ¹	1.960	0.049	0.002	1.864-2.056	<0.05
	Lu et al. ²⁶	1.460	0.050	0.003	1.362-1.558	<0.05
	Wei et al. ⁵¹	3.300	0.059	0.003	3.185-3.415	<0.05
	Sahman and Sisman ⁵⁷	2.130	0.064	0.004	2.005-2.255	<0.05
	Sridhar et al. ⁶²	1.560	0.110	0.012	1.345-1.775	<0.05
	Sridhar et al. ⁶²	2.510	0.124	0.015	2.267-2.753	<0.05
	Kheir and Sheikhi ⁶⁴	2.770	0.152	0.023	2.473-3.067	<0.05
	Moghddam et al. ⁸²					
Total	2.239	0.274	0.075	1.702-2.776	<0.05	

in Iran, and 4 studies in each of the 3 countries of China, the United States, and Turkey. There were fewer than 4 studies conducted in other countries.

The countries with the largest sample sizes were India with 2,558, China with 1,786, Brazil with 1,607, Iran with 1,201, and Germany with 1,010 participants. These countries made up almost 60% of the total sample size. Other countries had sample sizes of less than 1,000.

The included studies reported the prevalence of AL in 2 ways: patient-based and/or side-based. Patient-based reporting considers a participant or a CBCT image of a complete mandible as a sample unit, whereas side-based reporting considers a hemimandible as a sample unit. For example, Xie et al.²¹ reported that there were 147 subjects with ALs (patient-based); by adding 106 right-side and 106 left-side ALs, a total of 212 ALs were detected in the hemiman-

Table 4. Pooled estimates of the prevalence and length of anterior loops within different subgroups

Subgroup		No. of studies	Point estimate (95% confidence interval)	<i>P</i> -value	<i>I</i> ² (%)	Heterogeneity <i>P</i> -value		
Patient-based prevalence	Total	-	41	40.6% (32.8-48.9)	<0.05	97.6	<0.05	
	Sex	Male	14	34.9% (25.1-46.2)	<0.05	97.6	<0.05	
		Female	14	34.5% (23.5-47.4)	<0.05	95.0	<0.05	
	Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05
	Continent	Asia	24	46.6% (34.1-59.4)	>0.05	98.11	<0.05	
		Europe	7	44.2% (28.1-61.5)	>0.05	98.21	<0.05	
		South America	6	22.4% (11.6-38.7)	<0.05	94.99	<0.05	
		North America	2	16.8% (6.7-36.2)	<0.05	87.08	<0.05	
		Africa	1	55.0% (46.0-63.7)	>0.05	-	-	
		Australia	1	47.7% (38.5-57.1)	>0.05	-	-	
		Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity						
	Side-based prevalence	Total	-	43	36.0% (27.5-45.5)	<0.05	99.18	<0.05
		Sex	Male	19	47.8% (34.9-60.9)	>0.05	98.11	<0.05
Female			19	39.7% (27.0-54.0)	>0.05	98.64	<0.05	
Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05	
Side		Right	32	32.3% (22.8-43.7)	<0.05	98.70	<0.05	
		Left	32	29.7% (21.0-40.2)	<0.05	98.60	<0.05	
Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05	
Voxel size		<0.3 mm	14	34.4% (21.6-49.8)	<0.05	99.01	<0.05	
		≥0.3 mm	9	36.0% (16.5-61.7)	>0.05	99.60	<0.05	
Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05	
Field of view		Small	8	28.2% (15.1-46.4)	<0.05	98.49	<0.05	
		Medium	7	36.0% (11.7-70.5)	>0.05	99.46	<0.05	
		Large	9	57.7% (37.6-75.5)	>0.05	99.49	<0.05	
Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05	
Anterior loop length	CBCT-reconstruction image type	Cross-sectional	9	1.92 mm (1.33-2.51)	<0.05	99.50	<0.05	
		Panoramic	4	2.90 mm (2.38-3.42)	<0.05	94.44	<0.05	
		Other planes	7	2.23 mm (1.70-2.77)	<0.05	99.06	<0.05	
		Total	20	2.39 mm (2.07-2.70)	<0.05	99.39	<0.05	
	Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							<0.05
	Side	Right	13	2.05 mm (1.56-2.53)	<0.05	98.99	<0.05	
		Left	13	1.97 mm (1.50-2.45)	<0.05	98.97	<0.05	
	Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05
	Sex	Male	11	2.13 mm (1.54-2.73)	<0.05	99.28	<0.05	
		Female	11	1.85 mm (1.35-2.36)	<0.05	99.19	<0.05	
	Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							>0.05
	Continent	Asia	11	2.50 mm (1.79-3.22)	<0.05	99.53	<0.05	
		Europe	5	1.85 mm (1.24-2.46)	<0.05	99.25	<0.05	
South America		3	2.11 mm (1.02-3.21)	<0.05	98.80	<0.05		
North America		1	1.46 mm (1.36-1.55)	<0.05	-	-		
Africa		-	-	-	-	-		
Australia		-	-	-	-	-		
Between-subgroup difference: <i>P</i> -value by Q-test for heterogeneity							<0.05	

Table 4. Continued

Subgroup	No. of studies	Point estimate (95% confidence interval)	P-value	I ² (%)	Heterogeneity P-value	
Voxel size	<0.3 mm	7	2.38 mm (1.84-2.93)	<0.05	98.82	<0.05
	≥0.3 mm	5	1.58 mm (0.88-2.27)	<0.05	99.40	<0.05
	Between-subgroup difference: P-value by Q-test for heterogeneity					>0.05
Field of view	Small	4	2.75 mm (2.18-3.32)	<0.05	94.20	<0.05
	Medium	2	2.02 mm (-0.19-4.23)	>0.05	99.46	<0.05
	Large	6	2.08 mm (1.24-2.92)	<0.05	99.78	<0.05
	Between-subgroup difference: P-value by Q-test for heterogeneity					>0.05

dibles (side based). In most studies, it was clear how the prevalence was reported; however, in some studies, both ways could be implied. These exceptions are discussed in detail in the meta-analysis part of the Results section.

Overall, the highest and the lowest patient-based prevalence rates of AL were reported by Haghanifar et al.¹² (95.2%) and de Brito et al.¹¹ (7.7%), respectively. The highest side-based prevalence was 93.5% , reported by Yang et al.,²² and the lowest was 6.5%, reported by Sinha et al.²³ The age range of participants was 4 to 98 years. AL lengths ranged from 0.25 mm¹³ to 19 mm.¹¹

Thirteen studies used a small FOV,^{12,24-35} 15 studies used a medium FOV,^{14,23,26,36-47} and 16 studies used a large FOV.^{1,11,21,22,44-46,48-56} Moreover, 30 studies used a voxel size of less than 0.3 mm,^{11-14,22,24,25,28-31,33-35,39-42,44,45,47,49-51,55,57-61} whereas 26 studies used voxels that were 0.3 mm or larger.^{1,10,14,21,23,26-28,36,37,40-42,45-47,50-54,58,62-65}

Risk of bias assessment

The risk of bias assessment was conducted using the Joanna Briggs Institute checklist designed for prevalence studies. The ninth question (“Was the response rate adequate, and if not, was the low response rate managed appropriately?”) was considered irrelevant for the current research, because the studies evaluated CBCT images and there was no need for participants to answer a questionnaire; therefore, we omitted this question. The overall risk of bias summary is also shown in Figure 2. Two studies^{66,67} were considered to have a high risk of bias, 54 had a moderate risk of bias, and 7 had a low risk of bias.^{13,23,24,26,34,36,60} As shown in Figure 2, the sample size calculation and randomization processes were the primary origins of bias. The first, fifth, and eighth questions had low risks, and there were some concerns regarding the fourth, sixth, and seventh questions.

Meta-analysis results

The summary results of the analyses conducted in this study are presented in Table 4. AL was detected in 40.6% (95% CI: 32.8%-48.9%, *P*<0.05) of the total population. Forty-one studies were meta-analyzed, and a high level of between-study heterogeneity was observed (I²=97.6%, *P*<0.05). The subgroup analysis results based on sex and continent are available in Table 4.

Thirty-six percent (95% CI: 27.5%-45.5%, *P*<0.05) of all sides had ALs. Forty-three articles were included in the meta-analysis, and the heterogeneity between the studies was high (I²=99.1%, *P*<0.05). Regarding the subgroup meta-analysis based on sides, some studies reported the prevalence of AL on the left side, right side, and bilaterally; however, some did not report these items separately and mixed bilateral ALs with those on the right and left. Therefore, a subgroup analysis was conducted only for the right and the left sides. If a study reported bilateral ALs, they were merged into the categories of the right and left and included in the meta-analysis.

As presented in Table 3, 20 studies were included in the AL length meta-analysis. The total mean extension of the AL was 2.23 mm (95% CI, 1.83-2.62 mm, *P*<0.05). A subgroup analysis based on the CBCT-reconstruction image type revealed that the length was 1.92 mm (95% CI, 1.33-2.51 mm, *P*<0.05) in cross-sectional images, 2.90 mm (95% CI, 2.38-3.42 mm, *P*<0.05) in CBCT-reconstruction panoramic images, and 2.23 mm (95% CI, 1.70-2.77 mm, *P*<0.05) in other planes. Tangential, axial, and lateral oblique views were included in the “other planes” subgroup. The difference between subgroups was statistically significant (*P*<0.05), and the heterogeneity for each subgroup was high.

As seen in Table 4, significant differences were found in the prevalence and length (*P*<0.05) of ALs in individuals from different continents, as well as the measured

length of ALs on different CBCT-reconstruction images ($P < 0.05$). However, there were no significant differences among other analyzed subgroups.

Discussion

Perhaps the most significant impact of CBCT has been on presurgical assessment and treatment planning in dental implant fixture placement using conventional, digitally planned, or guided surgery, which necessitates a correct depiction of vital structures such as the mandibular canal, its mental foramen, and the sinus floor in the maxilla.⁶⁸ Therefore, a precise preoperative evaluation of the mental nerve loop trajectory, incidence, and extension is essential.¹ An inaccurate inspection of the surgical site could increase the risk of inadvertent damage to the anterior loop segment. Almost 37% of patients undergoing implant placement surgery in the mandibular premolar region experienced noticeable changes for up to 2 weeks; symptoms persisted in 10%-15% of these cases.^{1,69} The present systematic review and meta-analysis summarized the available data on the prevalence and length of the AL of the mental nerve based on CBCT images through a comprehensive search and critical appraisal of existing evidence.

To the best of the authors' knowledge, 1 systematic review and meta-analysis of the characteristics of AL of the mental foramen has been previously published by Mishra et al.¹⁵ In the current study, more major databases were covered, including Embase, Web of Science, Scopus, and ProQuest, resulting in 31 additional articles (a total of 63 articles), 13,743 participants and 27,075 sides. Moreover, data relevant to imaging protocols (voxel size and field of view) was gathered to help researchers reproduce and improve future studies to examine AL and other anatomic landmarks. Furthermore, 12 subgroup analyses were performed to reveal possible differences between different subgroups and the impacts of population-based characteristics and imaging settings on prevalence and extension of AL, which may be considered in clinical applications.

According to the results, AL was seen in 40.6% of the population, including 34.9% of males and 34.5% of females. The prevalence of AL was 36.0% on both sides, 32.3% on the right side, and 29.7% on the left side. The average extension of AL was 2.39 mm in the total population, 2.13 mm in males, 1.85 mm in females, 2.05 mm on the right side, and 1.97 mm on the left side. No significant difference was found according to sex, side, voxel size, and field of view in terms of prevalence or length.

The studies included in the present review originated from 6 continents. Most studies (36) were conducted in Asia, followed by Europe (11), South America (11), North America (4), Africa, and Australia (1 study each). Interestingly, the analyses' results showed that these 6 continents were significantly different in terms of AL prevalence and length ($P < 0.05$). Geographically, the longest ALs were seen in Asia (2.50 mm), followed by South America (2.11 mm), Europe (1.85 mm), and North America (1.46 mm). However, due to the high level of between-study heterogeneity, these data should be used conservatively in clinical practice. No study evaluated the extension of ALs in Africa and Australia. A statistically significant difference in prevalence was found between South America and North America ($P < 0.05$), with a higher prevalence of AL in South America (22.4%) than in North America (16.8%). However, the prevalence of AL was not significantly different between other continents.

Most of the series of images used to evaluate the mean AL length in the present study were cross-sections or reformatted panoramic images, with a smaller group of compromised tangential planes, oblique axial planes, and 3-dimensional reconstructions. Significant differences were found between the groups ($P < 0.05$). The mean AL length in cross-sectional images was 1.92 mm, while it was 2.90 mm in reformatted panoramic images; these results showed a 1-mm difference between the 2 image types. However, most clinicians believe that CBCT images are reliable and free from distortion and are unaware of the possibility of imprecisions or inconsistencies when taking linear measurements or assessing anatomical landmarks before implant insertion.⁷⁰ A systematic review by Fokas et al. assessed the precision of linear measurements on CBCT images and reported that a range of absolute error exceeding a threshold of 1 mm could have clinical implications. Moreover, they suggested that the sub-millimeter accuracy of CBCT may be insufficient in some cases of implant surgery, potentially resulting in clinical complications.⁷⁰ In contrast, some studies have recommended a 2-mm safety zone for measurements on panoramic radiography.^{3,11} The results obtained by Fokas et al.⁷⁰ were consistent with the current results, which showed a 1 mm difference between 3 CBCT-reconstruction image types. Therefore, to interpret the results cautiously, it is suggested that clinicians should consider a 2-mm safety margin from crucial anatomical structures observed on CBCT.

The studies included in this review that used a voxel size of less than 0.3 mm did not show significant differences in the prevalence and mean length of the AL compared to

those that used voxel sizes of at least 0.3 mm. Voxel sizes ranged from 0.07 mm to 0.4 mm among 46 studies which reported this parameter. Smaller voxel sizes have diagnostic accuracy in other applications, such as endodontics and the detection of various periodontal defects;^{71,72} however, for prevalence and mean length of AL, a voxel size of 0.3-0.4 mm seems to be sufficiently precise.

The findings of this study align with those of other studies reporting no significant differences between voxel sizes (0.2, 0.3, and 0.4 mm) when evaluating linear bone measurements or implant placement.⁷³⁻⁷⁵ No significant differences were found in the prevalence or mean length of ALs depending on the type of field of view (large: > 15 cm; medium: 10-15 cm; small: ≤ 10 cm). This indicates that radiation dose reduction settings, such as a limitation of the field of view, may be applied with minimal effects on the accuracy of measurements and the diagnostic outcome.⁷⁶

The overall risk of bias was moderate in the current study. The included studies contained 2 high-risk,^{66,67} 7 low-risk,^{13,23,24,26,34,36,60} and 54 moderate-risk studies. According to the Joanna Briggs Institute risk of bias assessment checklist, the most biased aspect was the randomization process. This can be ignored because the included studies used CBCT images as samples, and no systematic error could jeopardize the results. Another field of bias in the included studies was related to the method used to determine sample size, which generally seemed more arbitrary than determined through statistical calculations and power analyses. This could affect the external validity of the obtained results. The sample size of the studies ranged from 20 participants in the study of Lee et al.⁵⁴ to 1010 in that by Ritter et al.⁴¹

Furthermore, some studies did not describe their study settings and standard measurement methods of the outcomes in detail, which could jeopardize the reproducibility and reliability of their results. For example, some articles did not report settings such as the CBCT device manufacturer, software, voxel size, or field of view. It is recommended that researchers conduct future studies in adherence to the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) statement⁷⁷ to provide a more standardized and reliable report of results.

Overall, it seems the voxel size and field of view of CBCT devices are adequate for accurately identifying the AL; however, due to potential variations between individuals, an overall 2-mm safety margin from such crucial anatomical structures is suggested to be considered when using CBCT images. The overall heterogeneity between the studies of all evaluated subgroups was high, so the results

should be used in clinical practice with caution. Moreover, the aspect of the included studies with the greatest concern for bias was the sample size calculation process, which should be considered in designing future studies.

Conflicts of Interest: None

References

1. Velasco-Torres M, Padial-Molina M, Avila-Ortiz G, García-Delgado R, Catena A, Galindo-Moreno P. Inferior alveolar nerve trajectory, mental foramen location and incidence of mental nerve anterior loop. *Med Oral Patol Oral Cir Bucal* 2017; 22: e630-5.
2. Iyengar AR, Patil S, Nagesh KS, Mehkri S, Manchanda A. Detection of anterior loop and other patterns of entry of mental nerve into the mental foramen: a radiographic study in panoramic images. *J Dent Implant* 2013; 3: 21-5.
3. Greenstein G, Tarnow D. The mental foramen and nerve: clinical and anatomical factors related to dental implant placement: a literature review. *J Periodontol* 2006; 77: 1933-43.
4. Hogg KD. Mandibular anatomical structures. In: Ho CC. *Practical procedures in implant dentistry*. Hoboken: Wiley-Blackwell; 2021. p. 59-72.
5. Balaguer-Martí JC, Peñarrocha-Oltra D, Balaguer-Martínez J, Peñarrocha-Diogo M. Immediate bleeding complications in dental implants: a systematic review. *Med Oral Patol Oral Cir Bucal* 2015; 20: e231-8.
6. von Arx T, Häfliger J, Chappuis V. Neurosensory disturbances following bone harvesting in the symphysis: a prospective clinical study. *Clin Oral Implants Res* 2005; 16: 432-9.
7. Padmanabhan H, Kumar AV, Shivashankar K. Incidence of neurosensory disturbance in mandibular implant surgery - a meta-analysis. *J Indian Prosthodont Soc* 2020; 20: 17-26.
8. Aminoshariae A, Su A, Kulild JC. Determination of the location of the mental foramen: a critical review. *J Endod* 2014; 40: 471-5.
9. Gawlikowska-Sroka A, Stocki Ł, Dąbrowski P, Kwiatkowska B, Szczurowski J, Czerwiński F. Topography of the mental foramen in human skulls originating from different time periods. *Homo* 2013; 64: 286-95.
10. Vujanovic-Eskenazi A, Valero-James JM, Sánchez-Garcés MA, Gay-Escoda C. A retrospective radiographic evaluation of the anterior loop of the mental nerve: comparison between panoramic radiography and cone beam computerized tomography. *Med Oral Patol Oral Cir Bucal* 2015; 20: e239-45.
11. de Brito AC, Nejaim Y, de Freitas DQ, de Oliveira Santos C. Panoramic radiographs underestimate extensions of the anterior loop and mandibular incisive canal. *Imaging Sci Dent* 2016; 46: 159-65.
12. Haghanifar S, Moudi E, Bijani A, Lavasani S, Lameh A. Mandibular canal and its incisive branch: a CBCT study. *World Fam Med J* 2017; 15: 133-40.
13. do Nascimento EH, Dos Anjos Pontual ML, Dos Anjos Pontual A, da Cruz Perez DE, Figueiroa JN, Frazão MA, et al. Assessment of the anterior loop of the mandibular canal: a study

- using cone-beam computed tomography. *Imaging Sci Dent* 2016; 46: 69-75.
14. Apostolakis D, Brown JE. The anterior loop of the inferior alveolar nerve: prevalence, measurement of its length and a recommendation for interforaminal implant installation based on cone beam CT imaging. *Clin Oral Implants Res* 2012; 23: 1022-30.
 15. Mishra SK, Nahar R, Gaddale R, Chowdhary R. Identification of anterior loop in different populations to avoid nerve injury during surgical procedures - a systematic review and meta-analysis. *Oral Maxillofac Surg* 2021; 25: 159-74.
 16. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71.
 17. Munn Z, Stern C, Aromataris E, Lockwood C, Jordan Z. What kind of systematic review should I conduct? A proposed typology and guidance for systematic reviewers in the medical and health sciences. *BMC Med Res Methodol* 2018; 18: 5.
 18. Munn Z, Moola S, Lisy K, Riitano D, Tufanaru C. Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data. *Int J Evid Based Healthc* 2015; 13: 147-53.
 19. Ferreira Barbosa DA, Barros ID, Teixeira RC, Menezes Pimenta AV, Kurita LM, Barros Silva PG, et al. Imaging aspects of the mandibular incisive canal: a PROSPERO-registered systematic review and meta-analysis of cone beam computed tomography studies. *Int J Oral Maxillofac Implants* 2019; 34: 423-33.
 20. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; 21: 1539-58.
 21. Xie L, Li T, Chen J, Yin D, Wang W, Xie Z. Cone-beam CT assessment of implant-related anatomy landmarks of the anterior mandible in a Chinese population. *Surg Radiol Anat* 2019; 41: 927-34.
 22. Yang XW, Zhang FF, Li YH, Wei B, Gong Y. Characteristics of intrabony nerve canals in mandibular interforaminal region by using cone-beam computed tomography and a recommendation of safe zone for implant and bone harvesting. *Clin Implant Dent Relat Res* 2017; 19: 530-8.
 23. Sinha S, Kandula S, Sangamesh NC, Rout P, Mishra S, Bajoria AA. Assessment of the anterior loop of the mandibular canal using cone-beam computed tomography in Eastern India: a record-based study. *J Int Soc Prev Community Dent* 2019; 9: 290-5.
 24. do Carmo Oliveira M, Tedesco TK, Gimenez T, Allegrini S Jr. Analysis of the frequency of visualization of morphological variations in anatomical bone features in the mandibular interforaminal region through cone-beam computed tomography. *Surg Radiol Anat* 2018; 40: 1119-31.
 25. Phraisukwisarn P, Asvanund P, Kretapirom K. Measurement of anterior loop of inferior alveolar nerve using cone beam computed tomography (CBCT). *M Dent J* 2017; 37: 81-7.
 26. Lu CI, Won J, Al-Ardah A, Santana R, Rice D, Lozada J. Assessment of the anterior loop of the mental nerve using cone beam computerized tomography scan. *J Oral Implantol* 2015; 41: 632-9.
 27. Chen JC, Lin LM, Geist JR, Chen JY, Chen CH, Chen YK. A retrospective comparison of the location and diameter of the inferior alveolar canal at the mental foramen and length of the anterior loop between American and Taiwanese cohorts using CBCT. *Surg Radiol Anat* 2013; 35: 11-8.
 28. von Arx T, Friedli M, Sendi P, Lozanoff S, Bornstein MM. Location and dimensions of the mental foramen: a radiographic analysis by using cone-beam computed tomography. *J Endod* 2013; 39: 1522-8.
 29. Krishnan U, Monsour P, Thaha K, Lalloo R, Moule A. A limited field cone-beam computed tomography-based evaluation of the mental foramen, accessory mental foramina, anterior loop, lateral lingual foramen, and lateral lingual canal. *J Endod* 2018; 44: 946-51.
 30. Vieira CL, Veloso SD, Lopes FF. Location of the course of the mandibular canal, anterior loop and accessory mental foramen through cone-beam computed tomography. *Surg Radiol Anat* 2018; 40: 1411-7.
 31. Shalash M, Khallaf ME, Ali AR. Position and dimensions of the mental foramen and presence of the anterior loop in the Egyptian population: a retrospective CBCT study. *Bull Natl Res Cent* 2020; 44: 110.
 32. Abidullah M, Kaur P, Karthikeyan P, Koppolu P, Kaur R. Assessment of different patterns of entry of mental nerve in mental foramen: a cone beam computed tomography study. *World J Dent* 2018; 9: 382-6.
 33. Genú PR, Vasconcellos RJ, Oliveira BP, Vasconcelos BC, Delgado NC. Analysis of anatomical landmarks of the mandibular interforaminal region using CBCT in a Brazilian population. *Braz J Oral Sci* 2014; 13: 303-7.
 34. Siddiqui Z, Rai S, Ranjan V. Efficacy and evaluation of cone beam computed tomography in determining the prevalence and length of anterior loop of inferior alveolar nerve in North Indian population. *J Indian Acad Oral Med Radiol* 2018; 30: 32-7.
 35. Panjnoush M, Rabiee ZS, Kheirandish Y. Assessment of location and anatomical characteristics of mental foramen, anterior loop and mandibular incisive canal using cone beam computed tomography. *J Dent (Tehran)* 2016; 13: 126-32.
 36. Karnasuta P, Plianrunsi J, Denkongpon I, Horsimasathaporn N, Chayutthanabun P, Weerachartwattana J, et al. Cone-beam computed tomography investigation of crucial mandibular canal variations in Thais. *Oral Radiol* 2017; 33: 219-26.
 37. Shokry SM, Salam ZA, Alshaib S, Al Mohaimeed ZZ. Comparing the apparent length of the anterior loop on panoramic radiographs and a new tracing technique on three-dimensional cone-beam computed tomographic images. *Int J Clin Den Sci* 2016; 7: 1-5.
 38. Sakhdari S, Hafezi L, Esmaili M. Prevalence of the inferior alveolar nerve's anterior loop and mandibular incisive canal by use of cone beam computed tomography (CBCT) in an Iranian population. *J Res Dent Maxillofac Sci* 2016; 1: 14-21.
 39. do Couto-Filho CE, de Moraes PH, Alonso MB, Haiter-Neto F, Olate S, de Albergaria-Barbosa JR. Accuracy in the diagnosis of the mental nerve loop. a comparative study between panoramic radiography and cone beam computed tomography. *Int J Morphol* 2015; 33: 327-32.
 40. Chen Z, Chen D, Tang L, Wang F. Relationship between the position of the mental foramen and the anterior loop of the inferior alveolar nerve as determined by cone beam computed tomography combined with mimics. *J Comput Assist Tomogr*

- 2015; 39: 86-93.
41. Ritter L, Neugebauer J, Mischkowski RA, Dreiseidler T, Rothamel D, Richter U, et al. Evaluation of the course of the inferior alveolar nerve in the mental foramen by cone beam computed tomography. *Int J Oral Maxillofac Implants* 2012; 27: 1014-21.
 42. Raju N, Zhang W, Jadhav A, Ioannou A, Eswaran S, Weltman R. Cone-beam computed tomography analysis of the prevalence, length, and passage of the anterior loop of the mandibular canal. *J Oral Implantol* 2019; 45: 463-8.
 43. Prakash O, Srivastava PK, Jyoti B, Mushtaq R, Vyas T, Usha P. Radiographic evaluation of anterior loop of inferior alveolar nerve: a cone-beam computer tomography study. *Niger J Surg* 2018; 24: 90-4.
 44. Puri A, Verma P, Mahejan P, Bansal A, Kohli S, Faraz S. CBCT evaluation of the vital mandibular interforaminal anatomical structures. *Ann Maxillofac Surg* 2020; 10: 149-57.
 45. Kastala RK, David CM, Jayapal N. Momentousness of the mental loop: a comparative study. *Contemp Clin Dent* 2019; 10: 86-92.
 46. Chibber N, Singh S, Sudan S, Anand M. CBCT use to evaluate vital mandibular inter-foramin anatomical structures. *Indian J Public Health Res Dev* 2020; 11: 650-4.
 47. Eren H, Orhan K, Bagis N, Nalcaci R, Misirli M, Hincal E. Cone beam computed tomography evaluation of mandibular canal anterior loop morphology and volume in a group of Turkish patients. *Biotechnol Biotechnol Equip* 2016; 30: 346-53.
 48. Dhumad OS, Saliem SS. Anterior loop presence and extension using cone-beam computed tomography. *Indian J Public Health Res Dev* 2019; 10: 764-7.
 49. Demir A, Izgi E, Pekiner F. Anterior loop of the mental foramen in a Turkish subpopulation with dentate patients: a cone beam computed tomography study. *J Marmara Univ Inst Health Sci* 2015; 5: 231-8.
 50. Filo K, Schneider T, Locher MC, Kruse AL, Lübbers HT. The inferior alveolar nerve's loop at the mental foramen and its implications for surgery. *J Am Dent Assoc* 2014; 145: 260-9.
 51. Wei X, Gu P, Hao Y, Wang J. Detection and characterization of anterior loop, accessory mental foramen, and lateral lingual foramen by using cone beam computed tomography. *J Prosthet Dent* 2020; 124: 365-71.
 52. Rodricks D, Phulambrikar T, Singh S, Gupta A. Evaluation of incidence of mental nerve loop in Central India population using cone beam computed tomography. *Indian J Dent Res* 2018; 29: 627-33.
 53. Tofiño-Medina JH, Arriola-Guillén LE, Rodríguez-Cárdenas YA, Aliaga-Del Castillo A, Ruíz-Mora GA, Guerrero ME. Frequency of accessory mental foramen and anatomical variability of mental nerve anterior loop in a Peruvian population: a retrospective cone-beam computed tomography study. *J Oral Res* 2020; 9: 202-11.
 54. Lee JE, Lee YJ, Jin SH, Kim Y, Kook YA, Ko Y, et al. Topographic analysis of the mandibular symphysis in a normal occlusion population using cone-beam computed tomography. *Exp Ther Med* 2015; 10: 2150-6.
 55. Dos Santos Oliveira R, Maria Gomes Oliveira A, Cintra Junqueira JL, Kühn Panzarella F. Association between the anatomy of the mandibular canal and facial types: a cone-beam computed tomography analysis. *Int J Dent* 2018; 2018: 5481383.
 56. Chappidi V, Swapna L, Dheeraj V, Nikitha G, Kanakagiri M. Evaluation of morphometric variations in mental foramen and prevalence of anterior loop in South Indian population - a CBCT study. *J Indian Acad Oral Med Radiol* 2019; 31: 134-9.
 57. Sahman H, Sisman Y. Anterior loop of the inferior alveolar canal: a cone-beam computerized tomography study of 494 cases. *J Oral Implantol* 2016; 42: 333-6.
 58. Koivisto T, Chiona D, Milroy LL, McClanahan SB, Ahmad M, Bowles WR. Mandibular canal location: cone-beam computed tomography examination. *J Endod* 2016; 42: 1018-21.
 59. de Oliveira-Santos C, Souza PH, de Azambuja Berti-Couto S, Stinkens L, Moyaert K, Rubira-Bullen IR, et al. Assessment of variations of the mandibular canal through cone beam computed tomography. *Clin Oral Investig* 2012; 16: 387-93.
 60. Kung CY, Wang YM, Chan CP, Ju YR, Pan WL. Evaluation of the mandibular lingual canal and anterior loop length to minimize complications associated with anterior mandibular surgeries: a cone-beam computed tomography study. *J Oral Maxillofac Surg* 2017; 75: 2116.e1-13.
 61. Alyami OS, Alotaibi MS, Koppolu P, Alosaimy A, Abdulghani A, Swapna LA, et al. Anterior loop of the mental nerve in Saudi sample in Riyadh, KSA. A cone beam computerized tomography study. *Saudi Dent J* 2021; 33: 124-30.
 62. Sridhar M, Dhanraj M, Nesappan T, Jain A. A retrospective radiographic evaluation of incisive canal and anterior loop of mental nerve using cone beam computed tomography. *Drug Invent Today* 2018; 10: 1656-60.
 63. Goller Bulut D, Köse E. Available bone morphology and status of neural structures in the mandibular interforaminal region: three-dimensional analysis of anatomical structures. *Surg Radiol Anat* 2018; 40: 1243-52.
 64. Kheir MK, Sheikhi M. Assessment of the anterior loop of mental nerve in an Iranian population using cone beam computed tomography scan. *Dent Res J (Isfahan)* 2017; 14: 418-22.
 65. Al-Mahalawy H, Al-Aithan H, Al-Kari B, Al-Jandan B, Shujaat S. Determination of the position of mental foramen and frequency of anterior loop in Saudi population. A retrospective CBCT study. *Saudi Dent J* 2017; 29: 29-35.
 66. J PC, Marimuthu T, C K, Devadoss P, Kumar SM. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: a cross sectional study. *Clin Implant Dent Relat Res* 2018; 20: 531-4.
 67. Roshene R, Kumar J. Radiological study on the anterior loop of inferior alveolar nerve. *Indian J Public Health Res Develop* 2019; 10: 595-9.
 68. Freire-Maia B, Machado VD, Valerio CS, Custódio AL, Manzi FR, Junqueira JL. Evaluation of the accuracy of linear measurements on multi-slice and cone beam computed tomography scans to detect the mandibular canal during bilateral sagittal split osteotomy of the mandible. *Int J Oral Maxillofac Surg* 2017; 46: 296-302.
 69. Rosa MB, Sotto-Maior BS, Machado Vde C, Francischone CE. Retrospective study of the anterior loop of the inferior alveolar nerve and the incisive canal using cone beam computed tomography. *Int J Oral Maxillofac Implants* 2013; 28: 388-92.
 70. Fokas G, Vaughn VM, Scarfe WC, Bornstein MM. Accuracy

- of linear measurements on CBCT images related to presurgical implant treatment planning: a systematic review. *Clin Oral Implants Res* 2018; 29 Suppl 16: 393-415.
71. Icen M, Orhan K, Şeker Ç, Geduk G, Cakmak Özlü F, Cengiz M. Comparison of CBCT with different voxel sizes and intra-oral scanner for detection of periodontal defects: an in vitro study. *Dentomaxillofac Radiol* 2020; 49: 20190197.
 72. Kamburoğlu K, Kursun S. A comparison of the diagnostic accuracy of CBCT images of different voxel resolutions used to detect simulated small internal resorption cavities. *Int Endod J* 2010; 43: 798-807.
 73. Damstra J, Fourie Z, Huddleston Slater JJ, Ren Y. Accuracy of linear measurements from cone-beam computed tomography-derived surface models of different voxel sizes. *Am J Orthod Dentofacial Orthop* 2010; 137: 16.e1-6.
 74. Mukhia N, Birur NP, Shubhasini AR, Shubha G, Keerthi G. Dimensional measurement accuracy of 3-dimensional models from cone beam computed tomography using different voxel sizes. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2021; 132: 361-9.
 75. Torres MG, Campos PS, Segundo NP, Navarro M, Crusoé-Rebello I. Accuracy of linear measurements in cone beam computed tomography with different voxel sizes. *Implant Dent* 2012; 21: 150-5.
 76. Ganguly R, Ramesh A, Pagni S. The accuracy of linear measurements of maxillary and mandibular edentulous sites in cone-beam computed tomography images with different fields of view and voxel sizes under simulated clinical conditions. *Imaging Sci Dent* 2016; 46: 93-101.
 77. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008; 61: 344-9.
 78. Kajan ZD, Salari A. Presence and course of the mandibular incisive canal and presence of the anterior loop in cone beam computed tomography images of an Iranian population. *Oral Radiol* 2012; 28: 55-61.
 79. Parnia F, Moslehifard E, Hafezeqoran A, Mahboub F, Mojaver-Kahnemoui H. Characteristics of anatomical landmarks in the mandibular interforaminal region: a cone-beam computed tomography study. *Med Oral Patol Oral Cir Bucal* 2012; 17: e420-5.
 80. Marieiro LM, Deluiz D, Ferreira DC, Tannure PN, editors. Measurement of distance between the mental foramina using cone-beam computed tomography: a pilot study with a possible method for planning mandibular implants. *Pesqui Bras Odontopediatria Clin Integr* 2017; 17: e3328.
 81. Eachempati P, Nagraj SK, Chien DC, Lin CC, Soe HH, KS KK. Radiographic evaluation of the anterior loop of the mental nerve: comparison between orthopantomograph and cone-beam CT - a pilot study. *Res J Pharm Biol Chem Sci* 2017; 8: 2044-50.
 82. Moghddam MR, Davoudmanesh Z, Azizi N, Rakhshan V, Shariati M. Prevalence and length of the anterior loop of the inferior alveolar nerve in Iranians. *J Oral Implantol* 2017; 43: 333-6.
 83. Gala V, Tirpude V, Shah D, Doshi A, Fernandes G. A retrospective cone beam computed tomography (CBCT) study of the assessment of the length of the anterior loop of the inferior alveolar nerve. *Int J S Res Sci Tech* 2018; 4: 828-32.
 84. Wong SK, Patil PG. Measuring anterior loop length of the inferior alveolar nerve to estimate safe zone in implant planning: a CBCT study in a Malaysian population. *J Prosthet Dent* 2018; 120: 210-3.
 85. Bosykh YY, Turkina AY, Franco RP, Franco A, Makeeva MK. Cone beam computed tomography study on the relation between mental foramen and roots of mandibular teeth, presence of anterior loop and satellite foramina. *Morphologie* 2019; 103: 65-71.
 86. Gupta A, Pubreja L, Malik R, Gupta N. Evaluation of the dimensions of anterior loop of mental nerve in CBCT: a radiographic analysis. *J Maxillofac Oral Surg* 2020; 19: 168-72.