

RESEARCH ARTICLE

Open Access

Characterization of the *MLO* gene family in Rosaceae and gene expression analysis in *Malus domestica*

Stefano Pessina^{1,2}, Stefano Pavan³, Domenico Catalano⁴, Alessandra Gallotta³, Richard GF Visser², Yuling Bai², Mickael Malnoy^{1*} and Henk J Schouten^{2*}

Abstract

Background: Powdery mildew (PM) is a major fungal disease of thousands of plant species, including many cultivated Rosaceae. PM pathogenesis is associated with up-regulation of *MLO* genes during early stages of infection, causing down-regulation of plant defense pathways. Specific members of the *MLO* gene family act as PM-susceptibility genes, as their loss-of-function mutations grant durable and broad-spectrum resistance.

Results: We carried out a genome-wide characterization of the *MLO* gene family in apple, peach and strawberry, and we isolated apricot *MLO* homologs through a PCR-approach. Evolutionary relationships between *MLO* homologs were studied and syntenic blocks constructed. Homologs that are candidates for being PM susceptibility genes were inferred by phylogenetic relationships with functionally characterized *MLO* genes and, in apple, by monitoring their expression following inoculation with the PM causal pathogen *Podosphaera leucotricha*.

Conclusions: Genomic tools available for Rosaceae were exploited in order to characterize the *MLO* gene family. Candidate *MLO* susceptibility genes were identified. In follow-up studies it can be investigated whether silencing or a loss-of-function mutations in one or more of these candidate genes leads to PM resistance.

Keywords: Rosaceae, *MLO*, Powdery mildew, *Malus domestica*

Background

Powdery mildew (PM) is a major fungal disease for thousands of plant species [1], including cultivated Rosaceae such as apple (*Malus domestica*), apricot (*Prunus armeniaca*), peach (*Prunus persica*), and strawberry (*Fragaria x ananassa*). Powdery mildew occurs in all major growing regions of Rosaceous crops, leading to severe losses [2]. The major PM causal agents are *Podosphaera leucotricha* in apple [2], *Sphaerotheca pannosa* var. *persicae* in peach [3], *Podosphaera tridactyla* in apricot [4] and *Podosphaera aphanis* (syn. *Sphaerotheca macularis* f. sp. *fragariae*) in strawberry [5]. Powdery mildew shows similar symptoms in the four species: white spots appear on young green tissues, particularly leaves in the first days after opening,

whereas mature leaves show some resistance. Infected leaves crinkle, curl, and prematurely drop. Blossoms and fruits are not the primary targets of PM fungi, but infections of these tissues are possible [2,3,5]. In peach, apricot and apple, PM spores overwinter in buds and then in spring, with the reprise of vegetative growth, the spores start a new infection [2,3].

Cultivars resistant to PM are fundamental in order to reduce the use of pesticides in agricultural practices. The usual strategy in breeding focuses on dominant plant resistance genes (R-genes), however these genes often originate from wild-relatives of the cultivated species, and thus interspecific crossability barriers could prevent their introgression [6]. Moreover, in case of a successful cross, several undesirable traits are incorporated with the R-gene, making extensive backcrossing necessary, which is time-consuming in woody species. Finally, the durability of R-genes is generally limited due to the appearance of virulent strains of the pathogen, which can

* Correspondence: mickael.malnoy@fmach.it; henk.schouten@wur.nl

¹Department of Genomics and Biology of Fruit Crops, Fondazione Edmund Mach, via E. Mach 1, 38010 San Michele all'Adige, Italy

²Wageningen UR Plant Breeding, Wageningen University and Research Centre, P.O. Box 16, 6700 AA Wageningen, The Netherlands

Full list of author information is available at the end of the article

overcome resistance in a few years [7]. Two examples are *Venturia inaequalis* race 6, which is able to overcome *Rvi6* resistance to scab in apple [8], and *P. leucotricha* strains able to breakdown *Pl-1* and *Pl-2*, two major PM R-genes of apple [9].

An alternative to the use of R-genes is based on plant susceptibility genes (S-genes), defined as genes whose loss-of-function results in recessively inherited resistance [10]. Barley *mlo* PM resistance, first characterized in 1942, is a remarkable example of immunity due to the absence of an S-gene, as it derives from a loss-of-function mutation in the *MLO* (*Mildew Locus O*) gene, encoding for a protein with seven transmembrane domains [11,12]. *mlo* resistance has long been considered as a unique form of immunity, characterized by durability, broad-spectrum effectiveness and recessive inheritance [13]. However, the characterization of the sources of resistance in other plant species, such as *Arabidopsis* [14], pea [15,16] and tomato [17], has revealed that resistance resulting from loss-of-function mutations in *MLO* functional orthologs is more common than previously thought. Therefore, it has been suggested that the inactivation of *MLO* susceptibility genes could represent a valid strategy to introduce PM resistance across a broad range of cultivated species [10].

Histological characterization of *mlo* resistance revealed that it is based on a pre-penetration defense system, associated with the formation of cell-wall appositions [14,18] and at least partially dependent on the actin cytoskeleton [19]. It has been suggested that functional *MLO* proteins negatively regulate vesicle-associated and actin-dependent defense pathways at the site of attempted PM penetration [20]. *MLO* proteins are therefore targeted by PM fungi as a strategy to induce pathogenesis. The early stages of PM infection are associated with an increase in transcript abundance of *MLO* susceptibility genes, showing a peak at six hours after inoculation. This has been shown to occur in tomato [17], barley [21], pepper [22] and grape [23,24].

MLO susceptibility genes are members of a gene family which shows tissue specific expression patterns and are involved in a variety of physiological processes, besides the response to PM fungi: one of the 15 *MLO* genes of *Arabidopsis*, *AtMLO7*, is involved in pollen tube reception by the embryo sac and its mutation results in reduced fertility [25]. Two other *Arabidopsis* genes, named *AtMLO4* and *AtMLO11*, are involved in the control of root architecture, as mutants with null alleles of these two genes display asymmetrical root growth and exaggerated curvature [26].

Previous phylogenetic analysis of the *MLO* protein family identified six clades [23]. Currently, all *MLO* proteins functionally related to PM susceptibility in dicot species appear in a single clade, namely Clade V [14,17,23,24]. Similarly, Clade IV harbours all characterized PM susceptibility proteins from monocots [20,27].

MLO genes have been intensively studied in many monocots and dicots, but very little has been performed in Rosaceae. In this investigation, we characterized the *MLO* gene family in a number of Rosaceous species, with respect to their structural, genomic and evolutionary features. Moreover, we monitored the transcript abundances of apple *MLO* homologs following *P. leucotricha* inoculation in three apple cultivars.

Results

In silico and *in vitro* characterization of Rosaceae

MLO homologs

A database search for Rosaceae *MLO* homologs produced 21 significant matches in peach, 23 in strawberry and 28 in apple. Of these, six (five from *M. domestica* and one from *F. vesca*) showed a very limited alignment region with other *MLO* genes, whereas eight (two from *M. domestica*, two from *P. persica* and four from *F. vesca*) were characterized by markedly different length with respect to *MLO* homologs reported in the genomes of *Arabidopsis* and grapevine [23,28], i.e. less than 350 amino acids (aa) or more than 700 aa. Details on genomic localization amino acid number, putative transmembrane domains and predicted exon/intron structure of the remaining homologs, together with information about the *MLO* homologs nomenclature chosen in this study is provided in Tables 1, 2 and 3.

Peach and apricot are evolutionary very close to each other, and show a high degree of homology in DNA sequence. Phylogenetic analysis (see next paragraph) indicated peach homologs *PpMLO1*, *PpMLO3* and *PpMLO4* as candidates for being required for PM susceptibility. Therefore, we used the sequences of these genes to design primers to identify full-length apricot *MLO* genes. This approach resulted in the amplification and the successive characterization of three *MLO* sequences, which were by analogy named *PaMLO1*, *PaMLO3*, and *PaMLO4* (deposited in the NCBI database with the accession numbers KF177395, KF177396, and KF177397, respectively).

Phylogenetic relations and inference of orthology

We performed a phylogenetic study on the newly identified Rosaceae *MLO* proteins. The dataset was completed with four homologs recently characterized in *Rosa hybrida* (rose) [30] (RhMLO1, RhMLO2, RhMLO3 and RhMLO4), the complete *Arabidopsis thaliana* *AtMLO* protein family [14], a series of *MLO* homologs which have been functionally associated with PM susceptibility, namely tomato (*Solanum lycopersicum*) SIMLO1 [17], pea (*Pisum sativum*) PsMLO1 [15,16], pepper (*Capsicum annuum*) CaMLO2 [27], lotus (*Lotus japonicus*) LjMLO1 [15], barrel clover (*Medicago truncatula*) MtMLO1 [15], barley (*Hordeum vulgare*) HvMLO [11], rice (*Oryza sativa*) OsMLO2 [31], wheat (*Triticum aestivum*) TaMLO_B1 and TaMLO_A1b [31], and grapevine (*Vitis vinifera*) VvMLO14,

Table 1 Members of the MdMLO gene family as predicted in *M. domestica* cv. Golden Delicious genome sequence

Gene	Accession number ^a	Chr.	Starting position (Mb)	Clade	Introns	TM ^b	Amino acids	Conserved aa ^c
MdMLO1	MDP0000177099	2	1.02	II	11	3	487	25
MdMLO2	MDP0000240125	2	11.10	I	11	3	571	20
MdMLO3	MDP0000168575	2	11.11	I	13	7	670	22
MdMLO4	MDP0000207002	2	8.79	III	16	7	634	28
MdMLO5	MDP0000163089	9	15.26	V	14	6	579	30
MdMLO6	MDP0000119433	3	33.95	II	0	7	504	30
MdMLO7	MDP0000123907	n.d.	n.d.	V	n.d.	6	561	28
MdMLO8	MDP0000218520	2	11.11	I	9	4	390	14
MdMLO9	MDP0000320797	2	27.20	II	10	5	454	28
MdMLO10	MDP0000196373	3	26.97	I	13	6	539	28
MdMLO11	MDP0000239643	4	9.84	V	12	8	575	28
MdMLO12	MDP0000133162	6	0.81	III	13	5	516	28
MdMLO13	MDP0000142608	7	7.48	II	12	6	351	18
MdMLO14	MDP0000191469	8	29.25	II	10	5	395	23
MdMLO15	MDP0000141595	9	7.54	III	15	6	647	24
MdMLO16	MDP0000191848	9	21.12	VI	14	6	606	29
MdMLO17	MDP0000145097	11	27.97	I	13	7	523	28
MdMLO18	MDP0000928368	10	27.97	VII	12	7	502	30
MdMLO19	MDP0000168714	12	16.23	V	13	7	590	30
MdMLO20	MDP0000134649	13	11.61	VIII	13	5	589	27
MdMLO21	MDP0000133760	15	24.99	VI	15	6	560	28

^aAvailable at http://www.rosaceae.org/gb/gbrowse/malus_x_domestica/.

^bNumber of transmembrane domains in the predicted protein, as determined by InterPro prediction software (<http://www.ebi.ac.uk/interpro/>).

^cnumber of conserved amino acids out of the 30 identified by Elliot et al. [29].

the only dicot MLO homolog known to belong to clade IV [23]. Clustering analysis using the UPGMA algorithm resulted in a total of eight distinct clades and no divergent lineage (Figure 1). Clade numbers from I to VI were assigned based on the position of *Arabidopsis* AtMLO homologs and barley HvMLO, according to the previous study of Feechan et al. [23]. The two additional clades (named VII and VIII) were found to include Rosaceae MLO homologs only, both having one homolog from *P. persica*, one from *F. vesca* and one from *M. domestica*. Further clustering analysis with a Neighbour-Joining algorithm resulted in merging clade VII and VIII (not shown).

Four apple MLO homologs (MdMLO5, MdMLO7, MdMLO11 and MdMLO19) and three MLO homologs from peach (PpMLO1, PpMLO3 and PpMLO4), apricot (PaMLO1, PaMLO3 and PaMLO4) and woodland strawberry (FvMLO1, FvMLO4 and FvMLO12) were found to cluster together in the phylogenetic clade V, containing all the dicot MLO proteins experimentally shown to be required for PM susceptibility (e.g. [16,23]). One homolog from strawberry (FvMLO17) and one from peach (PpMLO12) were found to group, together with grapevine VvMLO14, in clade IV, which contains all

monocot MLO proteins acting as PM susceptibility factors (Figure 1).

We used the GBrowse-Syn tool to detect syntenic blocks encompassing *P. persica*, *F. vesca* and *M. domestica* MLO genes. As syntenic blocks derive from the evolution of the same chromosomal region after speciation, orthology between MLO genes could be inferred. In total, twelve orthologous relationships were predicted between *P. persica* and *F. vesca*, nine between *P. persica* and *M. domestica* and eight between *F. vesca* and *M. domestica* (Table 4, Figure 2 and Additional file 1).

Transcription of putative apple MLO genes in response to *Podosphaera leucotricha* inoculation

To identify MLO genes that respond to the PM fungus *P. leucotricha*, we measured the transcript abundance of 19 out of 21 apple MLO genes in leaves 4, 6, 8 and 24 hours after artificial inoculation with the pathogen, and compared these data with the ones of non-inoculated leaves. Three cultivars, Golden Delicious, Braeburn and Gala, were analysed in order to investigate whether up-regulation was comparable among them and results could therefore be generalized for all apple cultivars. Three genes, namely

Table 2 Members of the *PpMLO* gene family as predicted in *Prunus persica* genome sequence

Gene	Accession number ^a	Chr.	Starting position (Mb)	Clade	Introns	TM ^b	Amino acids	Conserved aa ^c
<i>PpMLO1</i>	ppa003207m	6	6.82	V	14	7	593	30
<i>PpMLO2</i>	ppa003435m	7	18.38	III	14	8	574	30
<i>PpMLO3</i>	ppa003437m	6	21.99	V	13	7	574	30
<i>PpMLO4</i>	ppa003466m	2	21.03	V	14	7	572	30
<i>PpMLO5</i>	ppa003706m	4	10.92	I	14	8	555	30
<i>PpMLO6</i>	ppa004012m	7	22.64	II	14	6	535	29
<i>PpMLO7</i>	ppa004508m	8	21.17	II	0	7	506	29
<i>PpMLO8</i>	ppa004621m	6	22.01	VI	14	6	499	29
<i>PpMLO9</i>	ppa004687m	4	2.59	VII	11	7	496	29
<i>PpMLO10</i>	ppa004866m	2	13.73	II	11	7	488	29
<i>PpMLO11</i>	ppa020172m	1	43.04	I	14	4	561	30
<i>PpMLO12</i>	ppa020311m	5	0.82	IV	13	7	566	30
<i>PpMLO13</i>	ppa021048m	4	15.57	VIII	12	5	510	24
<i>PpMLO14</i>	ppa022847m	6	6.80	VI	14	6	550	29
<i>PpMLO15</i>	ppa024476m	7	17.63	I	14	8	539	26
<i>PpMLO16</i>	ppa024488m	5	0.76	III	14	6	504	30
<i>PpMLO17</i>	ppa024581m	6	8.95	II	13	6	463	27
<i>PpMLO18</i>	ppa026565m	6	22.00	VI	13	6	416	25
<i>PpMLO19</i>	ppb024523m	1	42.04	II	13	5	446	23

^aAvailable at http://www.rosaceae.org/gb/gbrowse/prunus_persica/.

^bNumber of transmembrane domains in the predicted protein, as determined by InterPro prediction software (<http://www.ebi.ac.uk/interpro/>).

^cnumber of conserved amino acids out of the 30 identified by Elliot et al. [29].


Table 3 Members of the *FvMLO* gene family as predicted in *Fragaria vesca* genome sequence

Gene	Accession number ^a	Chr.	Starting position (Mb)	Clade	Introns	TM ^b	Amino acids	Conserved aa ^c
<i>FvMLO1</i>	mrna02774.1-v1.0-hybrid	n.d.	n.d.	V	14	7	632	28
<i>FvMLO2</i>	mrna03210.1-v1.0-hybrid	3	14.46	II	11	5	528	20
<i>FvMLO3</i>	mrna09651.1-v1.0-hybrid	6	35.88	III	14	6	542	28
<i>FvMLO4</i>	mrna09653.1-v1.0-hybrid	6	35.90	V	14	7	573	30
<i>FvMLO5</i>	mrna10166.1-v1.0-hybrid	1	1.34	II	14	3	688	26
<i>FvMLO6</i>	mrna10346.1-v1.0-hybrid	3	12.52	II	7	2	385	15
<i>FvMLO7</i>	mrna10363.1-v1.0-hybrid	3	12.49	II	9	2	442	21
<i>FvMLO8</i>	mrna10558.1-v1.0-hybrid	2	19.08	II	n.d.	6	514	28
<i>FvMLO9</i>	mrna11028.1-v1.0-hybrid	n.d.	n.a.	I	10	4	434	18
<i>FvMLO10</i>	mrna13023.1-v1.0-hybrid	1	7.96	III	13	6	557	27
<i>FvMLO11</i>	mrna14592.1-v1.0-hybrid	1	8.77	I	13	7	548	28
<i>FvMLO12</i>	mrna23198.1-v1.0-hybrid	7	15.89	V	14	7	507	29
<i>FvMLO13</i>	mrna26428.1-v1.0-hybrid	7	17,79	VIII	11	5	558	20
<i>FvMLO14</i>	mrna28541.1-v1.0-hybrid	n.d.	n.a.	III	11	4	481	26
<i>FvMLO15</i>	mrna29770.1-v1.0-hybrid	3	7.36	VII	13	7	538	28
<i>FvMLO16</i>	mrna31264.1-v1.0-hybrid	3	30.51	I	16	8	579	28
<i>FvMLO17</i>	mrna31498.1-v1.0-hybrid	5	20.23	IV	11	5	531	27
<i>FvMLO18</i>	mrna29285.1-v1.0-hybrid	5	19.12	V	6	4	357	18

^aAvailable at http://www.rosaceae.org/gb/gbrowse/fragaria-vesca_v1.0-1g/ (hybrid).

^bNumber of transmembrane domains in the predicted protein, as determined by InterPro prediction software (<http://www.ebi.ac.uk/interpro/>).

^cnumber of conserved amino acids out of the 30 identified by Elliot et al. [29].



MdMLO11, *MdMLO18* and *MdMLO19*, were found to be significantly up-regulated after inoculation with the pathogen (Figure 3 and Additional file 2). Up-regulation of these genes was about 2-fold compared to non-inoculated plants, with peaks of 4-fold up-regulation at very early time points ('Braeburn'- *MdMLO11* - 6 hpi; 'Gala'- *MdMLO18* - 4 hpi; 'Golden Delicious'- *MdMLO19* - 6hpi). *MdMLO11* and *MdMLO18* were up-regulated in all cultivars, while *MdMLO19* was only up-regulated in 'Braeburn' and 'Golden Delicious'.

Two of the genes, *MdMLO11* and *MdMLO19* belong to Clade V, while *MdMLO18* belongs to the newly identified Clade VII (Figure 1).

Discussion

Genomic organization and phylogenetic relations between Rosaceae MLO homologs

We report here the identification, through an *in silico* approach, of 19 MLO homologs in the genome of peach and 18 in the genome of strawberry. This is consistent with the results of previous genome-wide studies carried out on dicotyledonous species, indicating the presence of 15 MLO homologs in *Arabidopsis*, 17 in grapevine and 16 in tomato [9,13]; Appiano et al., unpublished results; [24]. Conversely, the number of MLO homologs detected in apple (21) is lower than expected, considering that a relatively recent genome-wide duplication event has occurred in the Pyreae tribe [32].

Table 4 Relations of orthology inferred between *P. persica*, *F. vesca* and *M. domestica* MLO homologs

<i>P. persica</i> genes	<i>F. vesca</i> orthologs	<i>M. domestica</i> orthologs
<i>PpMLO2</i>	<i>FvMLO10</i>	<i>MdMLO15</i>
<i>PpMLO3</i>	<i>FvMLO4</i>	<i>MdMLO19</i>
<i>PpMLO4</i>	<i>FvMLO12</i>	-
<i>PpMLO5</i>	<i>FvMLO16</i>	<i>MdMLO10</i> , <i>MdMLO17</i>
<i>PpMLO6</i>	<i>FvMLO5</i>	<i>MdMLO1</i>
<i>PpMLO7</i>	<i>FvMLO8</i>	-
<i>PpMLO8</i>	<i>FvMLO3</i>	-
<i>PpMLO9</i>	<i>FvMLO15</i>	<i>MdMLO18</i>
<i>PpMLO10</i>	<i>FvMLO2</i>	<i>MdMLO9</i>
<i>PpMLO14</i>	<i>FvMLO14</i>	<i>MdMLO21</i>
<i>PpMLO15</i>	<i>FvMLO11</i>	-
<i>PpMLO16</i>	-	<i>MdMLO12</i>
<i>PpMLO18</i>	<i>FvMLO3</i>	-

Relations of orthology between *PpMLO1*, *PpMLO3*, *PpMLO4* and apricot *PaMLO1*, *PaMLO3*, *PaMLO4* were clearly suggested by the high percentage of sequence identity between these homolog genes, which was 97,3%, 98,8% and 96,7%, respectively.

Most *PpMLO*, *FvMLO* and *MdMLO* homologs appeared to be widely distributed within the respective genomes (Tables 1, 2 and 3), indicating segmental duplication as the prevailing evolutionary mechanism for the Rosaceae *MLO* gene family. However, we also found cases of clusters of adjacent homologs (*PpMLO3*, *PpMLO8* and *PpMLO18*, *PpMLO12* and *PpMLO16*, *PpMLO1* and *PpMLO14*, *FvMLO3* and *FvMLO4*, *FvMLO6* and *FvMLO7*, *MdMLO2*, *MdMLO3* and *MdMLO8*), which are likely the result of tandem duplication events.

Inference of phylogenetic relationships between *MLO* proteins revealed the presence of apple, strawberry, peach and apricot homologs in the clade V, containing all dicot *MLO* homologs shown so far to be involved in PM susceptibility, thus making them candidates to act as susceptibility factors. Although the simple clustering in clade V is not enough to recognize a gene as a susceptibility factor, it does provide the first evidence for functionality and allows for the reduction in the number of candidates for further functional analysis. Clade IV, that contains functional *MLO* susceptibility homologs from monocots, was found to include one homolog from *F. vesca* (*FvMLO17*) and one from *P. persica* (*PpMLO12*). In accordance with this finding, a *MLO* homolog from the dicot species *V. vinifera* also clusters in clade IV [23,24]. Figure 1). Interestingly, phylogenetic analyses carried out in this study also revealed the presence of one or two additional clades, depending on the type of phylogenetic reconstruction (UPGMA or Neighbour-Joining), which were not reported to occur in earlier investigations. Moreover, they appear to be characteristic of Rosaceae, since they contain only


homologs from this family. Clearly, the exclusivity for Rosaceae of these clade(s) needs to be confirmed by further studies containing a larger dataset of *MLO* proteins. Additional studies could be also addressed to the functional characterization of Rosaceae *MLO* homologs grouped in clade VII. Indeed, this appears to be basal to both clade IV and clade V (Figure 1), and thus might have contained ancestral proteins which later on evolved into PM susceptibility factors.

Synteny between apple, peach and woodland strawberry *MLO* genes

We took advantage of recent developments in Rosaceae genomics in order to detect synteny between *P. persica*, *F. vesca* and *M. domestica* chromosomal regions containing *MLO* homologs. This permitted the inference of orthologous relationships between *MLO* genes in these species. Notably, all predicted *MLO* orthologs from different Rosaceae species, fell in the same phylogenetic clade (Tables 1, 2 and 3; Figure 1 and Additional file 1). This is to be expected, since orthologs generally share the same function and thus are characterized by a high level of sequence conservation. It is noteworthy that the chromosomal localization of predicted *MLO* orthologs between *P. persica*, *M. domestica* and *F. vesca* is in accordance with the results of the synteny study performed after the release of the three genomes [33,34]. In particular, genes situated on peach scaffold 2, 7 and 8 were predicted to have orthologs on strawberry chromosome 7, 1 and 2, respectively, whereas genes on peach scaffold 4 were predicted to have orthologs on strawberry chromosomes 2 or 3 (Table 4). *FvMLO3* was predicted to be orthologous to two peach *MLO* genes, *PpMLO8* and *PpMLO18*, which are localised in close proximity to each other on peach scaffold 6 and grouped together in clade VI. In this case, we hypothesize a relation of co-orthology due to the occurrence of a recent tandem duplication event in the peach genome. Similarly, *PpMLO5* and *FvMLO16* were predicted to be orthologs of two apple *MLO* genes, *MdMLO10* and *MdMLO17*, located on chromosomes 3 and 11. This is consistent with indications of duplications of large segments of these two chromosomes during the evolution of the apple genome [32].


Transcription of apple putative *MLO* genes in response to *P. leucotricha* inoculation

In barley, pea and tomato, only one of the clade V *MLO* homologs seems to be involved in powdery mildew susceptibility, whereas in *A. thaliana* three *MLO* genes in Clade V are required to be inactivated in order to achieve a fully resistant phenotype [16,27]. This implies that, within Clade V *MLO* genes, a further selection might be required to identify PM susceptibility genes. Accumulating



evidence indicates that *MLO* susceptibility genes are up-regulated upon challenge with powdery mildew fungi [17]. Therefore, we analysed the expression level of apple *MLO* genes identified in this study in response to the interaction with *P. leucotricha*. Three pathogen-dependent gene up-regulations were detected. Two up-regulated *MLO* homologs, *MdMLO11* and *MdMLO19*, encode for proteins falling in clade V, thus making them likely candidates to act as PM susceptibility genes in apple. *MdMLO11* and *MdMLO19* are located on chromosomes 4 and 12 respectively, and are therefore both generated from a duplication event in the 9-

chromosome ancestor of apple [32]. A third pathogen-dependent up-regulated gene, *MdMLO18*, was found, which encodes a protein grouping in the newly identified Clade VII (Figure 1). The presence of a PM upregulated gene outside clade V is consistent with transcriptome analyses recently performed in tomato (Appiano *et al.*, unpublished results). Apple clade V also contains two genes, *MdMLO5* and *MdMLO7*, which show no significant changes in expression following inoculation. Accordingly, the lack of up-regulation of some clade V *MLO* genes has been observed in grapevine and tomato [23,24]; Appiano *et al.*, unpublished results). The



possible role of these genes as susceptibility factors has not yet been highlighted.

PpMLO3, *PaMLO3* and *FvMLO4* are likely to represent true orthologs of *MdMLO19* (Table 4). Since orthologs often maintain the same function during evolution, we conjecture that the expression of these genes might also be responsive to PM fungi attacking corresponding species. Moreover, *FvMLO15* and *PpMLO9* are likely orthologs of *MdMLO18*, so they should also be considered as putative transcriptionally responsive genes to PM fungi attack. Further studies aimed at the functional characterization of these genes (e.g. through the application of reverse genetic approaches of targeted mutagenesis or gene silencing), in apple but also in peach and strawberry, might lead to the identification of resistant phenotypes, which could be used for the development of PM resistant cultivars. Particularly, studies on *MdMLO18*

could lead to the characterization of a possible role for clade VII in the interaction with PM fungi.

Conclusions

Our work led to the identification of 19 MLO homologs in peach, 17 in strawberry and 21 in apple. Three, three and four homologs, respectively, belong to clade V and therefore are candidates for being S-genes. Due to the high similarity between peach and apricot, we were able to amplify and characterize three Clade V apricot MLO genes.

The phylogenetic analysis revealed two new Rosaceae specific clades for the MLO family, although this needs to be confirmed by the use of a larger MLO proteins dataset.

Through inoculation of apple with *P. leucotricha*, we identified three up-regulated genes, i.e. *MdMLO11*,

MdMLO18 and *MdMLO19*. *MdMLO11* and *MdMLO19*, that belong to Clade V, are positioned in duplicated regions and have high sequence identity, therefore they are likely to be recent paralogs. *MdMLO18* belongs to the newly identified Clade VII.

Methods

In silico identification and comparison of MLO predicted proteins in peach, woodland strawberry and apple

Predicted peptides from the peach genome (v. 1.0) and the strawberry genome (v.1.0) gene model databases, available at the website of the Genomic Database for Rosaceae (www.rosaceae.org) [35], were queried for the presence of MLO homologs protein sequences. First, a BLAST search, using the tomato SIMLO1 amino acid sequence as query was carried out. A further search was performed with the HMMER programme, which uses a method for homolog searches based on the profile hidden Markov probabilistic model [36]. The sequences obtained with the previously mentioned BLAST search, were used together with other known MLO sequences from dicot and monocot species, namely: four RhMLOs from *Rosa hybrida*, 15 AtMLOs from *Arabidopsis thaliana*, SIMLO1 from *Solanum lycopersicum*, CaMLO2 from *Capsicum annuum*, PsMLO1 from *Pisum sativum*, MtMLO1 from *Medicago truncatula*, LjMLO1 from *Lotus japonicus*, VvMLO14 from *V. Vinifera*, HvMLO from *Hordeum vulgare*, TaMLO1_A1b and TaMLO_B1 from *Triticum aestivum* and OsMLO2 from *Oryza sativa*. MLO protein sequences from apple (*Malus domestica* cv. Golden Delicious) were identified by searching for the MLO domain profile (IPR004326) in the apple genome available at FEM-IASMA computational biology web resources (<http://genomics.research.iasma.it>). The resulting list was integrated with a BLAST search, carried out with the amino acid sequences previously listed for the HMMER search in peach and strawberry.

Chromosomal localization and predicted introns/exons structure of each MLO gene of apple, peach and strawberry was deduced based on the available genomic information at the GDR database. The presence and number of membrane spanning helices was predicted using the online software InterPro (<http://www.ebi.ac.uk/interpro>). Alignments for conserved amino-acids analysis were performed with the CLC Sequence Viewer v. 6.9 software (<http://clcbio.com>).

A total of 90 MLO protein sequences, including three apricot MLO sequences isolated *in vitro* (see next paragraph), were used for Clustal alignment (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>). UPGMA-based and Neighbour-Joining-based phylogenetic trees were obtained with the CLC sequence viewer software. The UPGMA clustering algorithm was further used as input for the Dendroscope software, suitable for the visualization of large phylogenetic trees [37].

Relationships of orthology between MLO candidate genes from peach, woodland strawberry and apple were inferred by running the GBrowse-Syn tool available at GDR (http://www.rosaceae.org/gb/gbrowse_syn/peach_apple_strawberry) [35,38]. This displays syntenic regions among the three available genomes of Rosaceae, as detected by the Mercator programme [35,39]. For 50 Kb chromosomal stretches flanking each *P. persica* *PpMLO* homolog, syntenic regions from *F. vesca* and *M. domestica* were searched. Orthology was called upon the identification of *F. vesca* or *M. domestica* MLO homologs within syntenic blocks.

In vitro isolation of apricot MLO homologs

RNA from apricot leaves (cultivar Orange Red) was extracted by using the SV Total RNA Isolation System Kit (Promega), and corresponding cDNA was synthesized by using the QuantiTect Reverse Transcription Kit (Qiagen) with oligo(dT) primers. Sequences of the peach MLO homologs *PpMLO1*, *PpMLO3* and *PpMLO4*, are phylogenetically close to MLO homologs functionally associated to PM susceptibility, and were therefore used to design the primer pairs 5'-ATGGCAGCCGCAACCTCAGG AAGA-3'/5'-TTATATACTTTGCCTATTGTCAAAC-3', 5'-ATGGCAGGGGGAAAAGAAGGACG-3'/5'-TCAAC TCCTTTCTGATTTCTCAA-3' and 5'-ATGGCCGA ACTAAGTAAAGA-3'/5'-TCAACTTCTTGATTTTCC TTTGC-3', respectively. These were employed to amplify full-length cDNA sequences of apricot putative orthologs, by using the AccuPrime Taq polymerase (Invitrogen). Amplicons were purified by using the NucleoSpin Extract II kit (Macherey-Nagel) and ligated (molar ratio 1:1) into the pGEM-T Easy vector (Promega). Recombinant plasmids were cloned in *E. coli* DH10 β chemically competent cells and recovered by using the Qiaprep spin miniprep kit (Qiagen). Sequencing reactions were performed twice, by using universal T7 and SP6 primers (Eurofins MWG Operon).

Glasshouse test with apple cultivars

A total of 192 apple plants from three cultivars (Braeburn, Golden Delicious and Gala) were used to measure transcript abundance of MLO genes. Budwoods from these cultivars were grafted on M9 rootstocks in January 2012. The grafts were kept at -1°C for 2 months, and potted at the beginning of March in greenhouse. The plants grew for 6 weeks in the greenhouse at 20°C during the day, 17°C during the night, relative humidity of 70% and natural day/night cycle.

P. leucothrica was collected from apple trees in an unsprayed test orchard and used to infect greenhouse grown apple seedlings from 'Gala Galaxy' seeds. Four weeks after inoculation, conidia were used for the inoculation experiment, or transferred to new seedlings, to keep them viable.

We inoculated by touching the plants with heavily infected apple seedlings. Control plants were not inoculated and kept separated in the same greenhouse of the inoculated plants. Inoculated and control plants were grown in the greenhouse at the growing conditions previously mentioned. The leaf samples were collected 4, 6, 8 and 24 hours post-inoculation (hpi).

Eight experimental repeats were performed and each sample contained three or four young leaves collected from each single plant. Every plant was used for sampling only once, to avoid any possible effect of wounding on the expression of *MLO* genes. The smallest statistical unit was a plant. The leaves were flash-frozen and ground in liquid nitrogen, and stored at -80°C until RNA extraction.

qPCR analysis of transcript levels

RNA extraction was carried out with the MagMAX-96 Total RNA isolation kit (Applied Biosystem) that includes DNase treatment. The kit yielded between 50 and 200 ng/ul, of good quality RNA per sample.

Primers for gene expression analysis were designed with NCBI Primer Designing Tool (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>). Four serial dilutions of cDNA (1/5 - 1/25 - 1/125 - 1/625) were used to calculate the

efficiency of each primer pair with iCycler software (Biorad). In case of efficiency lower than 1.80 or greater than 2.20, the primer pair was discarded and a new one tested, with the exception of *MdMLO9*, for which was not possible to design a primer pair with better efficiency. It was only possible to analyse 19 *MLO* genes because for *MdMLO12* and *MdMLO16* was not possible to design specific and efficient primer pairs, despite numerous attempts. Presence of a specific final dissociation curve was determined after each qPCR amplification reaction with progressive increment of temperature from 65°C to 95°C (0.5°C each step, 5 sec) and the size of the product was confirmed by agarose gel electrophoresis.

Quantitative Real Time-PCR (qPCR) was performed with SYBR greenER mix (Invitrogen) in a 15- μL reaction volume, using a Bio-Rad iCycler iQ detection system, run by the Bio-Rad iCycler iQ multicolor 3.1 software. The software applies comparative quantification with an adaptive baseline. Samples were run in two technical replicates with the following thermal cycling parameters: 95°C 3 min - 95°C 15 sec, 60°C 1 min (repeated 40 times) - 95°C 10 sec.

Reference genes β -actin (NCBI accession number DT002474; Plaza accession number MD00G171330 - <http://bioinformatics.psb.ugent.be/plaza/>), ubiquitin (Plaza

Table 5 Gene-specific primers and amplicon sizes in qRT-PCR detection of 19 *MdMLO*-like genes based on *Malus domestica* cv. Golden Delicious genome sequence

Gene	Forward primer (5' - 3')	Reverse primer (5' - 3')	Size (bp)	Efficiency
<i>MdMLO1</i>	GTGGGCTCGGTCGGCCAAAA	CCAGACCAGCACCAGAACCA	81	2.06
<i>MdMLO2</i>	CGTTGGATCAACCACTGCGCCT	TGAGCTGCAGCCAGTGGGATCT	87	1.83
<i>MdMLO3</i>	CCACTGCGCCTCTCTGAAGCA	CCACAAAACGGCTCTCCAGGT	93	2.12
<i>MdMLO4</i>	TGTTGCAGACACTATGCTGCCATGT	GCCAGCAGCTAAAGATCTGCGT	109	1.87
<i>MdMLO5</i>	TCGTCAGGCTCTCATTGGGGT	GTGCTGCTGCCACTCCCTC	132	1.80
<i>MdMLO6</i>	TTCGCGGAGGAGGGTCTGTT	TTCGAGCGACAGCAACGGCA	72	2.15
<i>MdMLO7</i>	TGGAGCAAGTCACCAGTCTCCAT	CGTTCTGGTGCCAAATGTGC	127	2.12
<i>MdMLO8</i>	GTC AAGCTAATCTTACCACGCGCT	GGCTGGAAGGAAGGACAGCCA	85	1.95
<i>MdMLO9</i>	GCTGCAACACGTAATCACCC	AGAACGCCATTTGAAAGCA	173	2.30
<i>MdMLO10</i>	GCGATCGTTGGCCTTGACTCC	TTCCGCGCTCGACAAGCAGA	86	1.92
<i>MdMLO11</i>	CCGTTCCATACCAAGACGA	ATTGCTCTCCGAGTTACGCC	102	1.90
<i>MdMLO13</i>	ACATTGTCCCCAGGCTTGTT	GCCCAACCAATAAGTCCCGA	151	2.00
<i>MdMLO14</i>	TGCACTTGTCAGCCAGATGGG	GCATCTCCCACCCACGAACCG	81	2.15
<i>MdMLO15</i>	GCGCCTTCTCTCTGCTGGGT	CGCGTGCAGGTGGTCTCTT	90	2.01
<i>MdMLO17</i>	TTGCCACTGTATGCCTTGGT	TGCTTGCTTCTGTGCGAATG	163	2.15
<i>MdMLO18</i>	AAGGAAGGCTCTCATTAGGCTCT	TGCAATTGGCTTTTGACCAACGGT	100	2.22
<i>MdMLO19</i>	CAGAGTGGCGACTGCACTTA	GGGACATGGAGTGCAAAGGA	110	1.97
<i>MdMLO20</i>	AAAAAGCTCCACCAACCCCA	TTTCTCTCCCATGACGCTCG	165	2.11
<i>MdMLO21</i>	CCTTGTTTCGAGGCCGTAGAG	ACCAAGTGCTTTGGTGGTTT	176	1.95
β -actin	CTATGTTCCCTGGTATTGCAGACC	GCCACAACCTTGATTTTCATGC	82	1.90
Ubiquitin	CATCCCCCAGACCAGCAGA	ACCACGGAGACGAAGCACCAA	349	1.91
Elongation Factor 1	TACTGGAACATCACAGGCTGAC	TGGACCTCTCAATCATGTTGTC	308	2.07

accession number MD05G001920) and elongation factor 1 (Plaza accession number MD09G014760) were used as reference genes (Table 5). All these three genes were used in previous works [40-42]. For additional control, we assessed the stability of our genes with the software geNorm (medgen.ugent.be/~jvdesomp/genorm/). An M-value lower than 1.5 is generally considered as stable enough [43-45] and all three reference genes in all three cultivars considered are within this threshold. We saw differences in stability between cultivars: 'Golden Delicious' was the most stable cultivar (actin: 0.824 – ubiquitin: 0,852 – elongation factor 1: 0,926), whereas 'Braeburn' was the less stable (actin: 1.246 – ubiquitin: 1,293 – elongation factor 1: 1,369) and 'Gala' showed intermediate stability (actin: 1.039 – ubiquitin: 1,152 – elongation factor 1: 1,078).

Each of the biological replicates was analysed in duplicate and the average of these two replicates was used for further analysis. In case of excessive difference between the two replicates (one Ct or more), the run was repeated. Considering the high number of samples and genes of interest, we opted for this approach in order to reduce the number of total runs. Data analysis was performed according to Hellemans *et al.* [46], using the statistical package SPSS (IBM). This analysis method takes into account the efficiency value of each primer pair. For some genes it was necessary to apply a natural log transformation to the data, in order to obtain normal distribution of residues. To investigate the differences between control and inoculated samples, we used *T*-test ($p \leq 0.05$).

Availability of supporting data

The following files are available on: mynotebook.labarchives.com.

Figure 1 – Phylogenetic tree of Rosaceae MLO.

DOI: 10.6070/H4Z60M0N.

Additional file 1 - Synteny between apple, peach and strawberry.

DOI: 10.6070/H4TD9V8C.

Additional files

Additional file 1: Synteny between apple, peach and strawberry.

Results of search for *F. vesca* and *M. domestica* regions syntenic to 50 kb *P. persica* chromosomal stretches containing the *PpMLO* homologs identified in this study. Shaded polygons indicate aligned regions between genomes. Grid lines are drawn to indicate insertions/deletions between the genomes of *F. vesca* and *M. domestica* with respect to the *P. persica* reference sequence. *P. persica*, *F. vesca* and *M. domestica* MLO homologs, named according to the nomenclature of the Genomic Database of Rosaceae, are boxed.

Additional file 2: Transcriptional variation of 19 apple MLO genes in three cultivars following inoculation with *P. leucotricha*.

Transcription abundances of 19 MLO-like genes following powdery mildew (PM) inoculation in 'Golden Delicious' (1a), 'Gala' (1b) and 'Braeburn' (1c) leaf samples. The graphs show expression values of inoculated samples relative to control samples, averaged from four to eight biological replicate, normalized, that are in turn the average of two

experimental replicates. The Ct values have been normalized with three reference genes: actin, ubiquitin and elongation factor 1. Statistical significance was determined with a *t*-test for each individual pair of inoculated and control samples at each time point (4, 6, 8 and 24 hpi). The error bars show standard errors of the means. Significant differences between inoculated samples and control samples are indicated with a * ($P < 0.05$).

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

SPe carried out the inoculation and gene expression analysis, identified MLO homologs in apple and wrote the major part of the manuscript. SPa contributed to design the experiment, identified MLO homologs in peach, apricot and strawberry, performed the analysis of synteny and contributed to write the manuscript. DC contributed to design the experiment, performed the phylogenetic analysis and revised the manuscript. AG contributed to design the experiment and revised the manuscript. YB contributed to the design of the experiments. RGFV contributed to the design of the experiments and revised the manuscript. MM contributed to the design of the experiments, contributed to the identification of MLO homologs in apple and revised the manuscript. HS contributed to the design of the experiments and revised the manuscript. All authors read and approved the final manuscript.

Acknowledgment

The authors would like to thank Rammelt Groenwold (WUR) for helping with apple inoculation with *P. leucotricha* and Lorenza Dalla Costa (FEM) for all the valuable advices about qPCR data analysis.

This work has been funded by Fondazione Edmund Mach and GMPF PhD programme.

Author details

¹Department of Genomics and Biology of Fruit Crops, Fondazione Edmund Mach, via E. Mach 1, 38010 San Michele all'Adige, Italy. ²Wageningen UR Plant Breeding, Wageningen University and Research Centre, P.O. Box 16, 6700 AA Wageningen, The Netherlands. ³Department of Soil, Plant and Food Science, University of Bari, Via Amendola 165/A, 70126 Bari, Italy. ⁴National Research Council, Institute of Plant Genetics, Via Amendola 165/A, 70126 Bari, Italy.

Received: 15 January 2014 Accepted: 10 July 2014

Published: 22 July 2014

References

1. Glawe DA: The powdery mildews: a review of the world most familiar (yet poorly known) plant pathogens. *Annu Rev Phytopathol* 2008, **46**:27–51.
2. Turechek WW, Carroll JE, Rosenberger DA: *Powdery Mildew of Apple*. New York State Integrated Pest Management Program: Cornell University; 2004 (www.nysipm.cornell.edu/factsheets/treefruit/diseases/pm/apple_pm.pdf).
3. Foulongne M, Pascal T, Pfeiffer F, Kervella J: QTLs for powdery mildew resistance in peach \times *Prunus davidiana* crosses: consistency across generations and environments. *Mol Breed* 2003, **12**:33–50.
4. Boesewinkel HJ: Differences between the conidial states of *Podosphaera tridactyla* and *Sphaerotheca pannosa*. *Annales de Phytopathologie* 1979, **11**:525–527.
5. Xiao CL, Chandler CK, Price JF, Duval JR, Mertely JC, Legard DE: Comparison of epidemics of botrytis fruit rot and powdery mildew of strawberry in large plastic tunnel and field production systems. *Plant Dis* 2001, **85**:901–909.
6. Fu XL, Lu Y, Liu XD, Li JQ: Crossability barriers in the interspecific hybridization between *Oryza sativa* and *O. Meyeriana*. *J Integr Plant Biol* 2009, **51**:21–28.
7. Parlevliet JE: What is durable resistance, a general outline. In *Durability of Disease Resistance*. Edited by Jacobs TH, Parlevliet JE. Dordrecht: Kluwer; 1993:23–29.
8. Parisi L, Lespinasse Y, Guillaumes SJ, Kruger J: A new race of *Venturia inaequalis* virulent to apples with resistance due to the *Vf* gene. *Phytopathology* 1993, **85**:533–537.

9. Krieghoff O: *Entwicklung einer In-vitro-selektionsmethode auf Resistenz von Malus-Genotypen gegenüber Podosphaera leucotricha (Ell. Et Ev.) Salm. und In-vitron-Differenzierung von Virulenzunterschieden des Erregers*. Berlin: PhD dissertation, Humboldt-Universität; 1995.
10. Pavan S, Jacobsen E, Visser RGF, Bai Y: **Loss of susceptibility as a novel breeding strategy for durable and broad-spectrum resistance**. *Mol Breed* 2010, **25**:1–12.
11. Büschges R, Hollricher K, Panstruga R, Simons G, Wolter M, Frijters A, van Daelen R, van der Lee T, Diergaarde P, Groenendijk J, Töpsch S, Vos P, Salamini F, Schulze-Lefert P: **The barley Mlo gene: a novel control element of plant pathogen resistance**. *Cell* 1997, **88**(5):695–705.
12. Devoto A, Piffanelli P, Nilsson I, Wallin E, Panstruga R, Von Heijne G, Schulze-Lefert P: **Topology, subcellular localization, and sequence diversity of the Mlo family in plants**. *J Biol Chem* 1999, **274**:34993–35004.
13. Jørgensen JH: **Discovery, characterization and exploitation of Mlo powdery mildew resistance in barley**. *Euphytica* 1992, **63**:141–152.
14. Consonni C, Humphry ME, Hartmann HA, Livaja M, Durner J, Westphal L, Vogel J, Lipka V, Kemmerling B, Schulze-Lefert P, Somerville SC, Panstruga R: **Conserved requirement for a plant host cell protein in powdery mildew pathogenesis**. *Nat Genet* 2006, **38**(6):716–720.
15. Humphry M, Reinstädler A, Ivanov S, Bisseling T, Panstruga R: **Durable broad-spectrum powdery mildew resistance in pea er1 plants is conferred by natural loss-of-function mutations in PsMLO1**. *Mol Plant Pathol* 2011, **12**(9):866–878.
16. Pavan S, Schiavulli A, Appiano M, Marcotrigiano AR, Cillo F, Visser RGF, Bai Y, Lotti C, Ricciardi L: **Pea powdery mildew er1 resistance is associated to loss-of-function mutations at a MLO homologous locus**. *Theor Appl Genet* 2011, **123**:1425–1431.
17. Bai Y, Pavan S, Zheng Z, Zappel N, Reinstädler A, Lotti C, De Giovanni C, Ricciardi L, Lindhout P, Visser RGF, Theres K, Panstruga R: **Naturally occurring broad-spectrum powdery mildew resistance in a Central American tomato accession is caused by loss of MLO function**. *Mol Plant Microbe Interact* 2008, **21**:30–39.
18. Aist JR, Bushnell WR: **Invasion of plants by powdery mildew fungi, and cellular mechanisms of resistance**. In *The Fungal Spore and Disease Initiation in Plants and Animals*. Edited by Cole GT, Hoch HC. New York: plenum press; 1991:321–345.
19. Miklis M, Consonni C, Bhat RA, Lipka V, Schulze-Lefert P, Panstruga R: **Barley MLO modulates actin-dependent and actin-independent antifungal defense pathways at the cell periphery**. *Plant Physiol* 2007, **144**:1132–1143.
20. Panstruga R: **Serpentine plant MLO proteins as entry portals for powdery mildew fungi**. *Biochem Soc Transact* 2005, **33**(Pt 2):389–392.
21. Piffanelli P, Zhou FS, Casais C, Orme J, Jarosch B, Schaffrath U, Collins NC, Panstruga R, Schulze-Lefert P: **The barley MLO modulator of defense and cell death is responsive to biotic and abiotic stress stimuli**. *Plant Physiol* 2002, **129**:1076–1085.
22. Zheng Z, Nonomura T, Bóka K, Matsuda Y, Visser RGF, Toyoda H, Kiss L, Bai Y: **Detection and quantification of *Leveillula taurica* growth in pepper leaves**. *Phytopathology* 2013, **103**:6.
23. Feechan A, Jermakow AM, Torregrosa L, Panstruga R, Dry IB: **Identification of grapevine MLO gene candidates involved in susceptibility to powdery mildew**. *Funct Plant Biol* 2008, **35**:1255–1266.
24. Winterhagen P, Howard SF, Qiu W, Kovács LG: **Transcriptional up-regulation of grapevine MLO genes in response to powdery mildew infection**. *Am J Enol Vit* 2008, **59**:2.
25. Kessler SA, Shimosato-Asano H, Keinath NF, Wuest SE, Ingram G, Panstruga R, Grossniklaus U: **Conserved molecular components for pollen tube reception and fungal invasion**. *Science* 2010, **330**:968.
26. Chen Z, Noir S, Kwaaitaal M, Hartmann A, Wu MJ, Mudgil Y, Sukumar P, Munday G, Panstruga R, Jones AM: **Two seven-transmembrane domain MILDEW RESISTANCE LOCUS O proteins function in *Arabidopsis* root thigmomorphogenesis**. *Plant Cell* 2009, **21**:1972–1991.
27. Reinstädler A, Müller J, Czembor JH, Piffanelli P, Panstruga R: **Novel induced mlo mutant alleles in combination with site-directed mutagenesis reveal functionally important domains in the heptahelical barley Mlo protein**. *BMC Plant Biol* 2010, **10**:31.
28. Devoto A, Hartmann HA, Piffanelli P, Elliott C, Simmons C, Taramino G, Goh CS, Cohen FE, Emerson BC, Schulze-Lefert P, Panstruga R: **Molecular phylogeny and evolution of the plant-specific seven-transmembrane MLO family**. *J Mol Evol* 2003, **56**(1):77–88.
29. Elliott C, Muller J, Miklis M, Bhat RA, Schulze-Lefert P, Panstruga R: **Conserved extracellular cysteine residues and cytoplasmic loop-loop interplay are required for functionality of the heptahelical MLO protein**. *Biochem J* 2005, **385**(Pt 1):243–254.
30. Kaufmann H, Qiu X, Wehmeyer J, Debener T: **Isolation, molecular characterization, and mapping of four rose MLO orthologs**. *Front Plant Sci* 2012, **3**:244.
31. Elliott C, Zhou F, Spielmeier W, Panstruga R, Schulze-Lefert P: **Functional conservation of wheat and rice Mlo orthologs in defense modulation to the powdery mildew fungus**. *Mol Plant Microbe Interact* 2002, **15**(10):1069–1077.
32. Velasco R, Zharkikh A, Affourtit J, Dhingra A, Cestaro A, Kalyanaraman A, Fontana P, Bhatnagar SK, Troggio M, Pruss D, Salvi S, Pindo M, Baldi P, Castelletti S, Cavaiuolo M, Coppola G, Costa F, Cova V, Dal Ri A, Goremykin V, Komjanc M, Longhi S, Magnago P, Malacarne G, Malnoy M, Micheletti D, Moretto M, Perazzolli M, Si-Ammour A, Vezzulli S, et al: **The genome of the domesticated apple (*Malus × domestica* Borkh)**. *Nat Genet* 2010, **42**:833–839.
33. Shulaev V, Sargent DJ, Crowhurst RN, Mockler TC, Folkerts O, Delcher AL, Jaiswal P, Mockaitis K, Liston A, Mane SP, Burns P, Davis TM, Slovin JP, Bassil N, Hellens RP, Evans C, Harkins T, Kodira C, Desany B, Crasta OR, Jensen RV, Allan AC, Michael TP, Setubal JC, Celton J, Rees DJG, Williams KP, Holt SH, Ruiz-Rojas JJ, Chatterjee M, et al: **The genome of woodland strawberry (*Fragaria vesca*)**. *Nat Genet* 2011, **43**:109–116.
34. Jung S, Cestaro A, Troggio M, Main D, Zheng P, Cho I, Folta K, Sosinski B, Abbot B, Celton JM, Arus P, Shulaev V, Verde I, Morgante M, Rokhsar D, Velasco R, Sargent DJ: **Whole genome comparisons of *Fragaria*, *Prunus* and *Malus* reveal different modes of evolution between Rosaceae subfamilies**. *BMC Genomics* 2012, **13**:129.
35. Jung S, Ficklin SP, Lee T, Cheng C, Blenda A, Zheng P, Yu J, Bombarely A, Cho I, Ru S, Evans KM, Peace C, Abbott A, Mueller L, Olmstead M, Main D: **The genome database for rosaceae (GDR): year 10 update**. *Nucleic Acid Res* 2014, **42**:D1237–D1244.
36. Finn RD, Clements J, Eddy SR: **HMMER web server: interactive sequence similarity searching**. *Nucleic Acids Res* 2011, **39**:W29–W37.
37. Huson D, Richter D, Rausch C, DeZulian T, Franz M, Rupp R: **Dendroscope: an interactive viewer for large phylogenetic trees**. *BMC Bioinformatics* 2007, **8**:460.
38. McKay SJ, Vergara IA, Stajich JE: **Using the generic synteny browser (GBrowse_syn)**. *Curr Protoc Bioinformatics* 2010, **Chapter 9**(Unit 9):12.
39. Dewey CN: **Aligning multiple whole genomes with Mercator and MAVID**. *Methods Mol Biol* 2007, **395**:221–236.
40. Kürkcüoğlu S, Degenhardt J, Lensing J, Al-Masri AN, Gaul AE: **Identification of differentially expressed genes in *Malus domestica* after application of the non-pathogenic bacterium *Pseudomonas fluorescens*Bk3 to the phyllosphere**. *J Exp Bot* 2007, **58**(3):733–741.
41. Giorno F, Guerriero G, Baric S, Mariani C: **Heat shock transcriptional factors in *Malus domestica*: identification, classification and expression analysis**. *BMC Genomics* 2012, **13**:639.
42. Dal Cin V, Danesin M, Boschetti A, Dorigoni A, Ramina A: **Ethylene biosynthesis and perception in apple fruitlet abscission (*Malus domestica* L. Borkh)**. *J Exp Bot* 2005, **56**(421):2995–3005.
43. Ling D, Salvaterra PM: **Robust RT-qPCR data normalization: validation and selection of internal reference genes during post-experimental data analysis**. *PLoS One* 2011, **6**:3.
44. Van Hiel MB, Van Wielendaele P, Temmerman L, Van Soest S, Vuerinckx K, Huybrechts R, Broeck JV, Simonet G: **Identification and validation of housekeeping genes in brains of the desert locust *Schistocerca gregaria* under different developmental conditions**. *BMC Mol Biol* 2009, **10**:56.
45. Strube C, Buschbaum S, Wolken S, Schnieder T: **Evaluation of reference genes for quantitative real-time PCR to investigate protein disulfide isomerase transcription pattern in the bovine lungworm *Dictyocaulus viviparus***. *Gene* 2008, **425**:36–43.
46. Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J: **qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data**. *Genome Biol* 2007, **8**:R19.

doi:10.1186/1471-2164-15-618

Cite this article as: Pessina et al.: Characterization of the MLO gene family in Rosaceae and gene expression analysis in *Malus domestica*. *BMC Genomics* 2014 **15**:618.