



# Discovery of a Eukaryotic Pyrroloquinoline Quinone-Dependent Oxidoreductase Belonging to a New Auxiliary Activity Family in the Database of Carbohydrate-Active Enzymes

Hirotohi Matsumura<sup>1,3\*</sup>, Kiwamu Umezawa<sup>2,3</sup>, Kouta Takeda<sup>3,3</sup>, Naohisa Sugimoto<sup>1</sup>, Takuya Ishida<sup>1,4</sup>, Masahiro Samejima<sup>1</sup>, Hiroyuki Ohno<sup>3</sup>, Makoto Yoshida<sup>2\*</sup>, Kiyohiko Igarashi<sup>1,4\*</sup>, Nobuhumi Nakamura<sup>3,4\*</sup>

**1** Department of Biomaterials Sciences, Graduate School of Agriculture and Life Sciences, The University of Tokyo, Tokyo, Japan, **2** Department of Environmental and Natural Resource Science, Tokyo University of Agriculture and Technology, Tokyo, Japan, **3** Department of Biotechnology and Life Science, Tokyo University of Agriculture and Technology, Tokyo, Japan, **4** Advanced Low Carbon Technology Research and Development Program (ALCA), Japan Science and Technology Agency (JST), Tokyo, Japan

## Abstract

Pyrroloquinoline quinone (PQQ) is a redox cofactor utilized by a number of prokaryotic dehydrogenases. Not all prokaryotic organisms are capable of synthesizing PQQ, even though it plays important roles in the growth and development of many organisms, including humans. The existence of PQQ-dependent enzymes in eukaryotes has been suggested based on homology studies or the presence of PQQ-binding motifs, but there has been no evidence that such enzymes utilize PQQ as a redox cofactor. However, during our studies of hemoproteins, we fortuitously discovered a novel PQQ-dependent sugar oxidoreductase in a mushroom, the basidiomycete *Coprinopsis cinerea*. The enzyme protein has a signal peptide for extracellular secretion and a domain for adsorption on cellulose, in addition to the PQQ-dependent sugar dehydrogenase and cytochrome domains. Although this enzyme shows low amino acid sequence homology with known PQQ-dependent enzymes, it strongly binds PQQ and shows PQQ-dependent activity. BLAST search uncovered the existence of many genes encoding homologous proteins in bacteria, archaea, amoebozoa, and fungi, and phylogenetic analysis suggested that these quinoproteins may be members of a new family that is widely distributed not only in prokaryotes, but also in eukaryotes.

**Citation:** Matsumura H, Umezawa K, Takeda K, Sugimoto N, Ishida T, et al. (2014) Discovery of a Eukaryotic Pyrroloquinoline Quinone-Dependent Oxidoreductase Belonging to a New Auxiliary Activity Family in the Database of Carbohydrate-Active Enzymes. PLoS ONE 9(8): e104851. doi:10.1371/journal.pone.0104851

**Editor:** Claudio M. Soares, Instituto de Tecnológica Química e Biológica, UNL, Portugal

**Received:** April 13, 2014; **Accepted:** July 16, 2014; **Published:** August 14, 2014

**Copyright:** © 2014 Matsumura et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

**Funding:** This work was supported financially by grants from JSPS (No. 21605004 to N. N.), by a Grant-in-Aid for Innovative Areas (No. 24114001 and 24114008 to K. I.) from the Japanese Ministry of Education, Culture, Sports, and Technology (MEXT), and by a grant from the Advanced Low Carbon Technology Research and Development Program (ALCA) of the Japan Science and Technology Agency (JST) to K. I. and N. N. H. M. was supported by a research fellowship from the Japan Society for the Promotion of Science (JSPS) for Young Scientists (Grant No. 208304) during his postdoc period at the University of Tokyo. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* Email: ymakoto@cc.tuat.ac.jp (MY); aquarius@mail.ecc.u-tokyo.ac.jp (KI); nobu1@cc.tuat.ac.jp (NN)

† These authors contributed equally to this work.


‡ Current address: Institute of Environmental Health, Oregon Health and Science University, Portland, Oregon, United States of America

## Introduction

Pyrroloquinoline quinone (Figure 1, PQQ), a cofactor of prokaryotic dehydrogenases, was discovered as a redox cofactor of bacterial glucose dehydrogenase in 1964 [1]. Its chemical structure was reported after its isolation from primary alcohol dehydrogenase of methylotrophic bacteria [2]. Some prokaryotic organisms are able to synthesize PQQ, whereas others require an exogenous source. The enzyme group to which the PQQ cofactor belongs is the quinoprotein group [3–5], and all PQQ-dependent quinoproteins reported to date have a characteristic propeller-fold superbarrel structure [3,5]. Bacterial PQQ-dependent dehydrogenases which have an eight-bladed structure contain the characteristic amino acid sequence, which has been usually used to identify PQQ-dependent proteins. Although PQQ is physiologi-

cally important in plants [6] as well as mammals (including humans) [7–10], there has been no evidence showing PQQ as a cofactor of any eukaryotic enzyme so far.

A new class, named “Auxiliary Activities” (AA), was recently introduced in the database of carbohydrate-active enzymes (CAZy) to cover the full extent of the lignocellulose breakdown machinery [11]. This class covers a broad group of catalytic modules associated with plant cell wall degradation. The members of the AA family 8 consist of a cytochrome domain of spectral class *b* (protoheme IX); the first example of this family was found as a domain of cellobiose dehydrogenase (CDH; EC 1.1.99.18) isolated from the wood-rotting basidiomycete *Phanerochaete chrysosporium*, in which the cytochrome domain was linked to a flavin-containing AA family 3 flavin-containing sugar dehydrogenase domain [12]. We are interested in the AA family 8 hemoproteins



**Figure 1. Chemical structure of PQQ.**  
doi:10.1371/journal.pone.0104851.g001


because the heme domain is the first cytochrome *b*-type heme bound through Met/His with a primarily  $\beta$ -structure [13], and appears to be unrelated to any known protein other than the cytochrome domain of CDHs.

The basidiomycete *Coprinopsis cinerea* (formerly known as *Coprinus cinereus*, Figure 2) produces many extracellular oxidoreductases, including peroxidases and laccases, as well as cellulolytic enzymes, and has many putative oxidoreductase genes. To identify novel proteins with this unique heme *b*-containing cytochrome domain, the amino acid sequence of the cytochrome domain of CDH from *P. chrysosporium* was subjected to BLAST search using the *C. cinerea* Okayama 7 (#130) genome database at the Broad Institute. We fortuitously discovered a novel PQQ-dependent protein that belongs to a category that is different from the known PQQ-quinoprotein family during a search for AA family 8 cytochrome domains.

## Materials and Methods

### Materials

*Coprinopsis cinerea* strain 5338 was kindly provided by Dr. Yasuhiro Ito (National Food Research Institute, Ibaraki, Japan) [14]. Pyrroloquinoline quinone was purchased from Sigma-



**Figure 2. Fruiting body of *Coprinopsis cinerea*.**  
doi:10.1371/journal.pone.0104851.g002

Aldrich (Tokyo, Japan). Bovine heart cytochrome *c* was purchased from Wako Pure Chemical Industries (Osaka, Japan).

### Cloning and transcript analysis


*C. cinerea* was grown in Kremer and Wood medium [15] containing 2% Avicel (Merck, Whitehouse Station, NJ) for 12 d, and then total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany). First-strand cDNA was synthesized using SuperScript III reverse transcriptase (Invitrogen, Carlsbad, CA) with an oligo(dT) primer (Takara Bio, Shiga, Japan). PCR using KOD plus version 2 DNA polymerase (Toyobo, Osaka, Japan) with the synthesized first-strand cDNA as a template was primed with oligonucleotides flanking the predicted translational start and stop codons (20 nucleotides upstream, 5'-TCGGGACCGACCACGAACG-3'; 63 nucleotides downstream, 5'-CGATTCTGTCTTGAAGCCCGACT-3'). The PCR products were subcloned into the pGEM-T Easy vector (Promega, Madison, WI) followed by sequencing with a 3130 Genetic Analyzer (Applied Biosystems, Foster City, CA). The nucleotide sequence reported in this paper has been deposited in the DDBJ database under accession number AB901366.

### Sequence analysis

The *C. cinerea* genome database ([http://www.broadinstitute.org/annotation/genome/coprinus\\_cinereus/MultiHome.html](http://www.broadinstitute.org/annotation/genome/coprinus_cinereus/MultiHome.html)) was searched with the amino acid sequence of CDH from *P. chrysosporium* (accession no. AAB92262) using the TBLASTN algorithm [16,17]. The search was carried out with standard settings and the BLOSUM 62 matrix. The amino acid sequences of candidate genes were scanned for the presence of signal peptides using SignalP version 4.1 software [18] at the Center for Biological Sequence Analysis (<http://www.cbs.dtu.dk/services/SignalP/>) and searched against the protein database at the National Center for Biotechnology Information (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The homology structure model was constructed using Protein Homology/analogy Recognition Engine (Phyre) version 2.0 (<http://www.sbg.bio.ic.ac.uk/phyre2/>) [19]. The *CcSDH* model was used for preparing a figure in which PQQ of sGDH from *Acinetobacter calcoaceticus* and heme in the cytochrome domain of *P. chrysosporium* CDH (PDBID 1CRU and 1D7C, respectively) were superimposed to the model. The structure based alignment was created during the process of homology modeling using the Phyre2 server, and the corresponding figure was prepared using ESPript ver. 2.3 [20].

### Production and purification of recombinant protein in *P. pastoris*


The expression vector was constructed as described [21,22]. Two different oligonucleotide primers were designed based on the nucleotide sequence of the mature protein to allow for ligation into the *EcoRI* and *NotI* sites of the *Pichia* expression vector pPICZ $\alpha$  (Invitrogen, Carlsbad, CA): 5' primer (5'-TTGAATTC-CAAGGCTCTCCCACTCAGTG-3') and 3' primer (5'-TTGCGGCCGCTATGCAGGAACACACTGAGAGTACC-3'). PCR was carried out using the primer pair with KOD plus version 2 DNA polymerase (Toyobo, Osaka, Japan) and the first-strand cDNA synthesized above as a template. The PCR product was subcloned into the pGEM-T Easy Vector (Promega, Madison, WI) followed by sequencing with a 3130 Genetic Analyzer (Applied Biosystems, Foster City, CA). The target gene was digested with *EcoRI* and *NotI* and ligated into the corresponding restriction sites of the pPICZ $\alpha$  vector. The nucleotide sequence of the inserted cDNA was confirmed by sequence analysis. Approx-



**Figure 3. Domain organization of *Cc*SDH, CDH from *P. chrysosporium*, CBM1-carrying CDH from *T. heterothallica*, and cellulose-binding cytochrome *b*<sub>562</sub> from *P. chrysosporium*.** Abbreviations: AA3, Auxiliary Activities (AA) family 3 enzymes defined as flavoproteins containing a flavin-adenine dinucleotide (FAD)-binding domain; AA8, AA family 8 enzymes with the cytochrome domain of spectral class *b*; CBM1, family 1 carbohydrate-binding module. doi:10.1371/journal.pone.0104851.g003

imately 10 µg of the DNA construct in pPICZα was linearized with Bpu 1102I (Takara, Japan) prior to transformation into *P. pastoris*. Electroporation, selection of the transformant, and production of recombinant protein were carried out using an EasySelect *Pichia* expression kit (version G; Invitrogen, Carlsbad, CA) according to the manufacturer’s instruction. The PQQ domain of the recombinant protein was constructed by deleting the coding regions for the heme domain and the carbohydrate-binding module (CBM) domain of the recombinant protein by PCR using pPICZα containing the full-length gene as a template. The two oligonucleotide primers used to truncate the heme domain (221 amino acids from the N-terminus) were 5’-

CGAGAAAAGAACCTTCGTCTCTTGC-3’ as the 5’ primer and 5’-GCAAGAGACGAAGGTTCTTTTCTCG-3’ as the 3’ primer; these primers included the α-factor signal sequence of the pPICZα vector upstream of the recombinant protein gene and the gene coding for the amino acids after residue T222. For truncation of the CBM domain, a stop codon was introduced by using the following primers: 5’-CATCATCAAGCGCTAGTCCGCCCC-TATTGTTTCAGC-3’ (5’ primer) and 5’-GCTGAACAA-TAGGGCCGGACTAGCGCTTGATGATG-3’ (3’ primer). The underlined codons represent mismatches that will introduce a stop codon and prevent the translation of the last 77 C-terminal amino acids, including the CBM domain. The mutations were



**Figure 4. A, SDS-PAGE of purified (lane 1) and deglycosylated (lane 2) *Cc*SDH. B, Effect of PQQ on cytochrome *c*-reducing activity of *Cc*SDH with L-fucose as a substrate. For the activity measurement in B, 40 nM *Cc*SDH was incubated with 20 mM L-fucose, 50 µM cytochrome *c* and 1 µM PQQ in HEPES buffer (pH 7.0), and the reduction of cytochrome *c* was monitored based upon the increase of absorption at 550 nm. C and D, Isothermal titration calorimetry analysis of the PQQ domain of *Cc*SDH and PQQ with the raw data (C) and plots of the integrated peaks (D). The PQQ domain of *Cc*SDH (6.73 µM) was titrated with an initial injection of 2 µL of PQQ (46.3 µM) followed by 69 consecutive injections of 4 µL of PQQ in the presence of 1 mM CaCl<sub>2</sub> and 20 mM sodium acetate (pH 6.0). The black line indicates the best fit to a single-site model. doi:10.1371/journal.pone.0104851.g004**

**Table 1.** Kinetic parameters of CcSDH for various monosaccharides.

	$k_{cat}$ ( $s^{-1}$ )	$K_m$ (mM)	$k_{cat}/K_m$ ( $\times 10^3 s^{-1}M^{-1}$ )
D-glucosone	74.1 ( $\pm 1.4$ )	7.9 ( $\pm 0.3$ )	9.37
L-fucose	56.4 ( $\pm 1.8$ )	24.8 ( $\pm 1.2$ )	2.27
D-arabinose	35.5 ( $\pm 0.6$ )	30.3 ( $\pm 0.8$ )	1.17
L-gulose	53.3 ( $\pm 1.1$ )	84.7 ( $\pm 2.6$ )	0.63
D-lyxose	12.9 ( $\pm 0.3$ )	66.8 ( $\pm 2.6$ )	0.19

doi:10.1371/journal.pone.0104851.t001

confirmed by DNA sequencing, and the recombinant plasmid was transformed into *P. pastoris* for subsequent protein expression according to the same protocol used for the full-length recombinant protein.

The culture was centrifuged (30 min, 1,500 $\times$ g), and the crude proteins (including the recombinant protein) were precipitated with ammonium sulfate (70% saturation) from the cell-free culture. The precipitates were dissolved in 20 mM Tris-HCl buffer containing 1 M ammonium sulfate (pH 8.0). The crude proteins were fractionated on a Toyopearl Phenyl-650S column equilibrated with 20 mM Tris-HCl buffer containing 1 M ammonium sulfate (pH 8.0). The recombinant protein was eluted using a linear reverse gradient to 20 mM Tris-HCl buffer (pH 8.0). The fraction containing the recombinant protein was collected and equilibrated against 20 mM Tris-HCl buffer (pH 8.5). The solution was loaded on a Toyopearl DEAE-650S column equilibrated with 20 mM Tris-HCl (pH 8.5). The enzyme was eluted from the column using a linear gradient in the same buffer (0 to 500 mM NaCl). The purity was confirmed by SDS-PAGE analysis on 12% polyacrylamide gels, and the N-terminal amino acid sequence was determined with a protein sequencer (model 491 cLC; Applied Biosystems, Foster City, CA) as described. The recombinant protein concentration in the solution was estimated from the absorbance at 420 nm ( $\epsilon_{420} = 130 \text{ mM}^{-1} \text{ cm}^{-1}$ ). Deglycosylation of the recombinant protein was performed using endo- $\beta$ -N-acetylglucosaminidase H (endo-H) as described [21,22].

### Enzyme activity

The enzyme activity was assayed using cytochrome *c* as an electron acceptor [23,24]. The assay was performed by photometric monitoring of the time-dependent reduction of 50  $\mu$ M cytochrome *c* at 550 nm ( $\epsilon_{550} = 17.5 \text{ mM}^{-1} \text{ cm}^{-1}$ ) at 30°C. One activity unit (U) corresponds to the amount of enzyme that can convert 1  $\mu$ mol of substrate per minute. The substrate specificity of the recombinant protein (40 nM) was assessed using 1 mM D-, L-allose, D-, L-galactose, D-, L-glucose, D-, L-gulose, D-, L-mannose, D-, L-talose, D-, L-arabinose, D-, L-lyxose, D-, L-xylose, D-, L-fructose, D-, L-tagatose, L-sorbose, cellobiose, maltose, sophorose, trehalose, D-, L-fucose, L-rhamnose, N-acetyl-D-glucosamine, D-glucosamine, and D-glucosone. To obtain the kinetic parameters of oxidation for L-gulose, D-arabinose, D-lyxose, L-fucose, and D-glucosone, the assay was performed with various concentrations of the saccharides toward 50 nM recombinant protein in 50 mM Tris-HCl buffer (pH 8.5) containing 1  $\mu$ M PQQ and 1 mM  $\text{CaCl}_2$ . To determine the Michaelis constant ( $K_m$ ) and catalytic rate ( $k_{cat}$ ), the experimental data were fitted to the Michaelis-Menten equation.

### Isothermal titration calorimetry (ITC)

ITC experiments were carried out at 25°C with a MicroCal VP-ITC (GE Healthcare, Northampton, MA, USA). A sample of the purified PQQ domain of the recombinant protein was prepared by dialysis against ITC buffer (20 mM sodium acetate, pH 6.0, 1 mM  $\text{CaCl}_2$ ). The concentration of the PQQ domain of the recombinant protein was determined from the UV absorbance at 280 nm using the calculated molar extinction coefficient  $7.241 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ . PQQ was dissolved in the same ITC buffer, and the concentration was measured based upon the absorbance at 257 nm using the molar extinction coefficient [25]  $2.15 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ . Solutions were de-gassed for 1 min using a vacuum degasser immediately prior to experimentation. The titration sequence consisted of an initial injection of 2  $\mu$ L of PQQ (46.3  $\mu$ M) followed by 69 additional injections of 4  $\mu$ L of PQQ at 220 s intervals into a calorimeter cell containing the PQQ domain of the recombinant protein solution (6.73  $\mu$ M) stirred at 300 rpm. Baseline-subtracted data were fitted according to Marquardt's single-site model using Origin ITC Analysis software (Microcal Software, Northampton, MA) to obtain stoichiometric and thermodynamic parameters.

### Construction of a phylogenetic tree


The National Center for Biotechnology Information protein database was searched for amino acid sequences corresponding to the catalytic domain of the recombinant protein using the PSI-BLAST algorithm. All searches were performed with standard settings and the BLOSUM 62 matrix. The sequences obtained and the sequences of known quinoproteins (PQQ-dependent methanol dehydrogenases, ethanol dehydrogenases, soluble glucose dehydrogenases, quinoxinoprotein alcohol dehydrogenases, polyvinyl alcohol dehydrogenase, membrane-bound glucose dehydrogenase, membrane-bound alcohol dehydrogenase, and aldose sugar dehydrogenases) were subjected to alignment analysis using MAFFT (ver. 6.85) [26,27]. Non-conserved regions were trimmed using SeaView (ver. 4.4.2) software. The phylogenetic tree was generated from the trimmed sequences using ClustalX (ver. 2.1) with the neighbor-joining method (Bootstrap value, 1000).

### Results

As a result of the BLAST search, several genes, including genes homologous to the cytochrome domain of CDH, were identified in the *C. cinerea* genome, and we cloned one of them (Chromosome 6:1274274–1277156). RT-PCR using total RNA of *C. cinerea* was performed to obtain the cDNA of the whole protein. The cDNA consisted of 2883 bp, which included an open reading frame for 726 amino acids. Based on the analysis of the predicted amino acid sequence using the SignalP 4.1 server [18], the N-terminal 18 amino acid sequence was identified as a signal peptide, suggesting







**Figure 6. 3D structure model of CcSDH showing its three domains the N-terminal AA family 8 cytochrome domain (orange), the PQQ-dependent dehydrogenase domain (magenta), and the C-terminal cellulose-binding domain belonging to CBM family 1 (green).** PQQ of sGDH from *Acinetobacter calcoaceticus* and heme in the cytochrome domain of *P. chrysosporium* CDH (PDBID 1CRU and 1D7C, respectively) were superimposed to the model.  
doi:10.1371/journal.pone.0104851.g006


cytochrome domain, we tested NAD(P), FAD, and PQQ as cofactors for two-electron oxidation reaction of various sugars, alcohols, aldehydes, and their derivatives. The assay was carried out by monitoring the rate of cytochrome *c* reduction (similar to the CDH assay [23,24]). Interestingly, when various sugars (including D-glucosone, L-fucose, and some rare sugars) were used as substrates, the reduction of cytochrome *c* was observed only in the presence of PQQ. CcSDH had catalytic activity toward L-galactose, L-gulose, D-talose, D-arabinose, D-lyxose, L-fucose, and D-glucosone in the presence of PQQ and CaCl<sub>2</sub>. CcSDH shows little or no activity toward abundant sugars such as D-glucose, D-fructose, and cellobiose. The steady-state kinetic parameters for L-gulose, D-arabinose, D-lyxose, L-fucose, and D-glucosone were measured using the cytochrome *c* assay under optimal condition as summarized in Table 1. Because D-glucosone has the reductive activity of cytochrome *c*, we determined the initial velocity in an enzymatic reaction after subtracting the rate of cytochrome *c* reduction by D-glucosone itself from an apparent rate of cytochrome *c* reduction. The catalytic efficiency measured was in the following order: D-glucosone > L-fucose > D-arabinose > L-gulose > D-lyxose. D-glucosone exhibited the highest catalytic efficiency,  $k_{cat}/K_m$ , calculated to be  $9.37 \times 10^3 \text{ s}^{-1} \text{ M}^{-1}$ . The  $K_m$  values for these sugars appear to be large. However, considering the fact that the PQQ-dependent sugar dehydrogenases, such as soluble glucose dehydrogenase from *Acinetobacter calcoaceticus* (sGDH) have a large  $K_m$  value (25 mM) for glucose, a physiological substrate [28], the  $K_m$  value for D-glucosone (7.9 mM) with CcSDH is reasonable for a quinoprotein. Figure 4B shows the time-course of the absorbance at 550 nm based upon the reduced cytochrome *c* for L-fucose as a typical example. When CcSDH was added to the assay solution without PQQ, no catalytic activity was observed. Upon addition of PQQ, the absorbance immediately increased. To determine the binding constant of the protein to PQQ, the PQQ domain alone was expressed in the *Pichia* expression system and the binding constant was estimated by ITC as shown in Figure 4C. The dissociation constant ( $K_d$ ) was 1.11 nM, and the molar ratio of PQQ to the domain was 1:1 (Figure 4D).

BLAST search revealed many genes that encode enzymes similar to CcSDH in bacteria, archaea, amoebozoa, and fungi. Therefore, this type of quinoprotein appears to be widely distributed not only in prokaryotes, but also in eukaryotes. A phylogenetic tree generated from the amino acid sequences showed that these quinoproteins are distinct from other classes of known quinoproteins (Figure 5), suggesting that they represent a new family.

## Discussion

The highest amino acid homology to the unknown domain of CcSDH was obtained with *Deinococcus radiodurans* L-sorbose dehydrogenase (26%). Glucose dehydrogenase and sorbose dehydrogenase, which have the highest degree of similarity to this protein, are both PQQ-dependent dehydrogenases. The catalytic activity for various sugars was observed only in the presence of PQQ. The strong binding activity of the protein to PQQ was shown by ITC. Based on these findings, we concluded that CcSDH is a PQQ-dependent enzyme, even though it shows low amino acid sequence homology with known PQQ-dependent enzymes.

The three-dimensional structures of PQQ-dependent quinoproteins are generally divided into two types: an eight-bladed beta-propeller structure with each blade consisting of a four-stranded anti-parallel beta-sheet, and a six-bladed beta-propeller structure [3,29]. These two types of enzymes have no amino acid sequence homology with each other. The three-dimensional structure of CcSDH was modeled using the Phyre2 protein fold recognition server and the predicted structure was compared with those of other quinoproteins and structurally similar proteins. CcSDH was predicted to have a six-bladed instead of an eight-bladed quinoprotein structure (Figure 6). As shown in Table 2, CcSDH shows some structural homology with known prokaryote PQQ-dependent sugar dehydrogenases, sGDH [30–32] and soluble aldose sugar dehydrogenases (Asd) from *Pyrobaculum aerophilum* [33] and *Escherichia coli* [34], although the amino acid sequence homology is low ( $\approx 20\%$ ). Among prokaryotic sugar dehydrogenases, CcSDH shows some structural similarity with human



**Figure 7. Alignment of the amino acid sequences of the catalytic domain of CcSDH and known structure of six-bladed quinoproteins.** Perfect matches are enclosed in boxes with a black background. Boxes with a magenta background indicate the amino acid residues interacting with PQQ via direct hydrogen bonds in known structures. The filled arrowhead indicates the position of the proposed catalytic histidine in bacterial quinoproteins. Conserved residues involved in PQQ-binding in known quinoproteins are indicated by open arrowheads if also conserved in CcSDH [32–34].  
doi:10.1371/journal.pone.0104851.g007

hedgehog-interacting protein (HHIP) [35]. As seen in the alignment of the amino acid sequence of CcSDH with those of known six-bladed quinoproteins (Figure 7), a putative catalytic histidine is conserved in all proteins, whereas several amino acids contributing to PQQ binding in the prokaryote sugar dehydrogenase are missing in CcSDH and HHIP. However, as demonstrated above, the activity of CcSDH is PQQ-dependent. Thus,

the amino acids involved in the adsorption of PQQ should be different in this enzyme family, indicating the diversity of PQQ-binding motifs in eukaryotic enzymes.

The PQQ-dependent sugar dehydrogenase that we discovered here has interesting enzymatic properties associated with characteristic cytochrome (AA family 8) and cellulose-binding domains (CBM family 1) as well as the enzymes (proteins) related to plant

**Table 2.** Summary of sequence homology modeling results using the Phyre2 server.

Enzyme	Organism	PDBID	Identity	Confidence	E-value <sup>†</sup>
soluble glucose dehydrogenase (sGDH)	<i>Acinetobacter calcoaceticus</i>	1CRU	16	100.0	1.90E-34
aldose sugar dehydrogenase	<i>Pyrobaculum aerophilum</i>	3A9G	22	100.0	1.10E-28
soluble aldose sugar dehydrogenase (Asd)	<i>Escherichia coli</i>	2G8S	15	100.0	1.30E-26
human hedge hog interacting protein (HHIP)	<i>Homo sapiens</i>	2WG3	16	100.0	1.60E-27
putative glucose dehydrogenase	<i>Thermus thermophilus</i>	2ISM	16	100.0	9.00E-27
aldose sugar dehydrogenase (Adh)	<i>Streptomyces coelicolor</i>	3DAS	17	100.0	9.40E-24

<sup>†</sup>E-value generated using the Phyre2 profile-profile alignment algorithm.  
doi:10.1371/journal.pone.0104851.t002



cell wall degradation, and shows high activity towards rare sugars. Therefore, the discovery of the PQQ-dependent domain in CcSDH could form the basis for a new AA family in the CAZy database. Detailed enzymatic studies will be necessary to evaluate why eukaryotes produce and secrete such enzymes, and how these enzymes acquire PQQ, because only a few prokaryotic organisms has been reported to be capable of synthesizing PQQ. Interestingly, genes encoding enzymes similar to CcSDH were found in various bacteria, whereas only a few archaea contained corresponding genes. Furthermore, all bacterial and archaeal CcSDH-like enzymes lack the cytochrome domain present in some of the fungal enzymes. This observation raises the question as to how eukaryotes such as fungi acquired genes encoding PQQ-dependent enzymes.

## References

- Hauge JG (1964) Glucose Dehydrogenase of Bacterium Anitratum: An enzyme with a novel prosthetic group. *J Biol Chem* 239: 3630–3639.
- Salisbury SA, Forrest HS, Cruse WB, Kennard O (1979) A novel coenzyme from bacterial primary alcohol dehydrogenases. *Nature* 280: 843–844.
- Anthony C, Ghosh M (1998) The structure and function of the PQQ-containing quinoprotein dehydrogenases. *Prog Biophys Mol Biol* 69: 1–21.
- Anthony C (2001) Pyrroloquinoline quinone (PQQ) and quinoprotein enzymes. *Antioxid Redox Signal* 3: 757–774.
- Matsushita K, Toyama H, Yamada M, Adachi O (2002) Quinoproteins: structure, function, and biotechnological applications. *Appl Microbiol Biotechnol* 58: 13–22.
- Choi O, Kim J, Kim JG, Jeong Y, Moon JS, et al. (2008) Pyrroloquinoline quinone is a plant growth promotion factor produced by *Pseudomonas fluorescens* B16. *Plant Physiol* 146: 657–668.
- Killgore J, Smidt C, Duich L, Romero-Chapman N, Tinker D, et al. (1989) Nutritional importance of pyrroloquinoline quinone. *Science* 245: 850–852.
- Sütes TE, Mitchell AE, Rucker RB (2000) Physiological importance of quinoenzymes and the O-quinone family of cofactors. *J Nutr* 130: 719–727.
- Misra HS, Rajpurohit YS, Khairnar NP (2012) Pyrroloquinoline-quinone and its versatile roles in biological processes. *J Biosci* 37: 313–325.
- Harris CB, Chohanadisaï W, Mishchuk DO, Satre MA, Slupsky CM, et al. (2013) Dietary pyrroloquinoline quinone (PQQ) alters indicators of inflammation and mitochondrial-related metabolism in human subjects. *J Nutr Biochem* 24: 2076–2084.
- Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B (2013) Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnol Biofuels* 6: 41.
- Ayers AR, Ayers SB, Eriksson KE (1978) Cellobiose oxidase, purification and partial characterization of a hemoprotein from *Sporotrichum pulverulentum*. *Eur J Biochem* 90: 171–181.
- Hallberg BM, Bergfors T, Backbro K, Pettersson G, Henriksson G, et al. (2000) A new scaffold for binding haem in the cytochrome domain of the extracellular flavocytochrome cellobiose dehydrogenase. *Structure* 8: 79–88.
- Yanagi SO, Kawasumi T, Takebe I, Takemaru T (1988) Genetic analyses of *Coprinus cinereus* strains derived through intraspecific protoplast fusion. *Agricultural and Biological Chemistry* 52: 281–284.
- Kremer SM, Wood PM (1992) Evidence that cellobiose oxidase from *Phanerochaete chrysosporium* is primarily an Fe(III) reductase. Kinetic comparison with neutrophil NADPH oxidase and yeast flavocytochrome *b<sub>2</sub>*. *Eur J Biochem* 205: 133–138.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215: 403–410.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, et al. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25: 3389–3402.
- Petersen TN, Brunak S, von Heijne G, Nielsen H (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8: 785–786.
- Kelley LA, Sternberg MJ (2009) Protein structure prediction on the Web: a case study using the Phyre server. *Nat Protoc* 4: 363–371.
- Gouet P, Courcelle E, Stuart DI, Metz F (1999) ESPript: analysis of multiple sequence alignments in PostScript. *Bioinformatics* 15: 305–308.
- Yoshida M, Igarashi K, Wada M, Kaneko S, Suzuki N, et al. (2005) Characterization of carbohydrate-binding cytochrome *b<sub>562</sub>* from the white-rot fungus *Phanerochaete chrysosporium*. *Appl Environ Microbiol* 71: 4548–4555.
- Yoshida M, Ohira T, Igarashi K, Nagasawa H, Aida K, et al. (2001) Production and characterization of recombinant *Phanerochaete chrysosporium* cellobiose dehydrogenase in the methylotrophic yeast *Pichia pastoris*. *Biosci Biotechnol Biochem* 65: 2050–2057.
- Samejima M, Eriksson KE (1992) A comparison of the catalytic properties of cellobiose:quinone oxidoreductase and cellobiose oxidase from *Phanerochaete chrysosporium*. *Eur J Biochem* 207: 103–107.
- Igarashi K, Verhagen MF, Samejima M, Schulein M, Eriksson KE, et al. (1999) Cellobiose dehydrogenase from the fungi *Phanerochaete chrysosporium* and *Hemicola insolens*. A flavohemoprotein from *Hemicola insolens* contains 6-hydroxy-FAD as the dominant active cofactor. *J Biol Chem* 274: 3338–3344.
- Jongejan JA (1989) Chemistry of PQQ. Delft, The Netherlands: Delft University of Technology.
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30: 3059–3066.
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30: 772–780.
- Igarashi S, Hirokawa T, Sode K (2004) Engineering PQQ glucose dehydrogenase with improved substrate specificity. Site-directed mutagenesis studies on the active center of PQQ glucose dehydrogenase. *Biomol Eng* 21: 81–89.
- Toyama H, Mathews FS, Adachi O, Matsushita K (2004) Quinohemoprotein alcohol dehydrogenases: structure, function, and physiology. *Arch Biochem Biophys* 428: 10–21.
- Oubrie A, Rozeboom HJ, Kalk KH, Olsthoorn AJ, Duine JA, et al. (1999) Structure and mechanism of soluble quinoprotein glucose dehydrogenase. *EMBO J* 18: 5187–5194.
- Oubrie A, Rozeboom HJ, Kalk KH, Duine JA, Dijkstra BW (1999) The 1.7 Å crystal structure of the apo form of the soluble quinoprotein glucose dehydrogenase from *Acinetobacter calcoaceticus* reveals a novel internal conserved sequence repeat. *J Mol Biol* 289: 319–333.
- Oubrie A, Rozeboom HJ, Dijkstra BW (1999) Active-site structure of the soluble quinoprotein glucose dehydrogenase complexed with methylhydrazine: a covalent cofactor-inhibitor complex. *Proc Natl Acad Sci U S A* 96: 11787–11791.
- Sakuraba H, Yokono K, Yoneda K, Watanabe A, Asada Y, et al. (2010) Catalytic properties and crystal structure of quinoprotein aldose sugar dehydrogenase from hyperthermophilic archaeon *Pyrobaculum aerophilum*. *Arch Biochem Biophys* 502: 81–88.
- Southall SM, Doel JJ, Richardson DJ, Oubrie A (2006) Soluble aldose sugar dehydrogenase from *Escherichia coli*: a highly exposed active site conferring broad substrate specificity. *J Biol Chem* 281: 30650–30659.
- Bishop B, Aricescu AR, Harlos K, O’Callaghan CA, Jones EY, et al. (2009) Structural insights into hedgehog ligand sequestration by the human hedgehog-interacting protein HHIP. *Nat Struct Mol Biol* 16: 698–703.
- Kasahara T, Kato T (2003) Nutritional biochemistry: A new redox-cofactor vitamin for mammals. *Nature* 422: 832.