



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

York N. Hsiang, MB, ChB, MHSc, FRCSC

Department of Surgery
University of British Columbia
Vancouver, BC, Canada

A. Srikantha Phani, PhD

Department of Mechanical Engineering
University of British Columbia
Vancouver, BC, Canada

REFERENCE

1. Gioia G, Chakraborty P. Turbulent friction in rough pipes and the energy spectrum of the 85 phenomenological theory. *Phys Rev Lett* 2006;96:044502.

<https://doi.org/10.1016/j.jvs.2022.06.086>

Searching for the reason why the results of infrarenal AAA open surgery are worsening



Pomy et al¹ compared the 30-day outcomes of open surgical repair (OSR) of infrarenal abdominal aortic aneurysms (IAAAs) from 2005 to 2010 and 2014 to 2019. They found that the outcomes were worse for both elective and emergency OSR in the latter period, despite the improvements in the preoperative assessments and advances in anesthesiology and intensive care techniques. In particular, they reported a significant increase in 30-day mortality for both the elective group (3.7% vs 3.2%; $P = .006$) and the ruptured cohort (41.4% vs 40%; $P < .0001$). Considering the limitations of a retrospective data review, their conclusions have raised important questions regarding the reasons for such a deterioration in the results.

The first consideration is that the worse outcomes could have been because in recent years many vascular surgeons have often opted for endovascular repair instead of performing OSR, not only for more anatomically complex IAAs, but also for standard IAAs in patients fit for OSR. In addition, the recommendation of an endovascular first approach for ruptured IAAs was determined from observational studies; however, randomized controlled trials did not conclude the superiority of endovascular repair.²⁻⁵

The second consideration is that small series can offer better results, such as a 30-day mortality rate of 12% in a series of 75 patients with ruptured IAAs.⁶ It could be interesting to investigate the reason for such discrepancies.

The last consideration is that the poor training of the newer generations of vascular surgeons for open surgical techniques is taking its toll. It has been documented that younger vascular surgeons have reduced skills in open vascular surgery. Thus, it is crucial to offer balanced training for two equally effective and valid techniques such as open and endovascular repair.

As claimed by the ancient philosopher Aristotle, the solution will probably be “in medio stat virtus” (virtue is found in the mean).

Antonio Bozzani, MD

UOC Chirurgia Vascolare
Fondazione IRCCS Policlinico San Matteo
Pavia, Italy

Rosa Rossini, MD

UOC Chirurgia Vascolare
Ospedale Maggiore di Crema
Crema, Italy

Vittorio Arici, MD

UOC Chirurgia Vascolare
Fondazione IRCCS Policlinico San Matteo
Pavia, Italy

REFERENCES

1. Pomy BJ, Devlin J, Lala S, Amdur RL, Ricotta JJ, Sidawy AN, et al. Comparison of contemporary and historical outcomes of elective and ruptured open abdominal aortic aneurysm repair. *J Vasc Surg* 2022;75:543-51.
2. Hinchliffe RJ, Bruijstens L, MacSweeney ST, Braithwaite BD. A randomised trial of endovascular and open surgery for ruptured abdominal aortic aneurysms: results of a pilot study and lessons learned for future studies. *Eur J Vasc Endovasc Surg* 2006;32:506-13.
3. Reimerink JJ, Hoornweg LL, Vahl AC, Wisselink W, van den Broek TA, Legemate DA, et al. Endovascular repair versus open repair of ruptured abdominal aortic aneurysms: a multicenter randomized controlled trial. *Ann Surg* 2013;258:248-56.
4. Powell JT, Sweeting MJ, Thompson MM, Ashleigh R, Bell R, Gomes M, et al. Endovascular or open repair strategy for ruptured abdominal aortic aneurysm: 30-day outcomes from IMPROVE randomized trial. *BMJ* 2014;348:f7661.
5. Desgranges P, Kobeiter H, Katsahian S, Bouffi M, Gouny P, Favre JP, et al; ECAR Investigators. ECAR (Endovasculaire ou Chirurgie dans les Anévrysmes aorto-iliaques Rompus): a French randomized controlled trial of endovascular versus open surgical repair of ruptured aorto-iliac aneurysms. *Eur J Vasc Endovasc Surg* 2015;50:303-10.
6. Arici V, Bozzani A, Rossi M, Corbetta R, Brunetto MB, Scudeller L, et al. Contemporary early and long-term results of open repair for ruptured and symptomatic unruptured infrarenal AAA: single center experience. *Ann Vasc Surg* 2020;64:99-108.

<https://doi.org/10.1016/j.jvs.2022.03.898>

From coronavirus disease 2019 to long coronavirus disease 2019 in vascular pathology



The interesting report by Faries et al¹ concerning the frequent complications in patients with coronavirus disease 2019 (COVID-19) who had been treated for arterial thrombosis merits some further discussion. SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), a multisystem disease, after an acute phase, can become protracted as prolonged COVID-19, lasting for 12 weeks, or for longer as long COVID-19. Medical treatments have often been unable to completely eradicate the viral load, neutralize all the negative effects, and restore the damaged physiologic function. In particular, COVID-19




Fig. Flowchart showing pathophysiologic steps from coronavirus disease 2019 (COVID-19) to long COVID-19.

micro-organisms can penetrate into the arterial, venous, and capillary endothelia, causing accelerated pyroptosis and generating or aggravating corresponding endothelial dysfunction. In capillaries, the parietal infiltration of inflammatory cells will cause endotheliitis and subsequent microthromboses, with a large production of cytokines, typically interleukin-6.² Moreover, interrelating B and T lymphocytes will activate plasma cells to produce antibodies, autoantibodies, and complement anaphylatoxins (C3a and C5a), sometimes in a dysregulated manner, leading to a type 3 hypersensitive vasculitis and a more dangerous necrotizing leukocytoclastic vasculitis. An often associated secondary antiphospholipid-like syndrome, together with increased production of platelets from megakaryocytes, also present in the lung parenchyma, can generate an immune-correlated hypercoagulable state (Fig).³⁻⁵ In medium- to large-size vessels, including coronary arteries, this results in an increased risk of thrombosis and, through a mechanism of platelet adhesion and deposition of inflammatory protein molecules, a rapid increase in atherosclerotic plaque formation. Aneurysms can increase in size owing to activated metalloproteinases, and typically in the aorta, can result in vasa vasorum thrombosis, increasing the risk of rupture.⁶ Similarly, in cardiomyocytes, COVID-19 virions will accelerate their apoptosis, followed by myocardial invasion of inflammatory cells, typically lymphocytes. The consequent myocarditis favors the occurrence of dysrhythmia, decreases the ventricles' ejection fraction, and can lead to a secondary myocardial fibrosis. Moreover, the

pulmonary lesions that can occur after COVID-19 owing to multiple areas of fibrosis can increase pulmonary resistance and the work of the right cardiac chambers.⁷

In conclusion, COVID-19 infection should be considered a disease not limited to an acute phase but with possible unpredictable long-term sequelae, correlating with a hyperregulated or dysregulated immune condition and a hypercoagulable state. Because of these factors, in addition to the often unknown power of the immune reaction of each patient, precise clinical studies should be performed before any decision for surgery, with close postoperative monitoring, especially for those with pre-existing cardiovascular risk factors.

Alberto Farinetti, MD
Antonio Manenti, MD
Gianrocco Manco, MD

Department of Surgery
University of Modena and Reggio Emilia
Polyclinic Hospital
Modena, Italy

Luca Roncati, MD

Department of Pathology
University of Modena and Reggio Emilia
Polyclinic Hospital
Modena, Italy

Francesca Coppi, MD
Anna Vittoria Mattioli, MD

Department of Cardiology
University of Modena and Reggio Emilia
Polyclinic Hospital
Modena, Italy

REFERENCES

1. Faries CM, Rao A, Ilonzo N, Hwang S, Krishnan P, Farhan S, et al. Follow-up after acute thrombotic events following COVID-19 infection. *J Vasc Surg* 2022;75:408-15.e1.
2. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet* 2020;395:1417-8.
3. Roncati L, Ligabue G, Fabbiani L, Malagoli C, Gallo G, Lusenti B, et al. Type 3 hypersensitivity in COVID-19 vasculitis. *Clin Immunol* 2020;217:108487.
4. Roncati L, Ligabue G, Nasillo V, Lusenti B, Gennari W, Fabbiani L, et al. A proof of evidence supporting abnormal immunothrombosis in severe COVID-19: naked megakaryocyte nuclei increase in the bone marrow and lungs of critically ill patients. *Platelets* 2020;31:1085-9.
5. Roncati L, Manenti A, Manco G, Farinetti A, Mattioli AV. COVID-19 thromboembolic complications: deepening immunoinflammatory features. *J Vasc Surg* 2021;74:1048-9.
6. Roncati L, Manenti A, Farinetti A, Manco G, Mattioli AV. Toward a unified pathophysiology in COVID-19 acute aortopathies. *J Vasc Surg* 2021;74:1771-2.
7. Guzik TJ, Mohiddin SA, Dimarco A, Patel V, Sawatis K, Marelli-Berg FM, et al. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovasc Res* 2020;116:1666-87.

<https://doi.org/10.1016/j.jvs.2022.03.897>