



Biosynthesis of Metal Nanoparticles via Microbial Enzymes: A Mechanistic Approach

Muhammad Ovais ^{1,2,*}, Ali Talha Khalil ³, Muhammad Ayaz ⁴, Irshad Ahmad ⁵, Susheel Kumar Nethi ⁶ and Sudip Mukherjee ^{7,*}

- ¹ CAS Key Laboratory for Biomedical Effects of Nanomaterials and Nanosafety, CAS Center for Excellence in Nanoscience, National Center for Nanoscience and Technology (NCNST), Beijing 100190, China
- ² University of Chinese Academy of Sciences, Beijing 100049, China
- ³ Department of Eastern Medicine and Surgery, Qarshi University, Lahore 54000, Pakistan; talhakhalil.qau@gmail.com
- ⁴ Department of Pharmacy, University of Malakand, Khyber Pakhtunkhwa (KPK), Chakdara 18000, Pakistan; ayazuop@gmail.com
- ⁵ Department of Life sciences, King Fahd University of Petroleum and Minerals (KFUPM), Dhahran 31261, Saudi Arabia; irshad@kfupm.edu.sa
- ⁶ Department of Experimental and Clinical Pharmacology, College of Pharmacy, University of Minnesota, Minneapolis, MN 55455, USA; susheel.nethi@gmail.com
- ⁷ Department of Bioengineering, Rice University, Houston, TX 77030, USA
- * Correspondence: movais@bs.qau.edu.pk (M.O.); sudip.mukherjee@rice.edu (S.M.); Tel.: +86-185-1347-5504 (M.O.); +1-702-957-9891 (S.M.)

Received: 30 November 2018; Accepted: 17 December 2018; Published: 18 December 2018



Abstract: During the last decade, metal nanoparticles (MtNPs) have gained immense popularity due to their characteristic physicochemical properties, as well as containing antimicrobial, anti-cancer, catalyzing, optical, electronic and magnetic properties. Primarily, these MtNPs have been synthesized through different physical and chemical methods. However, these conventional methods have various drawbacks, such as high energy consumption, high cost and the involvement of toxic chemical substances. Microbial flora has provided an alternative platform for the biological synthesis of MtNPs in an eco-friendly and cost effective way. In this article we have focused on various microorganisms used for the synthesis of different MtNPs. We also have elaborated on the intracellular and extracellular mechanisms of MtNP synthesis in microorganisms, and have highlighted their advantages along with their challenges. Moreover, due to several advantages over chemically synthesized nanoparticles, the microbial MtNPs, with their exclusive and dynamic characteristics, can be used in different sectors like the agriculture, medicine, cosmetics and biotechnology industries in the near future.

Keywords: metal nanoparticles; microbial flora; biosynthesis; microbial enzymes; action mechanism

1. Background and Role of Microbial Enzymes in Metal Nanoparticle (MtNP) Biosynthesis

Nanotechnology has accomplished enormous development during the last decades due to the synthesis and varied applications of metal nanoparticles (MtNPs) in different areas, such as biology, food, agriculture, engineering, electronics, cosmetics, medicine, and in food and biomedical devices. MtNPs have reached a momentous position due to their specific physicochemical characteristics and significant biotechnological applications [1,2]. According to a market report, the worldwide production of MtNPs is currently valued at 13.7 billon US dollars, which is expected to reach 50 billon US dollars by 2026. The wide-ranging use of MtNPs has significantly contributed to robust development in the

macroeconomic industry and the demand for MtNPs will continue to be high in three regions of the world: North America, Western Europe and the Asia-Pacific region [3].

During the last decade, many researchers have synthesized MtNPs by orthodox physical and chemical methods. The disadvantages of the physical method include expensive synthesis and little yield. Similarly, the chemical methods are unsafe due to the involvement of hazardous chemical substances that are attached to the surface of MtNPs, which possess detrimental side effects in biomedical applications. Keeping in view the aforementioned concerns, research has shifted towards the synthesis of MtNPs using biological constituents which are economical, biocompatible, non-toxic and eco-friendly [4–7]. The biological synthesis of MtNPs is mostly carried out by utilizing different types of plants and microorganisms. The phytogenic process of MtNP synthesis is economical and relatively simple. However, this process generates polydispersed nanoparticles, due to its diverse photochemistry [8,9]. In contrast, the microorganisms are considered as prospective bio-factories for the green synthesis of MtNPs, something that has gained vast consideration in recent times, as they are indispensable and of technological importance. This is due to the broad range of microorganisms that react in a different way with the metal ions for MtNP synthesis. Microbial MtNPs of different sizes and shapes have been reported in various species of bacteria, fungi and yeasts due to their enhanced growth rate, easy cultivation and their capability to grow under established conditions of temperature, pH and pressure [10,11]. The biosynthesis of MtNPs and their alloys (gold, silver, gold-silver alloy, selenium, tellurium, platinum, palladium, silica, titania, zirconia, quantum dots, magnetite and uraninite) has been reported in different microorganisms, including bacteria, actinomycetes, fungi, yeasts and viruses.

Different microorganisms have the ability to synthesize inorganic materials via both intracellular and extracellular methods. In the intracellular approach, the microbial cell is comprised of an amazing ion transport system. Due to electrostatic interaction, the bacterial cell wall which is negatively charged attracts the positively charged metal ions. Additionally, the bacterial cell wall contains enzymes that reduce the metal ions to their respective nanoparticles. Whereas in extracellular method, the microbial cell secretes reductases used in the bioreduction of metal ions into the corresponding MtNPs [12]. The following sections of this article will provide an overview on the green synthesis of MtNPs by using potential microbial flora, including bacteria, cyanobacteria, microalgae, actinomycetes, yeasts, fungi, viruses and diatoms. Furthermore, the intracellular and extracellular mechanisms that are adopted by the microorganism and the microbial enzymes for MtNP synthesis will be elaborated upon, along with the contemporary challenges and future prospects. Table 1 highlights the recent literature about the microbe assisted synthesis of MtNPs and their applications.

2. Biosynthesis of MtNPs by Microorganisms

Synthesizing uniform, ultrafine, well-dispersed functional nanoparticles under normal conditions through a controlled manner remains a great challenge [13,14]. This has caused a major surge in looking for alternative means of synthesizing nanoparticles that are devoid of such disadvantages. In recent years, biological resources have been frequently explored for the biosynthesis of metal or metal-based nanoparticles. These biological resources usually provide a versatile, economical and eco-friendly method to fabricate metal nanoparticles [14,15] that exhibit interesting physical, chemical and biological properties. Furthermore, other advantages such as the ease of production and scaling, well defined morphologies and enhanced biocompatibility (relative to the physiochemical based nanoparticles) have attracted many scientists to use such resources as nanofactories [16,17]. The bio-based methods for the synthesis of metal nanoparticles are based on the systematic use of plant extracts and microorganisms, like bacteria, yeasts, and fungi [18–22]. In biological synthesis, no capping or stabilizing agents are added as the biomolecules can perform this function themselves [23–25]. The properties of these nanoparticles (such as shape, size, etc.), are governed by these biomolecules [13]. These biomolecules also functionalize the nanoparticles, making it more effective relative to nanoparticles synthesized through nonbiological means [17].

The use of plants for biosynthesis has been discussed in detail previously by our own group as well as other scientists across the world [19,20,26,27]. The phytogenic and microgenic biosynthesis of nanoparticles both have their own sets of advantages and disadvantages. Phytogenic synthesis is time economical and relatively simple but usually leads to polydispersed nanoparticles, due to the involvement of diverse phytochemicals like phenols, flavonoids, terpenoids, etc. [8,9]. Moreover, seasonal variations can alter the phytochemical profile of the extracts used for biosynthesis [28]. On the contrary, microbial synthesis is devoid of such disadvantages, however, it requires the maintainance of a sterilized environment and culture conditions, making it relatively complex [29]. Herein, we have discussed the potential of using the microbial world as nanofactories for the biosynthesis of metal nanoparticles or metal based composite materials, both from functional and mechanistic perspectives. Microorganisms hold an exciting amount of potential for the biosynthesis of nanoparticles as the synthesis is eco-friendly and is devoid of the use of hazardous chemicals. Microorganisms are cost effective and do not have high energy requirements. In addition, they can accumulate and detoxify heavy metals through reductase enzymes, which reduce metal salts to their corresponding metal nanoparticle with less polydispersity and a narrow size range [16].

3. Bacterial and Cyanobacterial Biosynthesis of MtNPs

Bacterial cells have been readily employed as nanofactories for the synthesis of various metal Both the extracellular and intracellular approaches have been demonstrated. nanoparticles. Extracellular biosynthesis occurs outside the bacterial cell after applying diverse techniques, such as (a) using of the bacterial biomass, (b) using the supernatant of bacterial cultures and (c) using cell free extracts. Extracellular synthesis is preferred over intracellular synthesis because it is devoid of of complex downstream processing [28]. These nanoparticles have been used in a wide array of applications, however mostly for biomedical uses. Recently, silver nanoparticles (AgNPs) that were synthesized using Bacillus brevis showed excellent antimicrobial activities against the multidrug resistant strains of Staphylococcus aureus and Salmonella typhi [30]. Pseudomonas stutzeri is another bacterial strain which was found capable of accumulating AgNPs using an intracellular mechanism [31]. The *Bacillus* sp. was also found to synthesize silver nanoparticles in the intracellular periplasmic space [32]. In one study, two different isolated strains of Pseudomonas aeruginosa were used for the biosynthesis of gold nanoparticles (AuNPs), generating different sizes of AuNPs [33]. Spherical (10-50 nm) and triangular plate (50-400 nm) AuNPs were produced using Rhodopseudomonas capsulate [34]. Serratia ureilytica mediated the as-synthesized ZnO nanoflowers that were used on cotton fabrics to provide antimicrobial effects against S. aureus and E. coli [35]. Lactobacillus plantarum was also reported for the biosynthesis of ZnO nanoparticles [36]. Aeromonas hydrophila is a gram-negative bacterial strain that is used for the biosynthesis and antimicrobial applications of ZnO nanoparticles [37]. Recently, triangular CuO nanoparticles were produced using Halomonas elongate, and their antimicrobial activity was confirmed against *E. coli* and *S. aureus* [38]. In another recent report, super paramagnetic iron oxide nanoparticles (29 nm) were produced using Bacillus cereus and they were reported for their anti-cancer effects against the MCF-7 and 3T3 cell lines in a dose-dependent manner [39]. Furthermore, bimetallic Ag-Au nanostructures have also been demonstrated using bacterial strains [40]

Numerous researchers have used cyanobacteria for the biosynthesis of nanoparticles [41]. In one study, eight cyanobacterial strains were screened to investigate their potential for the biosynthesis of AgNPs. The results indicated that seven cyanobacterial strains were capable of producing AgNPs under light, however, only three were capable of producing AgNPs in the dark [42]. The ammonia-sensing potential of the cyanobacterial synthesized silver nanoparticles was also reported [43]. Highly monodispersed AgNPs (5–6.5 nm) were reported through the marine cyanobacterium *Phormidium fragile* [44]. The Nostoc species was also used for the biosynthesis of AgNPs, which indicated significant cytotoxic activity against MCF-7 anti-cancer cell lines, as well as good antimicrobial activities [45]. The intracellular synthesis of AuNPs was performed using *Lyngbya*

majuscule, which was isolated from the Arabian Gulf. The authors also revealed the interesting role of the use of AuNPs in combination with *Lyngbya majuscule* as an anti-myocardial infraction agent [46].

4. Mycosynthesis of Nanoparticles

Myco-nanotechnological approaches have been successfully applied for the biosynthesis of different metal nanoparticles. Likewise, in bacteria/cyanobacteria, the biosynthesis can be intracellular or extracellular. In intracellular synthesis, metal salts are converted into a less toxic form in the mycelia, which can be used by the fungi [47]. Extracellular biosynthesis includes the use of fungal extracts [48]. Fungi are relatively more resourceful then bacteria in the biosynthesis of nanoparticles due to the presence of a number of bioactive metabolites, high accumulation and enhanced production [16,49,50]. Recently, different filamentous fungi were reported to be proficient in the biosynthesis of AuNPs. This study employed different methods for the biosynthesis of AuNPs. The authors proposed that the fungal compounds and fungal media components potentially played a role in stabilizing the nanoparticles [47]. In another research article, three different strains of fungi (namely Aureobasidium pullulans, Fusarium oxysporum and the Fusarium sp.) were used to biosynthesize AuNPs. The authors indicated that the biosynthesis occurred in fungi vacuoles and that reducing sugars were involved to tailor spherical AuNPs. They also established the role of specific fungal proteins in the capping of the AuNPs [51]. A recent study indicated that *Rhizopus stolonifera* extracts mediated the synthesis of monodispersed AgNPs (9.4 nm), while condition optimization resulted in 2.86 nm AgNPs [52]. The extracellular synthesis of AgNPs using Candida glabrata indicated good antimicrobial potential [53]. Aspergillus niger mediated ZnO nanoparticles indicated excellent antibacterial potential, while they were also able to degrade the Bismarck brown dye by up to 90% [54]. Recently, cobalt oxide nanoparticles were produced using Aspergillus nidulans [55]. The nanoparticles were characterized via x-ray diffraction analysis (XRD), transmission electron microscopy (TEM), Fourier transform infrared spectroscopy (FTIR) and energy dispersive x-ray analysis (EDX) techniques, which revealed their spherical shape and average size of 20.29 nm. Interestingly, biogenic cobalt oxide nanoparticles have the potential to be applied in energy storage, lithium-ion batteries and in gas sensing, as well as in medicine.

5. Algae as Biosynthesis Factories

The use of algae is also increasingly popular for the biosynthesis of nanoparticles. *Sargassum muticum* was used for the biosynthesis of ZnO nanoparticles and was reported to decrease angiogenesis along with apoptotic effects in HepG2 cells [56]. AgNPs that were biosynthesized using *Gelidium amansii* indicated excellent antimicrobial properties by forming a diverse biofilm to combat bacterial strains [14]. *Sargassum crassifolium*, a macroalgae and sea weed, has been used in the biosynthesis of AuNPs. Furthermore, the authors observed a blue shift in the UV absorption after increasing the concentration of *S. crassifolium*, which was attributed to a decreased size due to increased nucleation centers in the reductant [57]. *Cystoseira trinodis* was used for the biosynthesis of CuO nanoparticles (7 nm) and was reported to have enhanced antibacterial activities and possess significant potential as an antioxidant, degrading the methylene blue [58]. Aluminum oxide nanoparticles (~20 nm) were produced using *Sargassum ilicifolium* [59]. Different algae strains were reported for the biosynthesis of gold nanoparticles, namely *Turbinaria conoides* [60], *Sargassum tenerrimum* [61], *Acanthophora spicifera* [62], *Laminaria japonica* [63], etc. These have been used for the biosynthesis of AuNPs. Novel core (Au)-shell (Ag) nanoparticle synthesis has also been reported using *Spirulina platensis* [64].

Serial Number Microorganisms		Nanoparticle Size/Shape		Application	Reference	
			Bacteria			
1	Actinobacter	Ag	13.2 nm/Spherical	Antibacterial	[65]	
2	Acinetobacter	Au	19 nm/Spherical-triangular-polyhedral	-	[66]	
3	Klebsiella pneumonia	Au	10–15 nm/Spherical	Antibacterial	[67]	
4	Sinomonas mesophila	Ag	4–50 nm/Spherical	Antibacterial	[68]	
5	Pseudomonas fluorescens	Au	5–50 nm/Spherical	Antibacterial	[69]	
6	Bacillus endophyticus	Ag	5.1 nm/Spherical	Antimicrobial	[70]	
7	Bacillus brevis	Ag	41–68 nm/Spherical	Antibacterial	[30]	
8	Streptomycesgriseoplanus	Ag	19.5–20.9 nm/Spherical	Antifungal	[71]	
9	Nocardiopsis flavascens	Ag	5 and 50/Spherical	Cytotoxicity	[72]	
10	Caldicellulosiruptor changbaiensis	Au	<20 nm/Spherical	Antibacterial, Antibiofilm	[73]	
11	Shewanella loihica	Cu	10–16 nm/Spherical	Antibacterial	[74]	
12	Shewanella loihica	Pt	1–10 nm/Spherical	Dye degradation	[75]	
13	Shewanella loihica	Pd	1–12 nm/Spherical	Dye degradation	[75]	
14	Shewanella loihica	Au	2–15 nm/Spherical	Dye degradation	[75]	
15	Micrococcus yunnanensis	Au	53.8 nm/Spherical	Antibacterial, Anticancer	[76]	
16	Mycobacterium sp.	Au	5–55 nm/Spherical	Anticancer	[77]	
17	Halomonas salina	Au	30–100 nm/Spherical	-	[78]	
			Fungi			
18	Aspergillus niger	ZnO	53–69 nm/Spherical	Antibacterial Dye degradation	[54]	
19	Trametes trogii	Ag	5–65 nm/Spherical- Ellipsoidal	-	[79]	
20	Trichoderma longibrachiatum	Ag	10 nm/Spherical	Antifungal against phyto-pathogenic fungi		
21	Trichoderma harzianum	Au	32–44 nm/Spherical	Antibacterial, Dye degradation		
22	Fusarium oxysporum	Ag	21.3–37 nm/Spherical	Antimicrobial		
23	Pleurotus ostreatus	Au	10–30 nm/Spherical	Antimicrobial, Anticancer		
24	Aspergillus terreus	Ag	16–57 nm/Spherical	Antibacterial		
25 26	Ganoderma sessiliforme Phenerochaete chrysosporium	Ag Ag	~45 nm/Spherical 34–90 nm/Spherical-Oval	Antibacterial, Antioxidant, Anticancer Antibacterial	[85] [86]	
27	Penicillium polonicum	Ag	10–15 nm/Spherical	Antibacterial	[87]	
28	Candida glabrata	Ag	2–15 nm/Spherical	Antibacterial	[53]	

Table 1. Studies from 2016 to date, indicating	different microorganisms for	or nanoparticle biosynthesis.

Serial Number	Microorganisms	Nanoparticle	Size/Shape	Application	Reference
			Fungi		
29	Macrophomina phaseolina	Ag/AgCl	5–30 nm/Spherical	Antibacterial	[88]
30	Aspergillus nidulans	CoO	20.29 nm/Spinel	-	[55]
31	Rhodotorula glutinis	Ag	15.45 nm/Spherical	Antifungal, Dye degradation, Cytotoxicity	[89]
32	Rhodotorula mucilaginosa	Ag	13.70 nm/Spherical	Antifungal, Dye degradation, Cytotoxicity	[89]
33	Cladosporium sp.	Ag	24 nm/Spherical	Antioxidant, Antidiabetic, Anti-Alzheimer	[90]
34	Cladosporium cladosporioides	Au	60 nm/Round	Antioxidant, Antibacterial	[91]
35	Nemania sp.	Ag	33.52 nm/Spherical	Antibacterial	[92]
36	Penicillium chrysogenum	Pt	5–40 nm/Spherical	Cytotoxicity	[93]
37	Aspergillus sp.	Au	2.5–6.7 nm/Spherical	Nitrophenol reduction	[94]
38	Rhizopus stolonifer	Ag	2.86 nm/Spherical	-	[52]
			Algae/Cyanobacteria		
39	Sargassum wightii	ZrO ₂	18 nm/Spherical	Antibacterial	[95]
40	Neochloris oleoabundans	Ag	40 nm/Spherical	Antibacterial	
41	Cystoseira baccata	Au	8.4 nm/Spherical	Anticancer	
42	Stephanopyxis turris	Au	10–30 nm/Spherical	-	
43	Galaxaura elongate	Au	3.85–77 nm/Spherical-rods-triangular	Antibacterial	
44	Chlorella vulgaris	Pd	5–20 nm nm/Spherical	-	
45	Enteromorpha compressa	Ag	4–24 nm/Spherical	Antimicrobial, Anticancer	
46	Nostoc linckia	Ag	5–60 nm/Spherical	Antibacterial	
47	Nostoc sp	Ag	51–100 nm/Spherical	Spherical	[45]
48	Leptolyngbya	Ag	5–50 nm/Spherical	Antibacterial, Anticancer	
49	Spyridia fusiformis	Ag	5–50 nm/Spherical	Antibacterial	
50	Chlorella pyrenoidosa	CdSe QD	4–5 nm	Imatinib sensing	
52	Sargassum ilicifolium	Al ₂ O ₃	20 nm/Spherical		
53	Padina pavonia	Ag	49.58–86.37 nm/spherical-triangular-rectangle-polyhedral-hexago		
53	Spirulina platensis	Pd	10–20 nm/Spherical	Adsorbent	[107]
54	Chlorella pyrenoidosa	TiO ₂	50 nm/Spherical	Dye degradation	[108]

Table 1. Cont.

6. Mechanisms of MtNP Synthesis by Microorganisms

The synthesis of nanosized substances using microbial cells is an emerging trend in the field of nanotechnology. Microbes including bacteria, fungi, viruses, actinomycetes and yeasts act as potential biofactories for the reduction of silver, gold, gold-silver alloy, cadmium, selenium, magnetite, silica, platinum, titania, palladium and other metals to their subsequent nanoparticles for biological applications [109]. Microbes synthesize these nanoparticles either extracellularly or intracellularly using various bio-reduction processes (Table 2).

7. Extracellular Enzymes

Extracellular microbial enzymes are known to play a significant role as reducing agents in the production of MtNPs [110] (Figure 1). Studies suggest that cofactors such as nicotinamide adenine dinucleotide (NADH) and reduced form of Nicotinamide adenine dinucleotide phosphate (NADPH) dependent enzymes both play vital roles as reducing agents via the transfer of the electron from NADH by NADH-reliant enzymes, which act electron carriers [111]. The extracellular synthesis of AuNPs by the bacterium *Rhodopseudomonas capsulata* is mediated via the secretion of NADH and NADH-reliant enzymes. The bioreduction of gold is initiated via electron transfer from NADH by NADH-reliant reductase enzymes present in *R. capsulata*. Consequently, gold ions accept electrons and get reduced (Au³⁺ to Au⁰), leading to the formation of gold nanoparticles [34]. Several other factors, including the concentration of the predecessor, the pH, the temperature and the duration of reaction are limiting factors in controlling the size of MtNPs. Beside these enzymes, several compounds, including naphthoquinones, anthraquinones and hydroquinones are involved in the production of MtNPs [19]. Microbes utilize various mechanisms for the synthesis of NPs, including changes in solubility, biosorption, metal complexation, extracellular precipitation, toxicity via oxidation-reduction, the absence of specific transporters and efflux pumps [19,112].




Figure 1. Role of NADH and NADH-dependent microbial enzymes in the synthesis of metal nanoparticles (MtNPs).

Several fungi produce extracellular enzymes like acetyl xylan esterase, cellobiohydrolase D, glucosidase and β -glucosidase, which are known to play a significant role in the biosynthesis of MtNPs [15]. One mechanism involved in the extracellular synthesis of AgNPs is the use of nitrate reductase which is secreted by fungi, which helps in the bioreduction and synthesis of MtNPs. Several studies reported the involvement of nitrate reductase in the extracellular synthesis of MtNPs [113,114]. Studies involving the use of commercially available nitrate reductase disks revealed that these NADH-reliant reductase enzymes were involved in the reduction of Ag^+ ions to $Ag^{(0)}$ and the subsequent formation of silver nanoparticles [115,116]. Fusarium oxysporum was used as source of reducing agents for the synthesis of gold and silver NPs. Results showed that extracellular reductases produced by the fungi caused the reduction of Au³⁺ and Ag¹⁺ to Au–Ag alloy NPs. Moreover, nitrate-reliant reductases and shuttle quinone obtained from various species of this fungi were utilized in the extracellular synthesis of NPs [117]. However, some species like *Fusarium moniliforme* failed to generate AgNPs, even upon the release of the reductase, indicating the Ag¹⁺ reduction via the involvement of conjugated oxidation-reduction reactions of electron carriers involving NADP-reliant nitrate reductase [116]. Furthermore, nitrate reductase from F. oxysporum was used in an in vitro study in oxygen free conditions in the presence of a cofactor (NADPH), a stabilizer protein (phytochelatin) and an electron carrier (4-hydroxyquinoline) in order to synthesize AgNPs. This fungi exhibited good extracellular production of AgNPs and can be considered an excellent candidate for the extracellular synthesis of other MtNPs [113,118]. Yet in other studies, F. oxysporum was used for the extracellular synthesis of semiconductor CdS nanoparticles, where extremely luminescent CdSe nanoparticles were synthesized using the reductase enzyme of the fungi [114,119]. Enzymes from other fungal stains, including *Fusarium semitectum* and *Fusarium solani*, were used for the extracellular production of AgNPs. The results of the study revealed that specific proteins might be responsible for the reduction of Ag⁺, thus forming AgNPs [120,121]. Cladosporium cladosporioides and Coriolus versicolor were effectively used for the extracellular synthesis of AgNPs that involved fungal proteins, organic acids and polysaccharides which effect the growth and shape of the nanocrystals [122]. Subsequent to incubation of Aspergillus niger in a AgNO₃ solution, the extracellular production of AgNPs was stabilized by fungal proteins [123]. Likewise, Aspergillus fumigatus extracellularly produced AgNPs in exceptionally less time (10 min), as compared to other physical and chemical techniques [124]. Thus, A. funigatus is an ideal candidate for industrial scale production of a variety of NPs. Penicillium fellutanum was also observed to reduce Ag¹⁺ ions in a very short amount of time (10 min). Further studies revealed that a protein of nitrate reductase was responsible for the reduction of Ag^{1+} ions [125]. *Penicillium brevicompactum* was reported to cause the reduction of Ag¹⁺ ions via the liberation of NADH-reliant enzyme nitrate reductases [126].

Nanotechnology was also applied to a new group of the plant kingdom known as algae. Among the algae, *Sargassum wightii greville* was reported to rapidly reduce Au³⁺ ions to form AuNPs 8–12 nm in size [127]. Another filamentous algae, *Chlorella vulgaris*, was used in the biosynthesis of gold nanoparticles, resulting in the formation of Au and Au⁺¹S nanoparticles [128].

Source of Extracellular Enzymes (Bacterial Species)	Nature of Organism	Metals Used	Shape	Size (nm)	Temperature (°C)	Reference
Desulfovibrio desulfuricans	G-ive Bacteria	Pd	Round	50	25	[129]
Pyrobaculum islandicum	G-ive Rods	U, Tc, Cr, Co, Mn	Round	NA	100	[130]
Escherichia coli	G-ive Bacteria	CdTe	Round	2–3.2	37	[131]
Escherichia coli	G-ive Bacteria	ive Bacteria Au		20-30	37	[132]
Bacillus licheniformis	G+ive mesophilic bacteria	Ag	NA	50	37	[133]
Shewanella species	Marine Bacteria	Se	Round	181	30	[134]
Ureibacillus thermosphaericus	G+ive Bacteria	Au	NA	50-70	60-80	[135]
Corynebacterium glutamicum	G+ive Bacteria Ag		Irregular	5-50	30	[136]
Rhodopseudomonas capsulate	Phototrophic Bacteria Au		Round	10-20	30	[34]
Pseudomonas aeruginosa	G-ive Bacteria	Au	NA	15–30	37	[33]
Shewanella Oneidensis	Facultative Bacteria	Au	Round	12	30	[137]
	Fungi	and Algae Species				
Plectonema boryanum UTEX 485	Filamentous Fungi	Au	Octahedral	10 nm–6 μm	25	[138]
Phaenerochaete chrysosporium	Fungi	Ag	Ag Pyramidal 50–200		37	[139]
Aspergillus flavus	Fungi	Ag	Round	8.92	25	[140]
Yeast	Fungi	Au, Ag	Polygonal	9–25	30	[131]
Fusarium oxysporum	Ascomycete fungus	Alloy of Au-Ag	Round	8-14	25	[117]
Sargassum wightii	Macro-algae	Au	Planar	8-12	NA	[127]
Neurospora crassa	Bread mold	Au–Ag, Au	Round	20–50	28	[50]
Verticillium sp.	Fungi	Ag	Round	25–32	25	[141]
Aspergillus fumigatus	Fungi	Ag Ag	Round NA	5–25 2–4	25 10–40	[124] [142]
Trichoderma viride	Fungi					
Yarrowia lipolytica	Fungi	Au	Triangles	15	30	[143]
Source of Intracellular Enzyme (Bacterial Species)	Nature of Microb.	Metals used	Shape	Size (nm)	Temperature (°C)	Referen
Shewanella algae	G-ive marine bacteria	Pt	NA	5	25	[144]
Enterobacter species	Anaerobic G-ive Bacilli	Hg	Round	2–5	30	[145]
Bacillus cereus	G+ive Bacteria	Ag	Round	4–5	37	[146]
Brevibacterium casei	Actinomycetales Bacteria	Ag, Au	Roud	10-50	37	[147]
Rhodococcus sp.	Actinobacteria	Au	Round	8-12	NA	[148]
	Fungi	and Algae Species				
Plectonema boryanum	Algae	Au	Cubic	<10-25	25-100	[128]
Neurospora crassa	Bread mold	Au–Ag, Au	Round	32	28	[50]
Verticillum luteoalbum	Ascomycota Fungi	Au	NA	NA	37	[149]
Candida utilis	Fungus	Au	NA	NA	25	[149]

Table 2. List of extracellular and intracellular bioreducing microbial enzymes and resulting nanoparticles.

NA: Not available; G+ive: Gram-positive; G-ive: Gram-negative.

In the intracellular mechanism of metal bioreduction, bacterial and fungal cells along with sugars molecules play a significant role. Mainly, the interactions of intracellular enzymes and positively charged groups are utilized in the gripping of metallic ions from the medium and the subsequent reduction inside the cell [150,151]. When observed microscopically, MtNPs accumulated in the

periplasmic space, the cytoplasmic membrane and the cell wall. This was due to the diffusion of metal

ions across the membranes and enzymatic reduction resulting in MtNPs. Among the actinomycetes, alkalo-tolerant (*Rhodococcus* sp.) and alkalo-thermophilic (Thermomonospora sp.) actinomycetes were used for the intracellular synthesis of AuNPs [148,152]. The intracellular production of AuNPs with uniform dimensions was carried out by reacting the *Rhodococcus* species with an aqueous solution of AuCl^{4–} ions. Reductions of Au³⁺ were effectively mediated by enzymes at the surface of the mycelia and the cytoplasmic membrane. The exposure of a Verticillium biomass to a Ag⁺ ionic solution resulted in intracellular reduction and the subsequent formation of AgNPs. Visualization via electron microscopy revealed that the AgNPs were formed under the surface of cell wall as a result of enzymatic bioreduction, which is non-toxic to the fungi [153]. The same procedure was adopted for the synthesis of AuNPs using the fungus *Verticillium* as the source of reducing enzymes. AuNPs were entrapped in the cytoplasmic membrane and the cell wall of the fungi, indicating that Au^{3+} was bio-reduced by reductase enzymes that were present there [154]. Southam and Beveridge reported that AuNPs were formed and precipitated inside bacterial cells after the incubation of the bacterial cells in a Au³⁺ ionic solution [155]. Pseudomonas stutzeri (AG259) when exposed to concentrated AgNO₃ solution has reduced Ag¹⁺ ions, with the subsequent formation of AgNPs in the bacterial periplasmic space [31]. A filamentous cyanobacterium (Plectonema boryanum) treated with AuCl⁴⁻ and Au(S₂O₃)₂³⁻ solutions resulted in the formation of AuNPs at the membrane level and gold sulfide residing intracellularly [128]. In another study, Phanerochaete chrysosporium incubation in an ionic Au³⁺ solution resulted in the formation of AuNPs 10–100 nm in particle size. The Laccase enzyme was used as an extracellular reducing agent, whereas ligninase was found to be responsible for the intracellular reduction of Au^{3+} ions [156]. Other factors, including the incubation age of the fungi, the concentration of the AuCl⁴⁻ solution and the incubation temperature demonstrated significant effects on the shape of AuNPs. The mesophilic bacterium Shewanella algae proved to be an efficient bioreducer of AuCl⁴⁻ ions to elemental gold. Au nanoparticles were found in the bacterial periplasmic space, mediated via intracellular enzymes [157]. Brevibacterium casei treated with aqueous solutions of Au³⁺ and Ag⁺ ions were reduced to intracellular enzymes, mediated by spherically shaped AuNPs and AgNPs respectively [147].

9. Challenges and Limitation of MtNP Synthesis by Microorganisms: A Possible Solution

Nanotechnology is an interdisciplinary field which involves close coordination between engineers, chemists and biologists to design and develop potential candidates for therapeutic applications. Research on nanoparticle biosynthesis using microbes has gained significant interest, owing to its simplicity, low cost of production and eco-friendliness. MtNPs have emerged as excellent agents for biomedical applications due to their unique physicochemical, optical, electronic and biological properties [158]. Expanding interest on the application of MtNPs in biology and healthcare has raised a need for the development of cheap, easy and eco-friendly methods for MtNP preparation, such as the bioinspired approaches [159,160]. The biosynthesis of MtNPs using microbes is well investigated by researchers globally, where the reduction of metal ions to MtNPs occurs as part of the heavy metal-resistance mechanism exhibited by microorganisms [161]. The primary challenge involved in microbial inspired synthesis is the selection of the best candidates, based on the intrinsic properties of microbes, such as growth rate, replication, biochemical activity etc. Controlling the shape, size and monodispersity of the biosynthesized MtNPs is another important criterion for achieving the desired therapeutic effects, which might be modulated by varying the concentration of biomolecules, the reduction time, the temperature and other factors. Another important challenge involved in

biosynthesis is the identification of the key components among the large pool of biomolecules obtained from microbial resources responsible for the reduction and stabilization of MtNPs from precursor salts. Maintaining the optimal conditions needed for the microbial growth and enzymatic activity is critical as these act as biocatalysts, playing a major role in the preparation of MtNPs. Optimizing the conditions, such as providing essential nutrients for growth, the size of inoculum, the temperature, the pH and the amount of light greatly enhance the efficiency of MtNP biosynthesis using microbes [162]. Yield of the final product and rate of biosynthesis of MtNPs using microorganisms are another set of important parameters to be considered to translate the measure of production to a large scale industrial level. In general, the large scale production of biosynthesized nanoparticles using microbes has several challenges in scaling up, due to medium to low yield compared to chemically synthesized nanoparticles. In general, the yield of the biosynthesis methods are 1/3rd of the yield obtained from the chemical synthesis methods. Also, we have to keep in mind that the laboratory scale bottom-up or top-down approaches for the synthesis of MtNPs varies significantly from industrial manufacturing. However, due to large availability, easy access and the high growth rate of microbes, the overall cost of the synthesis will be reduced at a large extent. Moreover, the cost of the biosynthesis gets further reduced as there is no requirement of organic solvents, chemical stabilizing agents, thermal heat or any fancy technique during these synthesis methods. Hence, it can be safely mentioned that, if the right techniques are properly utilized, the overall production cost of the NPs can be reduced by almost 1/10th compared to the chemical synthesis methods in the long run. Nevertheless, we strongly feel that during the initial stages of the development of NPs at the lab scale, researchers must consider and deploy appropriate methods useful for large-scale industrial manufacturing [163]. An advantage of this method is that the MtNPs produced using the biosynthesis route are more stable without any agglomeration, even at room temperatures for long periods of time when compared with MtNPs produced using chemical synthetic methods [164,165]. This stability might be due to the in situ capping of the microbial proteins or biomolecules over the MtNP surface during the biosynthesis process. Understanding the source and mechanism of biosynthesis is essential, as sometimes the utility of toxic microorganisms as a bioresource might impart pathogenicity to the prepared MtNPs. Above all, the protocols designed at the laboratory scale might yield well-characterized highly stable MtNPs. Scaling up this process to the industrial level is mandatory to fulfill the unmet medical challenges, which mainly depend upon optimizing the growth rate of the microbes, inoculation of the seed into the biomass, harvesting cells and effective MtNP preparation through reduction, as well as recovery of the final product with good yield [12,166,167]. Taken together, there are several important challenges and technical aspects that must be addressed to successfully translate this bioinspired method of MtNP synthesis for large scale industrial production.

10. Conclusions and Future Prospects

For the time being, it can be safely concluded that there is a bright future for the microbial synthesis of metal nanoparticles and their wide variety of applications in biomedical nanotechnology when compared with chemically assisted NPs, due to their low cost and eco-friendliness. The huge pool of bioresources (microbes and microbial enzymes), if appropriately exploited, could help biosynthesized NPs to turn out to be a potential game changer in the imminent future. However, several challenges must be dealt with before industrial scale production and wide use, which will take a little more time (5–10 years). Nevertheless, there is immense potential for microbial assisted metal nanoparticles as they are low in toxicity and cost and high in degradability and are useful in numerous therapeutic applications. It is also encouraging that several groups are concentrating on decoding the detailed mechanistic aspects of microbial biosynthesis, which will ultimately lead to better understanding and wiser applicability. Even though the mechanisms behind these nanoparticles' uptake, diffusion, long term toxicity and excretion remain uncertain, the potential biological applications make them a potential candidate to extensively substitute chemically synthesized NPs in the coming years.

Author Contributions: Conceptualization, M.O. and S.M.; writing-original draft preparation, A.T.K., M.A., I.A., S.K.N. and S.M.; writing-review and editing, M.O., S.M.; funding acquisition, S.M.

Funding: This research received no external funding. The APC was funded by Fondren Library, Rice University, Houston, TX, United States.

Acknowledgments: Sudip Mukherjee acknowledges Fondren Library, Rice University, Houston, TX, United States for financial support for the APC of this work. He also acknowledges Omid Veiseh (supervisor/mentor) for providing continuous support for his postdoctoral research.

Conflicts of Interest: There is no conflict of interest. The authors have no other pertinent affiliations or financial connection with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

References

- 1. Slavin, Y.N.; Asnis, J.; Häfeli, U.O.; Bach, H. Metal nanoparticles: Understanding the mechanisms behind antibacterial activity. *J. Nanobiotechnol.* **2017**, *15*, 65. [CrossRef] [PubMed]
- 2. Khan, I.; Saeed, K.; Khan, I. Nanoparticles: Properties, applications and toxicities. Arab. J. Chem. 2017. [CrossRef]
- 3. FutureMarketInsights Global Market for Metal & Metal Oxide Nanoparticles to Surge at More Than 10% CAGR. Available online: http://markets.businessinsider.com/news/stocks/Global-Market-forMetal-Metal-Oxide-Nanoparticles-to-Surge-at-More-Than-10CAGR-1001862836 (accessed on 21 November 2018).
- 4. Mukherjee, S.; Vinothkumar, B.; Prashanthi, S.; Bangal, P.R.; Sreedhar, B.; Patra, C.R. Potential therapeutic and diagnostic applications of one-step in situ biosynthesized gold nanoconjugates (2-in-1 system) in cancer treatment. *RSC Adv.* **2013**, *3*, 2318–2329. [CrossRef]
- Patra, S.; Mukherjee, S.; Barui, A.K.; Ganguly, A.; Sreedhar, B.; Patra, C.R. Green synthesis, characterization of gold and silver nanoparticles and their potential application for cancer therapeutics. *Mater. Sci. Eng. C* 2015, *53*, 298–309. [CrossRef] [PubMed]
- 6. Ovais, M.; Zia, N.; Ahmad, I.; Khalil, A.T.; Raza, A.; Ayaz, M.; Sadiq, A.; Ullah, F.; Shinwari, Z.K. Phyto-Therapeutic and Nanomedicinal Approaches to Cure Alzheimer's Disease: Present Status and Future Opportunities. *Front. Aging Neurosci.* **2018**, *10*. [CrossRef] [PubMed]
- Mukherjee, S.; Dasari, M.; Priyamvada, S.; Kotcherlakota, R.; Bollu, V.S.; Patra, C.R. A green chemistry approach for the synthesis of gold nanoconjugates that induce the inhibition of cancer cell proliferation through induction of oxidative stress and their in vivo toxicity study. *J. Mater. Chem. B* 2015, *3*, 3820–3830. [CrossRef]
- 8. Ovais, M.; Khalil, A.T.; Raza, A.; Khan, M.A.; Ahmad, I.; Islam, N.U.; Saravanan, M.; Ubaid, M.F.; Ali, M.; Shinwari, Z.K. Green synthesis of silver nanoparticles via plant extracts: beginning a new era in cancer theranostics. *Nanomedicine* **2016**, *12*, 3157–3177. [CrossRef] [PubMed]
- Salunke, G.R.; Ghosh, S.; Kumar, R.S.; Khade, S.; Vashisth, P.; Kale, T.; Chopade, S.; Pruthi, V.; Kundu, G.; Bellare, J.R. Rapid efficient synthesis and characterization of silver, gold, and bimetallic nanoparticles from the medicinal plant Plumbago zeylanica and their application in biofilm control. *Int. J. Nanomed.* 2014, 9, 2635.
- 10. Barabadi, H.; Ovais, M.; Shinwari, Z.K.; Saravanan, M. Anti-cancer green bionanomaterials: present status and future prospects. *Green Chem. Lett. Rev.* **2017**, *10*, 285–314. [CrossRef]
- 11. Emmanuel, R.; Saravanan, M.; Ovais, M.; Padmavathy, S.; Shinwari, Z.K.; Prakash, P. Antimicrobial efficacy of drug blended biosynthesized colloidal gold nanoparticles from Justicia glauca against oral pathogens: a nanoantibiotic approach. *Microb. Pathog.* **2017**, *113*, 295–302. [CrossRef] [PubMed]
- 12. Hulkoti, N.I.; Taranath, T. Biosynthesis of nanoparticles using microbes—A review. *Colloids Surf. B* 2014, 121, 474–483. [CrossRef] [PubMed]
- Tang, L.; Shi, J.; Wu, H.; Zhang, S.; Liu, H.; Zou, H.; Wu, Y.; Zhao, J.; Jiang, Z. In situ biosynthesis of ultrafine metal nanoparticles within a metal-organic framework for efficient heterogeneous catalysis. *Nanotechnology* 2017, 28, 365604. [CrossRef] [PubMed]
- 14. Ovais, M.; Raza, A.; Naz, S.; Islam, N.U.; Khalil, A.T.; Ali, S.; Khan, M.A.; Shinwari, Z.K. Current state and prospects of the phytosynthesized colloidal gold nanoparticles and their applications in cancer theranostics. *Appl. Microbiol. Biotechnol.* **2017**, *101*, 3551–3565. [CrossRef] [PubMed]

- Ovais, M.; Khalil, A.T.; Islam, N.U.; Ahmad, I.; Ayaz, M.; Saravanan, M.; Shinwari, Z.K.; Mukherjee, S. Role of plant phytochemicals and microbial enzymes in biosynthesis of metallic nanoparticles. *Appl. Microbiol. Biotechnol.* 2018, 102, 6799–6814. [CrossRef] [PubMed]
- 16. Singh, P.; Kim, Y.-J.; Zhang, D.; Yang, D.-C. Biological synthesis of nanoparticles from plants and microorganisms. *Trends Biotechnol.* **2016**, *34*, 588–599. [CrossRef] [PubMed]
- 17. Baker, S.; Rakshith, D.; Kavitha, K.S.; Santosh, P.; Kavitha, H.U.; Rao, Y.; Satish, S. Plants: Emerging as nanofactories towards facile route in synthesis of nanoparticles. *BioImpacts* **2013**, *3*, 111. [PubMed]
- 18. Kumari, R.; Barsainya, M.; Singh, D.P. Biogenic synthesis of silver nanoparticle by using secondary metabolites from Pseudomonas aeruginosa DM1 and its anti-algal effect on Chlorella vulgaris and Chlorella pyrenoidosa. *Environ. Sci. Pollut. Res.* **2017**, *24*, 4645–4654. [CrossRef] [PubMed]
- 19. Patra, C.R.; Mukherjee, S.; Kotcherlakota, R. Biosynthesized silver nanoparticles: A step forward for cancer theranostics? *Nanomedicine* **2014**, *9*, 1445–1448. [CrossRef] [PubMed]
- 20. Mukherjee, S.; Nethi, S.K.; Patra, C.R. Green synthesized gold nanoparticles for future biomedical applications. In *Particulate Technology for Delivery of Therapeutics*; Springer: Singapore, 2017; pp. 359–393.
- Khalil, A.T.; Ovais, M.; Ullah, I.; Ali, M.; Shinwari, Z.K.; Khamlich, S.; Maaza, M. Sageretia thea (Osbeck.) mediated synthesis of zinc oxide nanoparticles and its biological applications. *Nanomedicine* 2017, 12, 1767–1789. [CrossRef] [PubMed]
- Khalil, A.T.; Ovais, M.; Ullah, I.; Ali, M.; Shinwari, Z.K.; Hassan, D.; Maaza, M. Sageretia thea (Osbeck.) modulated biosynthesis of NiO nanoparticles and their in vitro pharmacognostic, antioxidant and cytotoxic potential. *Artif. Cells Nanomed. Biotechnol.* 2018, 46, 838–852. [CrossRef] [PubMed]
- 23. Makarov, V.; Love, A.; Sinitsyna, O.; Makarova, S.; Yaminsky, I.; Taliansky, M.; Kalinina, N. "Green" nanotechnologies: Synthesis of metal nanoparticles using plants. *Acta Nat.* **2014**, *6*, 35–44.
- 24. Sintubin, L.; Verstraete, W.; Boon, N. Biologically produced nanosilver: Current state and future perspectives. *Biotechnol. Bioeng.* **2012**, *109*, 2422–2436. [CrossRef] [PubMed]
- 25. Mukherjee, S.; Sushma, V.; Patra, S.; Barui, A.K.; Bhadra, M.P.; Sreedhar, B.; Patra, C.R. Green chemistry approach for the synthesis and stabilization of biocompatible gold nanoparticles and their potential applications in cancer therapy. *Nanotechnology* **2012**, *23*, 455103. [CrossRef] [PubMed]
- Mukherjee, S.; Sau, S.; Madhuri, D.; Bollu, V.S.; Madhusudana, K.; Sreedhar, B.; Banerjee, R.; Patra, C.R. Green synthesis and characterization of monodispersed gold nanoparticles: Toxicity study, delivery of doxorubicin and its bio-distribution in mouse model. *J. Biomed. Nanotechnol.* 2016, *12*, 165–181. [CrossRef] [PubMed]
- 27. Mukherjee, S.; Chowdhury, D.; Kotcherlakota, R.; Patra, S. Potential theranostics application of bio-synthesized silver nanoparticles (4-in-1 system). *Theranostics* **2014**, *4*, 316–335. [CrossRef] [PubMed]
- 28. Singh, R.; Wagh, P.; Wadhwani, S.; Gaidhani, S.; Kumbhar, A.; Bellare, J.; Chopade, B.A. Synthesis, optimization, and characterization of silver nanoparticles from Acinetobacter calcoaceticus and their enhanced antibacterial activity when combined with antibiotics. *Int. J. Nanomed.* **2013**, *8*, 4277.
- 29. Singh, R.; Shedbalkar, U.U.; Wadhwani, S.A.; Chopade, B.A. Bacteriagenic silver nanoparticles: Synthesis, mechanism, and applications. *Appl. Microbiol. Biotechnol.* **2015**, *99*, 4579–4593. [CrossRef] [PubMed]
- Saravanan, M.; Barik, S.K.; Mubarak Ali, D.; Prakash, P.; Pugazhendhi, A. Synthesis of silver nanoparticles from Bacillus brevis (NCIM 2533) and their antibacterial activity against pathogenic bacteria. *Microb. Pathog.* 2018, 116, 221–226. [CrossRef] [PubMed]
- 31. Klaus, T.; Joerger, R.; Olsson, E.; Granqvist, C.-G. Silver-based crystalline nanoparticles, microbially fabricated. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 13611–13614. [CrossRef] [PubMed]
- 32. Pugazhenthiran, N.; Anandan, S.; Kathiravan, G.; Prakash, N.K.U.; Crawford, S.; Ashokkumar, M. Microbial synthesis of silver nanoparticles by Bacillus sp. *J. Nanopart. Res.* **2009**, *11*, 1811. [CrossRef]
- 33. Husseiny, M.; El-Aziz, M.A.; Badr, Y.; Mahmoud, M. Biosynthesis of gold nanoparticles using Pseudomonas aeruginosa. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* **2007**, *67*, 1003–1006. [CrossRef] [PubMed]
- 34. He, S.; Guo, Z.; Zhang, Y.; Zhang, S.; Wang, J.; Gu, N. Biosynthesis of gold nanoparticles using the bacteria Rhodopseudomonas capsulata. *Mater. Lett.* **2007**, *61*, 3984–3987. [CrossRef]
- 35. Dhandapani, P.; Siddarth, A.S.; Kamalasekaran, S.; Maruthamuthu, S.; Rajagopal, G. Bio-approach: Ureolytic bacteria mediated synthesis of ZnO nanocrystals on cotton fabric and evaluation of their antibacterial properties. *Carbohyd. Polym.* **2014**, *103*, 448–455. [CrossRef] [PubMed]

- 36. Selvarajan, E.; Mohanasrinivasan, V. Biosynthesis and characterization of ZnO nanoparticles using Lactobacillus plantarum VITES07. *Mater. Lett.* **2013**, *112*, 180–182. [CrossRef]
- Jayaseelan, C.; Rahuman, A.A.; Kirthi, A.V.; Marimuthu, S.; Santhoshkumar, T.; Bagavan, A.; Gaurav, K.; Karthik, L.; Rao, K.B. Novel microbial route to synthesize ZnO nanoparticles using Aeromonas hydrophila and their activity against pathogenic bacteria and fungi. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 2012, 90, 78–84. [CrossRef] [PubMed]
- Rad, M.; Taran, M.; Alavi, M. Effect of Incubation Time, CuSO₄ and Glucose Concentrations on Biosynthesis of Copper Oxide (CuO) Nanoparticles with Rectangular Shape and Antibacterial Activity: Taguchi Method Approach. *Nano Biomed. Eng.* 2018, 10, 25–33. [CrossRef]
- 39. Fatemi, M.; Mollania, N.; Momeni-Moghaddam, M.; Sadeghifar, F. Extracellular biosynthesis of magnetic iron oxide nanoparticles by Bacillus cereus strain HMH1: Characterization and in vitro cytotoxicity analysis on MCF-7 and 3T3 cell lines. *J. Biotechnol.* **2018**, *270*, 1–11. [CrossRef] [PubMed]
- 40. Nair, B.; Pradeep, T. Coalescence of nanoclusters and formation of submicron crystallites assisted by Lactobacillus strains. *Cryst. Growth Des.* **2002**, *2*, 293–298. [CrossRef]
- 41. Lengke, M.F.; Fleet, M.E.; Southam, G. Biosynthesis of silver nanoparticles by filamentous cyanobacteria from a silver (I) nitrate complex. *Langmuir* **2007**, *23*, 2694–2699. [CrossRef] [PubMed]
- 42. Patel, V.; Berthold, D.; Puranik, P.; Gantar, M. Screening of cyanobacteria and microalgae for their ability to synthesize silver nanoparticles with antibacterial activity. *Biotechnol. Rep.* **2015**, *5*, 112–119. [CrossRef] [PubMed]
- 43. Tomer, A.K.; Rahi, T.; Neelam, D.K.; Dadheech, P.K. Cyanobacterial extract-mediated synthesis of silver nanoparticles and their application in ammonia sensing. *Int. Microbiol.* **2018**. [CrossRef]
- 44. Satapathy, S.; Shukla, S.P. Application of a marine cyanobacterium Phormidium fragile for green synthesis of silver nanoparticles. *Indian J. Biotechnol.* **2017**, *16*, 110–113.
- 45. Sonker, A.S.; Pathak, J.; Kannaujiya, V.; Sinha, R. Characterization and in vitro antitumor, antibacterial and antifungal activities of green synthesized silver nanoparticles using cell extract of Nostoc sp. strain HKAR-2. *Can. J. Biotechnol.* **2017**, *1*, 26–37. [CrossRef]
- Bakir, E.; Younis, N.; Mohamed, M.; El Semary, N. Cyanobacteria as Nanogold Factories: Chemical and Anti-Myocardial Infarction Properties of Gold Nanoparticles Synthesized by Lyngbya majuscula. *Mar. Drugs* 2018, 16, 217. [CrossRef] [PubMed]
- 47. Molnár, Z.; Bódai, V.; Szakacs, G.; Erdélyi, B.; Fogarassy, Z.; Sáfrán, G.; Varga, T.; Kónya, Z.; Tóth-Szeles, E.; Szűcs, R. Green synthesis of gold nanoparticles by thermophilic filamentous fungi. *Sci. Rep.* **2018**, *8*, 3943. [CrossRef] [PubMed]
- Zhao, X.; Zhou, L.; Riaz Rajoka, M.S.; Yan, L.; Jiang, C.; Shao, D.; Zhu, J.; Shi, J.; Huang, Q.; Yang, H. Fungal silver nanoparticles: Synthesis, application and challenges. *Crit. Rev. Biotechnol.* 2018, *38*, 817–835. [CrossRef] [PubMed]
- Alghuthaymi, M.A.; Almoammar, H.; Rai, M.; Said-Galiev, E.; Abd-Elsalam, K.A. Myconanoparticles: Synthesis and their role in phytopathogens management. *Biotechnol. Biotechnol. Eq.* 2015, 29, 221–236. [CrossRef] [PubMed]
- 50. Castro-Longoria, E.; Vilchis-Nestor, A.R.; Avalos-Borja, M. Biosynthesis of silver, gold and bimetallic nanoparticles using the filamentous fungus Neurospora crassa. *Colloids Surf. B* **2011**, *83*, 42–48. [CrossRef] [PubMed]
- 51. Zhang, X.; He, X.; Wang, K.; Yang, X. Different active biomolecules involved in biosynthesis of gold nanoparticles by three fungus species. *J. Biomed. Nanotechnol.* **2011**, *7*, 245–254. [CrossRef] [PubMed]
- AbdelRahim, K.; Mahmoud, S.Y.; Ali, A.M.; Almaary, K.S.; Mustafa, A.E.-Z.M.; Husseiny, S.M. Extracellular biosynthesis of silver nanoparticles using Rhizopus stolonifer. *Saudi J. Biol. Sci.* 2017, 24, 208–216. [CrossRef] [PubMed]
- Jalal, M.; Ansari, M.; Alzohairy, M.; Ali, S.; Khan, H.; Almatroudi, A.; Raees, K. Biosynthesis of Silver Nanoparticles from Oropharyngeal Candida glabrata Isolates and Their Antimicrobial Activity against Clinical Strains of Bacteria and Fungi. *Nanomaterials* 2018, *8*, 586. [CrossRef] [PubMed]
- 54. Kalpana, V.; Kataru, B.A.S.; Sravani, N.; Vigneshwari, T.; Panneerselvam, A.; Rajeswari, V.D. Biosynthesis of Zinc oxide nanoparticles using culture filtrates of Aspergillus niger: Antimicrobial textiles and dye degradation studies. *OpenNano* **2018**. [CrossRef]

- 55. Vijayanandan, A.S.; Balakrishnan, R.M. Biosynthesis of cobalt oxide nanoparticles using endophytic fungus Aspergillus nidulans. *J. Environ. Manag.* **2018**, *218*, 442–450. [CrossRef] [PubMed]
- Sanaeimehr, Z.; Javadi, I.; Namvar, F. Antiangiogenic and antiapoptotic effects of green-synthesized zinc oxide nanoparticles using Sargassum muticum algae extraction. *Cancer Nanotechnol.* 2018, *9*, 3. [CrossRef] [PubMed]
- 57. Maceda, A.F.; Ouano, J.J.S.; Que, M.C.O.; Basilia, B.A.; Potestas, M.J.; Alguno, A.C. Controlling the Absorption of Gold Nanoparticles via Green Synthesis Using Sargassum crassifolium Extract. In *Key Engineering Materials*; Trans Tech Publ: Clausthal-Zellerfeld, Germany, 2018; pp. 44–48.
- 58. Gu, H.; Chen, X.; Chen, F.; Zhou, X.; Parsaee, Z. Ultrasound-assisted biosynthesis of CuO-NPs using brown alga Cystoseira trinodis: Characterization, photocatalytic AOP, DPPH scavenging and antibacterial investigations. *Ultrason. Sonochem.* **2018**, *41*, 109–119. [CrossRef] [PubMed]
- 59. Koopi, H.; Buazar, F. A novel one-pot biosynthesis of pure alpha aluminum oxide nanoparticles using the macroalgae Sargassum ilicifolium: A green marine approach. *Ceram. Int.* **2018**, *44*, 8940–8945. [CrossRef]
- 60. Vijayaraghavan, K.; Mahadevan, A.; Sathishkumar, M.; Pavagadhi, S.; Balasubramanian, R. Biosynthesis of Au (0) from Au (III) via biosorption and bioreduction using brown marine alga Turbinaria conoides. *Chem. Eng. J.* **2011**, *167*, 223–227. [CrossRef]
- 61. Ramakrishna, M.; Babu, D.R.; Gengan, R.M.; Chandra, S.; Rao, G.N. Green synthesis of gold nanoparticles using marine algae and evaluation of their catalytic activity. *J. Nanostruct. Chem.* **2016**, *6*, 1–13. [CrossRef]
- 62. Swaminathan, S.; Murugesan, S.; Damodarkumar, S.; Dhamotharan, R.; Bhuvaneshwari, S. Synthesis and characterization of gold nanoparticles from alga Acanthophora spicifera (VAHL) Boergesen. *Int. J. Nanosci. Nanotechnol.* **2011**, *2*, 85–94.
- 63. Ghodake, G.; Lee, D.S. Biological synthesis of gold nanoparticles using the aqueous extract of the brown algae Laminaria japonica. *J. Nanoelectron. Optoelectron.* **2011**, *6*, 268–271. [CrossRef]
- 64. Govindaraju, K.; Basha, S.K.; Kumar, V.G.; Singaravelu, G. Silver, gold and bimetallic nanoparticles production using single-cell protein (Spirulina platensis) Geitler. J. Mater. Sci. 2008, 43, 5115–5122. [CrossRef]
- 65. Wypij, M.; Golinska, P.; Dahm, H.; Rai, M. Actinobacterial-mediated synthesis of silver nanoparticles and their activity against pathogenic bacteria. *IET Nanobiotechnol.* **2016**, *11*, 336–342. [CrossRef] [PubMed]
- Wadhwani, S.A.; Shedbalkar, U.U.; Singh, R.; Vashisth, P.; Pruthi, V.; Chopade, B.A. Kinetics of synthesis of gold nanoparticles by Acinetobacter sp. SW30 isolated from environment. *Indian J. Microbiol.* 2016, 56, 439–444. [CrossRef] [PubMed]
- 67. Prema, P.; Iniya, P.; Immanuel, G. Microbial mediated synthesis, characterization, antibacterial and synergistic effect of gold nanoparticles using Klebsiella pneumoniae (MTCC-4030). *RSC Adv.* **2016**, *6*, 4601–4607. [CrossRef]
- Manikprabhu, D.; Cheng, J.; Chen, W.; Sunkara, A.K.; Mane, S.B.; Kumar, R.; Hozzein, W.N.; Duan, Y.-Q.; Li, W.-J. Sunlight mediated synthesis of silver nanoparticles by a novel actinobacterium (Sinomonas mesophila MPKL 26) and its antimicrobial activity against multi drug resistant Staphylococcus aureus. J. Photochem. Photobiol. B 2016, 158, 202–205. [CrossRef] [PubMed]
- 69. Syed, B.; Prasad, N.M.; Satish, S. Endogenic mediated synthesis of gold nanoparticles bearing bactericidal activity. *J. Microsc. Ultrastruct.* **2016**, *4*, 162–166. [PubMed]
- 70. Gan, L.; Zhang, S.; Zhang, Y.; He, S.; Tian, Y. Biosynthesis, characterization and antimicrobial activity of silver nanoparticles by a halotolerant Bacillus endophyticus SCU-L. *Prep. Biochem. Biotechnol.* 2018. [CrossRef] [PubMed]
- 71. Vijayabharathi, R.; Sathya, A.; Gopalakrishnan, S. Extracellular biosynthesis of silver nanoparticles using Streptomyces griseoplanus SAI-25 and its antifungal activity against Macrophomina phaseolina, the charcoal rot pathogen of sorghum. *Biocatal. Agric. Biotechnol.* **2018**, *14*, 166–171. [CrossRef]
- 72. Ranjani, A.; Gopinath, P.M.; Ananth, S.; Narchonai, G.; Santhanam, P.; Thajuddin, N.; Dhanasekaran, D. Multidimensional dose–response toxicity exploration of silver nanoparticles from Nocardiopsis flavascens RD30. *Appl. Nanosci.* **2018**, *8*, 699–713. [CrossRef]
- 73. Bing, W.; Sun, H.; Wang, F.; Song, Y.; Ren, J. Hydrogen-producing Hyperthermophilic Bacteria Synthesized Size-controllable Fine Gold Nanoparticles with Excellence for Eradicating Biofilm and Antibacterial Applications. *J. Mater. Chem. B* 2018. [CrossRef]

- Lv, Q.; Zhang, B.; Xing, X.; Zhao, Y.; Cai, R.; Wang, W.; Gu, Q. Biosynthesis of copper nanoparticles using Shewanella loihica PV-4 with antibacterial activity: Novel approach and mechanisms investigation. *J. Hazard. Mater.* 2018, 347, 141–149. [CrossRef] [PubMed]
- 75. Ahmed, E.; Kalathil, S.; Shi, L.; Alharbi, O.; Wang, P. Synthesis of ultra-small platinum, palladium and gold nanoparticles by Shewanella loihica PV-4 electrochemically active biofilms and their enhanced catalytic activities. *J. Saudi Chem. Soc.* **2018**. [CrossRef]
- 76. Jafari, M.; Rokhbakhsh-Zamin, F.; Shakibaie, M.; Moshafi, M.H.; Ameri, A.; Rahimi, H.R.; Forootanfar, H. Cytotoxic and antibacterial activities of biologically synthesized gold nanoparticles assisted by Micrococcus yunnanensis strain J2. *Biocatal. Agric. Biotechnol.* 2018. [CrossRef]
- 77. Camas, M.; Camas, A.S.; Kyeremeh, K. Extracellular Synthesis and Characterization of Gold Nanoparticles Using Mycobacterium sp. BRS2A-AR2 Isolated from the Aerial Roots of the Ghanaian Mangrove Plant, Rhizophora racemosa. *Indian J. Microbiol.* 2018, *58*, 214–221. [CrossRef] [PubMed]
- 78. Shah, R.; Oza, G.; Pandey, S.; Sharon, M. Biogenic fabrication of gold nanoparticles using Halomonas salina. *J. Microbiol. Biotechnol.* **2017**, *2*, 485–492.
- 79. Kobashigawa, J.M.; Robles, C.A.; Ricci, M.L.M.; Carmarán, C.C. Influence of strong bases on the synthesis of silver nanoparticles (AgNPs) using the ligninolytic fungi Trametes trogii. *Saudi J.Biol. Sci.* **2018**. [CrossRef]
- Elamawi, R.M.; Al-Harbi, R.E.; Hendi, A.A. Biosynthesis and characterization of silver nanoparticles using Trichoderma longibrachiatum and their effect on phytopathogenic fungi. *Egypt. J. Biol. Pest Control* 2018, 28, 28. [CrossRef]
- Tripathi, R.M.; Shrivastav, B.R.; Shrivastav, A. Antibacterial and catalytic activity of biogenic gold nanoparticles synthesised byTrichoderma harzianum. *IET Nanobiotechnol.* 2018, 12, 509–513. [CrossRef] [PubMed]
- 82. Ahmed, A.-A.; Hamzah, H.; Maaroof, M. Analyzing formation of silver nanoparticles from the filamentous fungus Fusarium oxysporum and their antimicrobial activity. *Turk. J. Biol.* **2018**, *42*, 54–62. [CrossRef]
- El Domany, E.B.; Essam, T.M.; Ahmed, A.E.; Farghali, A.A. Biosynthesis Physico-Chemical Optimization of Gold Nanoparticles as Anti-Cancer and Synergetic Antimicrobial Activity Using Pleurotus ostreatus Fungus. *J. Appl. Pharm. Sci. Vol.* 2018, *8*, 119–128.
- 84. Singh, P.S.; Vidyasagar, G. Biosynthesis of antibacterial silver nano-particles from Aspergillus terreus. *World News Nat. Sci.* **2018**, *16*, 117–124.
- Mohanta, Y.K.; Nayak, D.; Biswas, K.; Singdevsachan, S.K.; Abd_Allah, E.F.; Hashem, A.; Alqarawi, A.A.; Yadav, D.; Mohanta, T.K. Silver Nanoparticles Synthesized Using Wild Mushroom Show Potential Antimicrobial Activities against Food Borne Pathogens. *Molecules* 2018, 23, 655. [CrossRef] [PubMed]
- 86. Saravanan, M.; Arokiyaraj, S.; Lakshmi, T.; Pugazhendhi, A. Synthesis of silver nanoparticles from Phenerochaete chrysosporium (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb. Pathog.* **2018**, *117*, 68–72. [CrossRef] [PubMed]
- Neethu, S.; Midhun, S.J.; Sunil, M.; Soumya, S.; Radhakrishnan, E.; Jyothis, M. Efficient visible light induced synthesis of silver nanoparticles by Penicillium polonicum ARA 10 isolated from Chetomorpha antennina and its antibacterial efficacy against Salmonella enterica serovar Typhimurium. *J. Photochem. Photobiol. B* 2018, 180, 175–185. [CrossRef] [PubMed]
- Spagnoletti, F.N.; Spedalieri, C.; Kronberg, F.; Giacometti, R. Extracellular biosynthesis of bactericidal Ag/AgCl nanoparticles for crop protection using the fungus Macrophomina phaseolina. *J. Environ. Manag.* 2019, 231, 457–466. [CrossRef] [PubMed]
- Cunha, F.A.; da CSO Cunha, M.; da Frota, S.M.; Mallmann, E.J.; Freire, T.M.; Costa, L.S.; Paula, A.J.; Menezes, E.A.; Fechine, P.B. Biogenic synthesis of multifunctional silver nanoparticles from Rhodotorula glutinis and Rhodotorula mucilaginosa: Antifungal, catalytic and cytotoxicity activities. *World J. Microbiol. Biotechnol.* 2018, 34, 127. [CrossRef] [PubMed]
- 90. Popli, D.; Anil, V.; Subramanyam, A.B.; Rao, S.N.; Rai, R.V.; Govindappa, M. Endophyte fungi, Cladosporium species-mediated synthesis of silver nanoparticles possessing in vitro antioxidant, anti-diabetic and anti-Alzheimer activity. *Artif. Cells Nanomed.* **2018**. [CrossRef] [PubMed]
- 91. Joshi, C.G.; Danagoudar, A.; Poyya, J.; Kudva, A.K.; Dhananjaya, B. Biogenic synthesis of gold nanoparticles by marine endophytic fungus-Cladosporium cladosporioides isolated from seaweed and evaluation of their antioxidant and antimicrobial properties. *Process Biochem.* **2017**, *63*, 137–144.

- 92. Farsi, M.; Farokhi, S. Biosynthesis of Antibacterial Silver Nanoparticles by Endophytic Fungus Nemania sp. Isolated From Taxus baccata L.(Iranian Yew). *Zahedan J. Res. Med. Sci.* **2018**, 20. [CrossRef]
- Subramaniyan, S.A.; Sheet, S.; Vinothkannan, M.; Yoo, D.J.; Lee, Y.S.; Belal, S.A.; Shim, K.S. One-Pot Facile Synthesis of Pt Nanoparticles Using Cultural Filtrate of Microgravity Simulated Grown P. chrysogenum and Their Activity on Bacteria and Cancer Cells. *J. Nanosci. Nanotechnol.* 2018, 18, 3110–3125. [CrossRef] [PubMed]
- Shen, W.; Qu, Y.; Pei, X.; Li, S.; You, S.; Wang, J.; Zhang, Z.; Zhou, J. Catalytic reduction of 4-nitrophenol using gold nanoparticles biosynthesized by cell-free extracts of Aspergillus sp. WL-Au. *J. Hazard. Mater.* 2017, 321, 299–306. [CrossRef] [PubMed]
- 95. Kumaresan, M.; Anand, K.V.; Govindaraju, K.; Tamilselvan, S.; Kumar, V.G. Seaweed Sargassum wightii mediated preparation of zirconia (ZrO₂) nanoparticles and their antibacterial activity against gram positive and gram negative bacteria. *Microb. Pathog.* **2018**, *124*, 311–315. [CrossRef] [PubMed]
- 96. Bao, Z.; Lan, C.Q. Mechanism of light-dependent biosynthesis of silver nanoparticles mediated by cell extract of Neochloris oleoabundans. *Colloids Surf. B* **2018**. [CrossRef] [PubMed]
- 97. González-Ballesteros, N.; Prado-López, S.; Rodriguez-Gonzalez, J.; Lastra, M.; Rodríguez-Argüelles, M. Green synthesis of gold nanoparticles using brown algae cystoseira baccata: Its activity in colon cancer cells. *Colloids Surf. B* 2017, 153, 190–198. [CrossRef] [PubMed]
- Pytlik, N.; Kaden, J.; Finger, M.; Naumann, J.; Wanke, S.; Machill, S.; Brunner, E. Biological synthesis of gold nanoparticles by the diatom Stephanopyxis turris and in vivo SERS analyses. *Algal Res.* 2017, 28, 9–15. [CrossRef]
- Abdel-Raouf, N.; Al-Enazi, N.M.; Ibraheem, I.B. Green biosynthesis of gold nanoparticles using Galaxaura elongata and characterization of their antibacterial activity. *Arab. J. Chem.* 2017, 10, S3029–S3039. [CrossRef]
- Arsiya, F.; Sayadi, M.H.; Sobhani, S. Green synthesis of palladium nanoparticles using Chlorella vulgaris. *Mater. Lett.* 2017, 186, 113–115. [CrossRef]
- 101. Ramkumar, V.S.; Pugazhendhi, A.; Gopalakrishnan, K.; Sivagurunathan, P.; Saratale, G.D.; Dung, T.N.B.; Kannapiran, E. Biofabrication and characterization of silver nanoparticles using aqueous extract of seaweed Enteromorpha compressa and its biomedical properties. *Biotechnol. Rep.* 2017, 14, 1–7. [CrossRef] [PubMed]
- 102. Vanlalveni, C.; Rajkumari, K.; Biswas, A.; Adhikari, P.P.; Lalfakzuala, R.; Rokhum, L. Green Synthesis of Silver Nanoparticles Using Nostoc linckia and its Antimicrobial Activity: A Novel Biological Approach. *Bionanoscience* 2018, *8*, 624–631. [CrossRef]
- 103. Zada, S.; Ahmad, A.; Khan, S.; Yu, X.; Chang, K.; Iqbal, A.; Ahmad, A.; Ullah, S.; Raza, M.; Khan, A. Biogenic synthesis of silver nanoparticles using extracts of Leptolyngbya JSC-1 that induce apoptosis in HeLa cell line and exterminate pathogenic bacteria. *Artif. Cells Nanomed. Biotechnol.* 2018. [CrossRef] [PubMed]
- 104. Murugesan, S.; Bhuvaneswari, S.; Sivamurugan, V. Green synthesis, characterization of silver nanoparticles of a marine red alga Spyridia fusiformis and their antibacterial activity. *Int. J. Pharm. Pharm. Sci.* 2017, 9, 192–197. [CrossRef]
- 105. Zhang, Z.; Chen, J.; Yang, Q.; Lan, K.; Yan, Z.; Chen, J. Eco-friendly intracellular microalgae synthesis of fluorescent CdSe QDs as a sensitive nanoprobe for determination of imatinib. *Sens. Actuators B Chem.* 2018, 263, 625–633. [CrossRef]
- 106. Abdel-Raouf, N.; Al-Enazi, N.M.; Ibraheem, I.B.M.; Alharbi, R.M.; Alkhulaifi, M.M. Biosynthesis of silver nanoparticles by using of the marine brown alga Padina pavonia and their characterization. *Saudi J. Biol. Sci.* 2018. [CrossRef]
- 107. Sayadi, M.H.; Salmani, N.; Heidari, A.; Rezaei, M.R. Bio-synthesis of palladium nanoparticle using Spirulina platensis alga extract and its application as adsorbent. *Surf. Interfaces* **2018**, *10*, 136–143. [CrossRef]
- 108. Sharma, M.; Behl, K.; Nigam, S.; Joshi, M. TiO₂-GO nanocomposite for photocatalysis and environmental applications: A green synthesis approach. *Vacuum* **2018**, *156*, 434–439. [CrossRef]
- Narayanan, K.B.; Sakthivel, N. Biological synthesis of metal nanoparticles by microbes. *Adv. Colloids Interface Sci.* 2010, 156. [CrossRef] [PubMed]
- 110. Subbaiya, R.; Saravanan, M.; Priya, A.R.; Shankar, K.R.; Selvam, M.; Ovais, M.; Balajee, R.; Barabadi, H. Biomimetic synthesis of silver nanoparticles from Streptomyces atrovirens and their potential anticancer activity against human breast cancer cells. *IET Nanobiotechnol.* 2017, *11*, 965–972. [CrossRef] [PubMed]
- 111. Bose, D.; Chatterjee, S. Biogenic synthesis of silver nanoparticles using guava (Psidium guajava) leaf extract and its antibacterial activity against Pseudomonas aeruginosa. *Appl. Nanosci.* **2016**, *6*, 895–901. [CrossRef]

- 112. Mukherjee, S.; Patra, C.R. Biologically synthesized metal nanoparticles: Recent advancement and future perspectives in cancer theranostics. *Future Sci.* **2017**. [CrossRef] [PubMed]
- Kumar, S.A.; Abyaneh, M.K.; Gosavi, S.; Kulkarni, S.K.; Pasricha, R.; Ahmad, A.; Khan, M. Nitrate reductase-mediated synthesis of silver nanoparticles from AgNO₃. *Biotechnol. Lett.* 2007, 29, 439–445. [CrossRef] [PubMed]
- 114. Kumar, S.A.; Ansary, A.A.; Ahmad, A.; Khan, M. Extracellular biosynthesis of CdSe quantum dots by the fungus, Fusarium oxysporum. *J. Biomed. Nanotechnol.* **2007**, *3*, 190–194. [CrossRef]
- 115. Ingle, A.; Gade, A.; Pierrat, S.; Sonnichsen, C.; Rai, M. Mycosynthesis of silver nanoparticles using the fungus Fusarium acuminatum and its activity against some human pathogenic bacteria. *Curr. Nanosci.* 2008, 4, 141–144. [CrossRef]
- 116. Durán, N.; Marcato, P.D.; Alves, O.L.; De Souza, G.I.; Esposito, E. Mechanistic aspects of biosynthesis of silver nanoparticles by several Fusarium oxysporum strains. *J. Nanobiotechnol.* **2005**, *3*, 8. [CrossRef] [PubMed]
- 117. Senapati, S.; Ahmad, A.; Khan, M.I.; Sastry, M.; Kumar, R. Extracellular biosynthesis of bimetallic Au–Ag alloy nanoparticles. *Small* 2005, 1, 517–520. [CrossRef] [PubMed]
- 118. Karbasian, M.; Atyabi, S.; Siadat, S.; Momen, S.; Norouzian, D. Optimizing nano-silver formation by Fusarium oxysporum PTCC 5115 employing response surface methodology. *Am. J. Agric. Biol. Sci.* 2008, *3*, 433–437.
- Ahmad, A.; Mukherjee, P.; Mandal, D.; Senapati, S.; Khan, M.I.; Kumar, R.; Sastry, M. Enzyme mediated extracellular synthesis of CdS nanoparticles by the fungus, Fusarium oxysporum. *J. Am. Chem. Soc.* 2002, 124, 12108–12109. [CrossRef] [PubMed]
- 120. Basavaraja, S.; Balaji, S.; Lagashetty, A.; Rajasab, A.; Venkataraman, A. Extracellular biosynthesis of silver nanoparticles using the fungus Fusarium semitectum. *Mater. Res. Bull.* **2008**, *43*, 1164–1170. [CrossRef]
- 121. Ingle, A.; Rai, M.; Gade, A.; Bawaskar, M. Fusarium solani: A novel biological agent for the extracellular synthesis of silver nanoparticles. *J. Nanopart. Res.* 2009, *11*, 2079. [CrossRef]
- 122. Balaji, D.; Basavaraja, S.; Deshpande, R.; Mahesh, D.B.; Prabhakar, B.; Venkataraman, A. Extracellular biosynthesis of functionalized silver nanoparticles by strains of Cladosporium cladosporioides fungus. *Colloids Surf. B* **2009**, *68*, 88–92. [CrossRef] [PubMed]
- 123. Gade, A.; Bonde, P.; Ingle, A.; Marcato, P.; Duran, N.; Rai, M. Exploitation of Aspergillus niger for synthesis of silver nanoparticles. *J. Biobased Mater. Biol.* **2008**, *2*, 243–247. [CrossRef]
- 124. Bhainsa, K.C.; D'souza, S. Extracellular biosynthesis of silver nanoparticles using the fungus Aspergillus fumigatus. *Colloids Surf. B* 2006, 47, 160–164. [CrossRef] [PubMed]
- 125. Kathiresan, K.; Manivannan, S.; Nabeel, M.; Dhivya, B. Studies on silver nanoparticles synthesized by a marine fungus, Penicillium fellutanum isolated from coastal mangrove sediment. *Colloids Surf. B* 2009, 71, 133–137. [CrossRef] [PubMed]
- Shaligram, N.S.; Bule, M.; Bhambure, R.; Singhal, R.S.; Singh, S.K.; Szakacs, G.; Pandey, A. Biosynthesis of silver nanoparticles using aqueous extract from the compactin producing fungal strain. *Process. Biochem.* 2009, 44, 939–943. [CrossRef]
- 127. Singaravelu, G.; Arockiamary, J.; Kumar, V.G.; Govindaraju, K. A novel extracellular synthesis of monodisperse gold nanoparticles using marine alga, Sargassum wightii Greville. *Colloids Surf. B* 2007, 57, 97–101. [CrossRef] [PubMed]
- 128. Lengke, M.F.; Fleet, M.E.; Southam, G. Morphology of gold nanoparticles synthesized by filamentous cyanobacteria from gold (I)- thiosulfate and gold (III)- chloride complexes. *Langmuir* 2006, 22, 2780–2787. [CrossRef] [PubMed]
- 129. Lloyd, J.R.; Yong, P.; Macaskie, L.E. Enzymatic recovery of elemental palladium by using sulfate-reducing bacteria. *Appl. Environ. Microbiol.* **1998**, *64*, 4607–4609. [PubMed]
- 130. Kashefi, K.; Lovley, D.R. Reduction of Fe (III), Mn (IV), and toxic metals at 100 C by Pyrobaculum islandicum. *Appl. Environ. Microbiol.* **2000**, *66*, 1050–1056. [CrossRef] [PubMed]
- 131. Bao, H.; Lu, Z.; Cui, X.; Qiao, Y.; Guo, J.; Anderson, J.M.; Li, C.M. Extracellular microbial synthesis of biocompatible CdTe quantum dots. *Acta Biomater.* **2010**, *6*, 3534–3541. [CrossRef] [PubMed]
- Du, L.; Jiang, H.; Liu, X.; Wang, E. Biosynthesis of gold nanoparticles assisted by Escherichia coli DH5α and its application on direct electrochemistry of hemoglobin. *Electrochem. Commun.* 2007, *9*, 1165–1170. [CrossRef]

- 133. Kalimuthu, K.; Babu, R.S.; Venkataraman, D.; Bilal, M.; Gurunathan, S. Biosynthesis of silver nanocrystals by Bacillus licheniformis. *Colloids Surf. B* **2008**, *65*, 150–153. [CrossRef] [PubMed]
- 134. Lee, J.-H.; Han, J.; Choi, H.; Hur, H.-G. Effects of temperature and dissolved oxygen on Se (IV) removal and Se (0) precipitation by Shewanella sp. HN-41. *Chemosphere* **2007**, *68*, 1898–1905. [CrossRef] [PubMed]
- 135. Juibari, M.M.; Abbasalizadeh, S.; Jouzani, G.S.; Noruzi, M. Intensified biosynthesis of silver nanoparticles using a native extremophilic Ureibacillus thermosphaericus strain. *Mater. Lett.* 2011, 65, 1014–1017. [CrossRef]
- 136. Sneha, K.; Sathishkumar, M.; Mao, J.; Kwak, I.; Yun, Y.-S. Corynebacterium glutamicum-mediated crystallization of silver ions through sorption and reduction processes. *Chem. Eng. J.* **2010**, *162*, 989–996. [CrossRef]
- 137. Suresh, A.K.; Pelletier, D.A.; Wang, W.; Broich, M.L.; Moon, J.-W.; Gu, B.; Allison, D.P.; Joy, D.C.; Phelps, T.J.; Doktycz, M.J. Biofabrication of discrete spherical gold nanoparticles using the metal-reducing bacterium Shewanella oneidensis. *Acta Biomater.* 2011, 7, 2148–2152. [CrossRef] [PubMed]
- Lengke, M.F.; Ravel, B.; Fleet, M.E.; Wanger, G.; Gordon, R.A.; Southam, G. Mechanisms of gold bioaccumulation by filamentous cyanobacteria from gold (III) – chloride complex. *Environ. Sci. Technol.* 2006, 40, 6304–6309. [CrossRef] [PubMed]
- Vigneshwaran, N.; Kathe, A.A.; Varadarajan, P.; Nachane, R.P.; Balasubramanya, R. Biomimetics of silver nanoparticles by white rot fungus, Phaenerochaete chrysosporium. *Colloids Surf. B* 2006, 53, 55–59. [CrossRef] [PubMed]
- Vigneshwaran, N.; Ashtaputre, N.; Varadarajan, P.; Nachane, R.; Paralikar, K.; Balasubramanya, R. Biological synthesis of silver nanoparticles using the fungus Aspergillus flavus. *Mater. Lett.* 2007, *61*, 1413–1418. [CrossRef]
- 141. Senapati, S.; Mandal, D.; Ahmad, A. Fungus mediated synthesis of silver nanoparticles: A novel biological approach. *Indian J. Phys. A* **2004**, *78*, 101.
- 142. Fayaz, A.M.; Balaji, K.; Kalaichelvan, P.; Venkatesan, R. Fungal based synthesis of silver nanoparticles—An effect of temperature on the size of particles. *Colloids Surf. B* **2009**, 74, 123–126. [CrossRef] [PubMed]
- 143. Agnihotri, M.; Joshi, S.; Kumar, A.R.; Zinjarde, S.; Kulkarni, S. Biosynthesis of gold nanoparticles by the tropical marine yeast Yarrowia lipolytica NCIM 3589. *Mater. Lett.* **2009**, *63*, 1231–1234. [CrossRef]
- 144. Konishi, Y.; Ohno, K.; Saitoh, N.; Nomura, T.; Nagamine, S.; Hishida, H.; Takahashi, Y.; Uruga, T. Bioreductive deposition of platinum nanoparticles on the bacterium Shewanella algae. *J. Biotechnol.* 2007, 128, 648–653. [CrossRef] [PubMed]
- 145. Sinha, A.; Khare, S.K. Mercury bioaccumulation and simultaneous nanoparticle synthesis by Enterobacter sp. cells. *Bioresour. Technol.* 2011, 102, 4281–4284. [CrossRef] [PubMed]
- 146. Babu, M.G.; Gunasekaran, P. Production and structural characterization of crystalline silver nanoparticles from Bacillus cereus isolate. *Colloids Surf. B* **2009**, *74*, 191–195. [CrossRef] [PubMed]
- 147. Kalishwaralal, K.; Deepak, V.; Pandian, S.R.K.; Kottaisamy, M.; BarathManiKanth, S.; Kartikeyan, B.; Gurunathan, S. Biosynthesis of silver and gold nanoparticles using Brevibacterium casei. *Colloids Surf. B* **2010**, *77*, 257–262. [CrossRef] [PubMed]
- 148. Ahmad, A.; Senapati, S.; Khan, M.I.; Kumar, R.; Ramani, R.; Srinivas, V.; Sastry, M. Intracellular synthesis of gold nanoparticles by a novel alkalotolerant actinomycete, Rhodococcus species. *Nanotechnology* 2003, 14, 824. [CrossRef]
- 149. Gericke, M.; Pinches, A. Biological synthesis of metal nanoparticles. *Hydrometallurgy* **2006**, *83*, 132–140. [CrossRef]
- 150. Thakkar, K.N.; Mhatre, S.S.; Parikh, R.Y. Biological synthesis of metallic nanoparticles. *Nanomedicine* **2010**, *6*, 257–262. [CrossRef] [PubMed]
- 151. Dauthal, P.; Mukhopadhyay, M. Noble metal nanoparticles: Plant-mediated synthesis, mechanistic aspects of synthesis, and applications. *Ind. Eng. Chem. Res.* **2016**, *55*, 9557–9577. [CrossRef]
- 152. Ahmad, A.; Senapati, S.; Khan, M.I.; Kumar, R.; Sastry, M. Extracellular biosynthesis of monodisperse gold nanoparticles by a novel extremophilic actinomycete, Thermomonospora sp. *Langmuir* 2003, *19*, 3550–3553. [CrossRef]
- 153. Mukherjee, P.; Ahmad, A.; Mandal, D.; Senapati, S.; Sainkar, S.R.; Khan, M.I.; Parishcha, R.; Ajaykumar, P.; Alam, M.; Kumar, R. Fungus-mediated synthesis of silver nanoparticles and their immobilization in the mycelial matrix: A novel biological approach to nanoparticle synthesis. *Nano Lett.* 2001, 1, 515–519. [CrossRef]

- 154. Mukherjee, P.; Ahmad, A.; Mandal, D.; Senapati, S.; Sainkar, S.R.; Khan, M.I.; Ramani, R.; Parischa, R.; Ajayakumar, P.; Alam, M. Bioreduction of AuCl4–ions by the fungus, Verticillium sp. and surface trapping of the gold nanoparticles formed. *Angew. Chem. Int. Ed.* **2001**, *40*, 3585–3588. [CrossRef]
- 155. Southam, G.; Beveridge, T.J. The occurrence of sulfur and phosphorus within bacterially derived crystalline and pseudocrystalline octahedral gold formed in vitro. *Geochim. Cosmochim. Acta* **1996**, *60*, 4369–4376. [CrossRef]
- 156. Sanghi, R.; Verma, P.; Puri, S. Enzymatic formation of gold nanoparticles using Phanerochaete chrysosporium. *Adv. Chem. Eng. Sci.* **2011**, *1*, 154. [CrossRef]
- 157. Konishi, Y.; Tsukiyama, T.; Tachimi, T.; Saitoh, N.; Nomura, T.; Nagamine, S. Microbial deposition of gold nanoparticles by the metal-reducing bacterium Shewanella algae. *Electrochim. Acta* 2007, 53, 186–192. [CrossRef]
- Mody, V.V.; Siwale, R.; Singh, A.; Mody, H.R. Introduction to metallic nanoparticles. *J. Pharm. Bioallied Sci.* 2010, 2, 282. [CrossRef] [PubMed]
- 159. Iravani, S. Green synthesis of metal nanoparticles using plants. Green Chem. 2011, 13, 2638–2650. [CrossRef]
- Nethi, S.K.; Mukherjee, S.; Veeriah, V.; Barui, A.K.; Chatterjee, S.; Patra, C.R. Bioconjugated gold nanoparticles accelerate the growth of new blood vessels through redox signaling. *Chem. Commun.* 2014, 50, 14367–14370. [CrossRef] [PubMed]
- Luo, C.-H.; Shanmugam, V.; Yeh, C.-S. Nanoparticle biosynthesis using unicellular and subcellular supports. NPG Asia Mater. 2015, 7, e209. [CrossRef]
- Korbekandi, H.; Iravani, S.; Abbasi, S. Production of nanoparticles using organisms. *Crit. Rev. Biotechnol.* 2009, 29, 279–306. [CrossRef] [PubMed]
- Mukherjee, S.; Patra, C.R. Therapeutic application of anti-angiogenic nanomaterials in cancers. *Nanoscale* 2016, *8*, 12444–12470. [CrossRef] [PubMed]
- 164. Wen, L.; Lin, Z.; Gu, P.; Zhou, J.; Yao, B.; Chen, G.; Fu, J. Extracellular biosynthesis of monodispersed gold nanoparticles by a SAM capping route. *J. Nanopart. Res.* **2009**, *11*, 279–288. [CrossRef]
- 165. Balakrishnan, S.; Mukherjee, S.; Das, S.; Bhat, F.A.; Raja, S.P.; Patra, C.R.; Arunakaran, J. Gold nanoparticles-conjugated quercetin induces apoptosis via inhibition of EGFR/PI3K/Akt-mediated pathway in breast cancer cell lines (MCF-7 and MDA-MB-231). *Cell Biochem. Funct.* 2017, 35, 217–231. [CrossRef] [PubMed]
- Moon, J.-W.; Rawn, C.J.; Rondinone, A.J.; Love, L.J.; Roh, Y.; Everett, S.M.; Lauf, R.J.; Phelps, T.J. Large-scale production of magnetic nanoparticles using bacterial fermentation. *J. Ind. Microbiol. Biotechnol.* 2010, 37, 1023–1031. [CrossRef] [PubMed]
- 167. Korbekandi, H.; Iravani, S.; Abbasi, S. Optimization of biological synthesis of silver nanoparticles using Lactobacillus casei subsp. casei. *J. Chem. Technol. Biotechnol.* **2012**, *87*, 932–937. [CrossRef]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).