

ORAL PRESENTATION

Open Access

TG4010 immunotherapy combined with first-line therapy in advanced non-small cell lung cancer (NSCLC): phase IIb results of the TIME study

Elisabeth Quoix¹, Lecia Sequist², John Nemunaitis³, Thaddeus Beck⁴, Piotr Jaskiewicz⁵, Jean-Philippe Oster⁶, Arnaud Scherpereel⁷, Erzsébet Juhász⁸, Zsuzsanna Mark⁹, Rosa Alvarez¹⁰, Saiama Waqar¹¹, Joseph Potz¹², Nandagopal Vrindavanam¹³, Anton Melnyk¹⁴, Helen Ross¹⁵, Jean-Marc Limacher^{16*}

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Background

TG4010 immunotherapy product is a poxvirus (MVA) coding for MUC1 tumor-associated antigen and interleukin-2. A previous study showed that a normal baseline level of Triple Positive Activated Lymphocytes (TrPAL, CD16+CD56+CD69+) might be a predictive biomarker for TG4010 efficacy in advanced NSCLC [1]. TIME is a double-blind Phase IIb/III study comparing the combination of first-line therapy with TG4010 or placebo (NCT01383148).

Patients and methods

Primary endpoint of the Phase IIb part of the study was to compare progression-free survival (PFS, according to RECIST 1.1) between TG4010 and placebo arms using a Bayesian design, in stage IV NSCLC patients whose tumor express MUC1. Secondary objectives were response rate, safety, survival and subgroup analyses according to stratification factors and level of TrPAL at baseline. A dynamic minimization procedure was applied at randomization for histology, prescription of Bevacizumab, type of chemotherapy, performance status and center.

Results

217 patients have been enrolled out of which 170 patients with a normal TrPAL level (pre-determined threshold) and an analysis of PFS was conducted in this cohort after 137 events of progression were recorded. The hazard ratio (HR) for PFS is 0.76 (95%CI: 0.54-1.06). This corresponds to a 97.5% Bayesian probability that the true HR

is <1, passing the threshold of 95% needed to consider the endpoint met in patients with normal TrPAL. TG4010 related adverse events were limited to mild or moderate fever and injection site reaction. Analysis in the 75% of patients with the lowest baseline level of TrPAL (three lowest quartiles, n = 152) shows a HR for PFS of 0.72 (95%CI: 0.50-1.03) consistent with the observation made in the previous study. Additional pre-planned analyses by subgroup show that patients with non-squamous tumors had a statistically significant improvement in PFS when treated with TG4010 (n = 145, HR = 0.67; CI: 0.46-0.97; p = 0.016) and especially when belonging to the three lowest quartiles (n = 131, HR = 0.63; 95% CI: 0.42-0.93, p = 0.009). Overall survival data will be presented at the time of the meeting.

Conclusion

These data confirm TG4010 efficacy and safety profile in stage IV NSCLC especially in patients with low level of TrPAL before treatment. They warrant the continuation of the TIME study with its Phase III part.

Authors' details

¹CHRU strasbourg, Strasbourg, France. ²Massachusetts General Hospital, Cambridge, MA, USA. ³Mary Crowley Medical Research Center, Dallas, TX, USA. ⁴Highlands Oncology Group, Fayetteville, AR, USA. ⁵Maria Skłodowska-Curie Memorial Cancer Center and Institute of Oncology - Warsaw, Warsaw, Poland. ⁶Centre Hospitalier de Colmar, Colmar, France. ⁷CHRU de Lille Hopital Calmette, Lille, France. ⁸Országos Korányi TBC és Pulmonológiai Intézet, Budapest, Hungary. ⁹Tudogyógyintézet Torokbalint, Torokbalint, Hungary. ¹⁰Hospital Gregorio Marañon, Madrid, Spain. ¹¹Washington University, St. Louis, WA, USA. ¹²Abington Hematology Oncology Associates Inc, Willow Grove, PA, USA. ¹³Signal Point Clinical Research Center, Middletown, OH, USA. ¹⁴Texas Oncology, P.A. - Abilene (South), Abilene, TX, USA. ¹⁵Mayo Clinic Arizona, Phoenix, AZ, USA. ¹⁶Transgene S.A., Illkirch Graffenstaden, France.

¹⁶Transgene S.A., Illkirch Graffenstaden, France
Full list of author information is available at the end of the article

Published: 6 November 2014

Reference

1. Quoix E, *et al*: Therapeutic vaccination with TG4010 and first-line chemotherapy in advanced NSCLC: a controlled phase 2B trial. *The Lancet Oncol* 2011, **12**:1125-33.

doi:10.1186/2051-1426-2-S3-O12

Cite this article as: Quoix *et al*: TG4010 immunotherapy combined with first-line therapy in advanced non-small cell lung cancer (NSCLC): phase IIb results of the TIME study. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):O12.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

