



# Genome Sequence of Human Cytomegalovirus Ig-KG-H2, a Variant of Strain KG Propagated in the Presence of Neutralizing Antibodies

Ahmed Al Qaffas,<sup>a</sup> Salvatore Camiolo,<sup>b</sup> Jenna Nichols,<sup>b</sup> Andrew J. Davison,<sup>b</sup> Amine Ourahmane,<sup>a</sup> Xiaohong Cui,<sup>a</sup> Mark R. Schleiss,<sup>c</sup> Laura Hertel,<sup>d</sup> Dirk P. Dittmer,<sup>e</sup> Michael A. McVoy<sup>a</sup>

<sup>a</sup>Department of Pediatrics, Virginia Commonwealth University, Richmond, Virginia, USA

<sup>b</sup>MRC-University of Glasgow Centre for Virus Research, Glasgow, United Kingdom

<sup>c</sup>Center for Infectious Diseases and Microbiology Translational Research, Division of Pediatric Infectious Diseases, Minneapolis, Minnesota, USA

<sup>d</sup>Department of Pediatrics, School of Medicine, University of California, San Francisco, Oakland, California, USA

<sup>e</sup>Lineberger Comprehensive Cancer Center Program in Global Oncology, Department of Microbiology and Immunology, Center for AIDS Research, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA

**ABSTRACT** Human cytomegalovirus shed in infant urine was isolated and serially passaged in fibroblasts in the presence or absence of neutralizing antibodies. Comparison of the genome sequences of representative viruses Ig-KG-H2 (passed with antibody) and  $\phi$ -KG-B5 (passed without antibody) revealed the presence of several mutations in each virus.

Human cytomegalovirus (HCMV) replicates poorly when first isolated from clinical material. However, upon serial passage in fibroblasts, improved replication and increased release of cell-free virus are conferred by disruptive mutations in the *RL13* gene and one or more of the contiguous genes *UL128*, *UL130*, and *UL131A* (1). The latter mutations disrupt assembly of a pentameric complex on the virion surface that is important for entry into epithelial and endothelial cells but not fibroblasts (2–6).

In our recent work, replicate fibroblast cultures were infected with HCMV in urine from a symptomatic congenitally infected infant (7). One lineage (Ig-KG) was passaged with HCMV-hyperimmune globulin (HIG) (CytoGam) in the culture medium, whereas the other ( $\phi$ -KG) was passaged in the absence of HIG.  $\phi$ -KG lost epithelial tropism and acquired frameshift mutations disrupting *RL13* and *UL131A*, whereas Ig-KG retained epithelial tropism and was intact in these genes after 22 passages. Long-term genetic stability of these lineages (and their mutations) was confirmed by isolating representative viruses by limiting dilution, i.e., Ig-KG-H2 from the Ig-KG passage 22 stock and  $\phi$ -KG-B5 from the  $\phi$ -KG passage 13 stock.

Preliminary Ion Torrent and targeted PCR/Sanger sequence analyses that were focused on protein-coding sequences identified mutations affecting five genes (*RL13*, *UL100*, *UL102*, *UL122*, and *UL131A*) in the parental Ig-KG and  $\phi$ -KG stocks that were also present in Ig-KG-H2 and  $\phi$ -KG-B5 (7). Here, we report the complete genome sequence of Ig-KG-H2 and compare it with that of  $\phi$ -KG-B5.

Ig-KG-H2 or  $\phi$ -KG-B5 virions were pelleted from culture supernatants by ultracentrifugation and treated with DNase I prior to DNA purification by proteinase K digestion, phenol-chloroform extraction, and ethanol precipitation (7). DNA samples (~100 ng) were sheared acoustically to an approximate size of 450 bp, and sequencing libraries were processed through seven PCR cycles using the LTP library preparation kit (KAPA Biosystems, Wilmington, MA, USA), employing NEBNext multiplex oligos for Illumina (New England Biolabs, Ipswich, MA, USA). Sequencing on an Illumina NextSeq mid-output 300-cycle cartridge generated 9,614,942 and 10,011,260 paired-end reads of 150

**Citation** Al Qaffas A, Camiolo S, Nichols J, Davison AJ, Ourahmane A, Cui X, Schleiss MR, Hertel L, Dittmer DP, McVoy MA. 2020. Genome sequence of human cytomegalovirus Ig-KG-H2, a variant of strain KG propagated in the presence of neutralizing antibodies. *Microbiol Resour Announc* 9:e00063-20. <https://doi.org/10.1128/MRA.00063-20>.

**Editor** Jelle Matthijssens, KU Leuven

**Copyright** © 2020 Al Qaffas et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Michael A. McVoy, [michael.mcvoy@vcuhealth.org](mailto:michael.mcvoy@vcuhealth.org).

**Received** 17 March 2020

**Accepted** 2 April 2020

**Published** 23 April 2020

**TABLE 1** Mutations identified in the Ig-KG-H2 and  $\phi$ -KG-B5 genomes

Gene	Protein	Mutant <sup>a</sup>	Mutation(s) <sup>b</sup>	Consequence
None	None	$\phi$ -KG-B5 <sup>c</sup>	1-bp deletion (C6372)	None
<i>RL13</i>	Membrane protein RL13	$\phi$ -KG-B5	10-bp deletion (CATTATTATT at positions 11661–11670)	Frameshift after residue 164
<i>UL57</i>	Single-stranded DNA-binding protein	Ig-KG-H2	C89864T substitution	Silent
<i>UL98</i>	DNase	Ig-KG-H2	C145699T substitution	Silent
<i>UL100</i>	Envelope glycoprotein M	Ig-KG-H2	C146566G substitution	E361D
			C146750A and T146751G substitutions	S300L
			C146794A substitution	Q286H
			C147608A substitution	S15I
<i>UL102</i>	Helicase-primase subunit	Ig-KG-H2	C147895G substitution	L23V
			C148861G substitution	L345V
			C149640T substitution	Silent
<i>UL122</i>	Regulatory protein IE2	$\phi$ -KG-B5	G171290C substitution	F384L
		Ig-KG-H2	G171315T substitution	S376Y
<i>UL131A</i>	Envelope protein UL131A	$\phi$ -KG-B5	1-bp insertion (T178079)	Frameshift after residue 27

<sup>a</sup> The virus in which each mutation occurred was identified by comparison with strain Merlin as a representative HCMV strain.

<sup>b</sup> Coordinates refer to the Ig-KG-H2 genome sequence.

<sup>c</sup> This is only nominally a mutant, as the mutation represents a difference in the number of nucleotides in a C tract that varies in length among HCMV strains.

nucleotides for Ig-KG-H2 and  $\phi$ -KG-B5, respectively. A pipeline included in the GRACy tool (<https://github.com/salvocamiolo/GRACy>) was used to perform *de novo* assembly of the Ig-KG-H2 reads. Briefly, reads that aligned with the Hg38 human reference sequence (GenBank [GCA\\_000001405.15](https://www.ncbi.nlm.nih.gov/nuccore/GCA_000001405.15)) using Bowtie 2 v. 2.3.1 (8) (with the end-to-end flag set) were removed, and sequencing adapters and low-quality reads were removed using Trim Galore v. 0.4.0 (<https://github.com/FelixKrueger/TrimGalore>) and PRINSEQ v. 0.20.4 (9), respectively. The remaining reads were normalized and assembled using SPAdes v. 3.12 (10), and the resulting contigs were ordered in relation to the HCMV reference strain Merlin genome sequence (GenBank accession number [AY446894.2](https://www.ncbi.nlm.nih.gov/nuccore/AY446894.2)). Gaps were closed using an overlap-layout-consensus algorithm implemented in GRACy, and the assembly was further refined by visualization in Tablet v. 1.19.09.03 (11) of a read alignment that had been generated using Bowtie 2. All tools were used with default parameters unless otherwise specified. The Ig-KG-H2 genome sequence consisted of 236,244 bp (G+C content, 57.4%) and was determined at an average coverage of 4,886 reads/nucleotide. The  $\phi$ -KG-B5 reads were aligned to the resulting Ig-KG-H2 genome sequence using Bowtie 2, and differences present in the entire population were identified manually using Tablet.

As reported previously (7),  $\phi$ -KG-B5 had disruptive mutations in *RL13* and *UL131A*, as well as a single amino acid substitution in *UL122* (Table 1). In contrast, Ig-KG-H2 lacked disruptive mutations in *RL13* and *UL128*, *UL130*, or *UL131A* but contained mutations resulting in four amino acid substitutions in *UL100*, two amino acid substitutions in *UL102*, and a distinct single amino acid substitution in *UL122*. Also, Ig-KG-H2 had two silent mutations in *UL57* and *UL98*. The availability of the genome sequences of Ig-KG-H2 and  $\phi$ -KG-B5 will facilitate studies of the relative importance of these mutations in the adaptation of Ig-KG-H2 to growth in the presence of HIG.

**Data availability.** The genome sequence of Ig-KG-H2 has been deposited in GenBank under accession number [MN274568](https://www.ncbi.nlm.nih.gov/nuccore/MN274568). Raw reads are available from the European Nucleotide Archive with accession numbers [ERR3988552](https://www.ebi.ac.uk/ena/record/ERR3988552) (Ig-KG-H2) and [ERR3988553](https://www.ebi.ac.uk/ena/record/ERR3988553) ( $\phi$ -KG-B5).

## ACKNOWLEDGMENTS

We thank CSL Behring (King of Prussia, PA, USA) for the kind gift of CytoGam.

This work was supported by grants R01AI088750 and R21AI073615 (National Institutes of Health; to M.A.M.), R01AI128912 (National Institutes of Health; to M.A.M. and L.H.), R01HD079918 (National Institutes of Health; to M.R.S.), P01CA019014 (National Institutes of Health; to D.P.D.), 6-FY17-849 (March of Dimes Birth Defects Foundation; to

M.R.S.), 204870/Z/16/Z (Wellcome; to A.J.D.), MC\_UU\_12014/3 (Medical Research Council; to A.J.D.), and LKR141973 and LKRD119165 (Merck & Co.; to M.A.M.).

## REFERENCES

1. Dargan DJ, Douglas E, Cunningham C, Jamieson F, Stanton RJ, Baluchova K, McSharry BP, Tomasec P, Emery VC, Percivalle E, Sarasini A, Gerna G, Wilkinson GW, Davison AJ. 2010. Sequential mutations associated with adaptation of human cytomegalovirus to growth in cell culture. *J Gen Virol* 91:1535–1546. <https://doi.org/10.1099/vir.0.018994-0>.
2. Hahn G, Revello MG, Patrone M, Percivalle E, Campanini G, Sarasini A, Wagner M, Gallina A, Milanese G, Koszinowski U, Baldanti F, Gerna G. 2004. Human cytomegalovirus UL131-128 genes are indispensable for virus growth in endothelial cells and virus transfer to leukocytes. *J Virol* 78:10023–10033. <https://doi.org/10.1128/JVI.78.18.10023-10033.2004>.
3. Wang D, Shenk T. 2005. Human cytomegalovirus virion protein complex required for epithelial and endothelial cell tropism. *Proc Natl Acad Sci U S A* 102:18153–18158. <https://doi.org/10.1073/pnas.0509201102>.
4. Adler B, Scrivano L, Ruzsics Z, Rupp B, Sinzger C, Koszinowski U. 2006. Role of human cytomegalovirus UL131A in cell type-specific virus entry and release. *J Gen Virol* 87:2451–2460. <https://doi.org/10.1099/vir.0.81921-0>.
5. Ryckman BJ, Rainish BL, Chase MC, Borton JA, Nelson JA, Jarvis MA, Johnson DC. 2008. Characterization of the human cytomegalovirus gH/gL/UL128-131 complex that mediates entry into epithelial and endothelial cells. *J Virol* 82:60–70. <https://doi.org/10.1128/JVI.01910-07>.
6. Freed DC, Tang Q, Tang A, Li F, He X, Huang Z, Meng W, Xia L, Finnefrock AC, Durr E, Espeseth AS, Casimiro DR, Zhang N, Shiver JW, Wang D, An Z, Fu TM. 2013. Pentameric complex of viral glycoprotein H is the primary target for potent neutralization by a human cytomegalovirus vaccine. *Proc Natl Acad Sci U S A* 110:E4997–E5005. <https://doi.org/10.1073/pnas.1316517110>.
7. Ourahmane A, Cui X, He L, Catron M, Dittmer DP, Al Qaffasaa A, Schleiss MR, Hertel L, McVoy MA. 2019. Inclusion of antibodies to cell culture media preserves the integrity of genes encoding RL13 and the pentameric complex components during fibroblast passage of human cytomegalovirus. *Viruses* 11:221. <https://doi.org/10.3390/v11030221>.
8. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
9. Schmieder R, Edwards R. 2011. Quality control and preprocessing of metagenomic datasets. *Bioinformatics* 27:863–864. <https://doi.org/10.1093/bioinformatics/btr026>.
10. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
11. Milne I, Stephen G, Bayer M, Cock PJ, Pritchard L, Cardle L, Shaw PD, Marshall D. 2013. Using Tablet for visual exploration of second-generation sequencing data. *Brief Bioinform* 14:193–202. <https://doi.org/10.1093/bib/bbs012>.