

RESEARCH ARTICLE

Urban air pollution and emergency department visits related to central nervous system diseases

Anna O. Lukina¹, Brett Burstein^{2,3}, Mieczysław Szyszkowicz^{1*}

1 Environmental Health Science and Research Bureau, Health Canada, Ottawa, Canada, **2** Division of Pediatric Emergency Medicine, Department of Pediatrics, Montreal Children's Hospital, McGill University Health Centre, Montreal, Quebec, Canada, **3** Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Quebec, Canada

* mszyszkowicz@yahoo.ca**OPEN ACCESS**

Citation: Lukina AO, Burstein B, Szyszkowicz M (2022) Urban air pollution and emergency department visits related to central nervous system diseases. PLoS ONE 17(6): e0270459. <https://doi.org/10.1371/journal.pone.0270459>

Editor: Julian Aherne, Trent University, CANADA

Received: June 4, 2021

Accepted: June 12, 2022

Published: June 27, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0270459>

Copyright: © 2022 Lukina et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All files are available from the <https://github.com/szyszkowiczm/NERVEToronto>.

Funding: The authors received no specific funding for this work.

Abstract

Ambient air pollution has been associated with adverse neurological health outcomes. Ambient pollutants are thought to trigger oxidative stress and inflammation to which vulnerable populations, such as elderly may be particularly susceptible. Our study investigated the possible association between concentrations of ambient air pollutants and the number of emergency department (ED) visits for nervous system disorders among people residing in a large Canadian city. A time-stratified case-crossover study design combining data from the National Ambulatory Care Reporting System (NACRS) and the National Air Pollution Surveillance (NAPS) between 2004 and 2015 was used. Two air quality health indices were considered in addition to specific pollutants, including carbon monoxide (CO), nitrogen dioxide (NO₂), sulfur dioxide (SO₂), ozone (O₃) and fine particulate matter (PM_{2.5}). Weather condition data were included in the models. ED visits with a discharge diagnosis were identified using ICD-10 codes (G00-G99). The analysis was stratified by sex and age, also by seasons. The associations were investigated in arrays organized as 18 strata and 15 time lags (in days) for each pollutant. Overall, 140,511 ED visits were included for the analysis. Most ED visits were related to episodic and paroxysmal diagnoses (G40-G47, 64%), with a majority of visits for migraines (G43, 39%). Among females, an increase of 0.1 ppm ambient CO was associated with an increased risk of paroxysmal diagnoses at day 1 (RR = 1.019 (95%CI 1.004–1.033)), day 6 (1.024 (1.010–1.039)) and day 7 (1.022 (1.007–1.036)). PM_{2.5} and SO₂, and air quality indices were similarly associated with ED visits for episodic and paroxysmal disorders in days 6 and 7. Findings highlight that ambient air pollution is associated with an increased number of ED visits for nervous system disorders, particularly visits for paroxysmal diagnoses.

1. Introduction

Air pollution is the fifth-leading cause of mortality, accounting for nearly 9% of associated mortalities worldwide [1]. The study of the detrimental health effects of ambient air pollution is complex, as the environment contains a mixture of various particles and gases, including volatile organic compounds, particulate matter (PM), nitrogen oxides (NO_x), ground-level

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: AD, Alzheimer's disease; AQHI, Air quality health index; CI, Confidence interval; CIHI, Canadian institute health information; CO, carbon monoxide; ED, Emergency department; ICD-10, International classification of diseases, 10th Edition; IQR, Interquartile range; IRB, Institutional review board; NACRS, National ambulatory care reporting system; NAPS, National air pollution surveillance; NO₂, nitrogen dioxide; O₃, ozone (O₃H8 = ozone measured as 8 hour maximum average); PD, Parkinson's disease; PM, Particulate matter; ppb, parts per billion; ppm, parts per million; RR, Relative risk; SO₂, sulfur dioxide; WHO, World health organization.

ozone (O₃), sulfur dioxide (SO₂), carbon monoxide (CO) and others. Moreover, pathogenesis is also dependent on weather conditions, including humidity, wind speed, atmospheric pressure, precipitation, temperature, and seasons. Fine PM with a particle aerodynamic diameter of 2.5 μm (PM_{2.5}) or smaller, NO₂, and O₃ are often studied because such pollutants pose a significant threat to human health [2]. Both NO₂ and PM_{2.5} are emitted directly into the atmosphere from industrial sources (i.e., burning of fossil fuels, including for power generation), forest fires, and vehicle exhaust, including heavy duty and diesel engine exhaust, whereas ground-level O₃ is formed mostly by photochemical reaction [3].

Exposure to air pollutants may result in more visits to doctors or emergency department (ED), hospital admissions, frequent use of prescription medications, loss of productivity, and general changes to personal quality of life. In the last decade it was discovered that exposure to some major air pollutants may lead to various neurological outcomes, especially in older adults, including dementia, Alzheimer's disease (AD) and Parkinson's disease (PD) [4–6], headaches and migraines [7–9], epilepsy and seizures [10, 11], overall cognitive decline [12], and depression [13]. For example, long-term exposure to PM_{2.5} for women over 70 years of age was associated with cognitive decline [14], as well as decreases in brain gray and white matter volume [15]. However, short-term exposure to ambient NO₂, PM_{2.5}, SO₂, and CO was associated with some neurological outcomes, including but not limited to depression [13], migraines [7], and epilepsy [10, 11, 16]. Two Canadian studies conducted in the largest and most populous provinces, Ontario and Quebec, found that when older Canadians are long-term exposed to ambient PM_{2.5} and NO₂, and other traffic-related air pollutants (i.e., ultrafine particles and black carbon), over time they may develop dementia [6, 17]. In a study conducted in five major Canadian cities, exposure to ambient SO₂ and PM_{2.5} was associated with more ED visits for headaches and migraine attacks, especially among females [7]. In China, the frequency of outpatient visits for epilepsy was higher for men exposed to ambient SO₂ and O₃, whereas women were more sensitive to NO₂ exposure [16]. However, in another Chinese study, acute exposure to ambient NO₂, CO, and PM_{2.5} was associated with increased hospitalization for epilepsy [10]. A recent review of studies in Europe, Asia, and America highlighted the relationship between exposure to ambient air pollution and neurologic development among the young, as well as cognitive decline among the elderly [18].

Vulnerable populations, including elderly people, newborns and young children, pregnant women, as well as those with underlying health conditions, appear to be particularly sensitive to the deleterious health effects of ambient air pollution [14, 19–22].

Understanding the relationship between air pollutants and human health is essential to estimate economic impacts and the burden of health system resource utilization. To date, there is a paucity of information addressing the association of air pollution exposure with a wide range of disorders of the nervous system over the full life course. Previous studies have focused predominantly on specific vulnerable populations and health outcomes [19, 20, 23], or on particular pollutants [4, 14, 23–26]. The objective of this study was to comprehensively examine potential associations between concentrations of ambient air pollution and the number of ED visits for central and peripheral nervous system diseases in a large urban Canadian city.

2. Materials and methods

2.1 Studied population

Health data on the number of daily ED visits related to physician-diagnosed central and peripheral nervous system diseases were based on the *International Classification of Diseases 10th Revision* (ICD-10) codes G00–G99 (Chapter VI: “Diseases of the nervous system”) [27], and were obtained from the National Ambulatory Care Reporting System (NACRS) database

[28] for the period of April 01, 2004 to December 31, 2015, inclusively (overall 4,292 days or 140 months). Briefly, the NACRS database is based on the Canadian Institute of Health Information (CIHI) reporting system in Canada, where the data are collected from hospitals and ambulatory care centres in the province of Ontario [29]. In 2016, the enumerated population of Toronto was 2,731,571 according to the Canadian Census Division 2016. The population density of this region is 4,334 people per square kilometer. The study population included all individuals from newborn to >60 years with a home addresses located in the area determined by the Census Division of Toronto. The primary outcome was the number of ED visits with a discharge diagnosis of all nervous system disorders (ICD-10 codes; G00-G99), and sub-analyzed according to diagnostic sub-categories, including “inflammatory diseases of the central nervous system” (G00-G09), “systemic atrophies primarily affecting the central nervous system” (G10-G14), “extrapyramidal and movement disorders” (G20-G26), “other degenerative diseases of the nervous system” (G30-G32), “demyelinating diseases of the central nervous system” (G35-G37), “episodic and paroxysmal disorders” (G40-G47), “nerve, nerve root and plexus disorders” (G50-G59), “polyneuropathies and other disorders of the peripheral nervous system” (G60-G64), “diseases of myoneural junction and muscle” (G70-G73), “cerebral palsy and other paralytic syndromes” (G80-G83), and “other disorders of the nervous system” (G90-G99).

2.2 Air pollutants and meteorological conditions data

Long-term air pollution data were obtained from the National Air Pollution Surveillance (NAPS) program, maintained by Environment and Climate Change Canada [30]. Hourly data from seven automated fixed-site monitoring stations with approximate maximum distance among them of 15 kilometers were averaged to estimate daily air pollution concentrations for the whole Census Division of Toronto. Five major ambient air pollutants were studied, which included CO, NO₂, PM_{2.5}, O₃, and SO₂. For each pollutant, daily measurements taken at hourly intervals were obtained. A 24-hour average was used for all air pollutants with the exception to O₃, which was calculated as a daily maximum 8-hour average. Additionally, the Air Quality Health Index (AQHI) was calculated, based on the composite of three pollutants, NO₂, PM_{2.5}, and O₃, measured as 24-hour averages in Toronto. The AQHI incorporates air pollutant concentrations and health risk estimations determined by mortality rates in large Canadian cities [31]. For the Canadian public, these values are rounded and shown as integer numbers on a scale (1–10 and 10+, and also expressed by colours from blue to brown) to demonstrate the risk related to ambient air quality, where lower-scale numbers represent lower health risks and higher-scale numbers represent higher health risks. The AQHI values are generated using the following formula:

$$AQHI = \frac{1000}{10.4} \times (e^{0.000537 \times O_3} + e^{0.000871 \times NO_2} + e^{0.000487 \times PM_{2.5}} - 3),$$

where the coefficients for air pollutants were estimated using mortality risks [31]. In addition, another value of the index, AQHI-x, was calculated using the same three air pollutants as AQHI, but O₃ was on a maximum 8-hour average. This index has a stronger representation of O₃ than in the AQHI calculations, because ground-level O₃ tends to reach higher concentrations during daytime, especially on hot sunny days. The constructed indices were applied to investigate simultaneous effects of three air pollutants on human health.

To control for seasonal fluctuations in the concentrations of some air pollutants, data were analyzed by season, defined as cold (October to March) and warm (April to September) periods, classified according to the mean temperatures for each month recorded in Toronto [30].

To control for variations in weather [32, 33], daily relative humidity and ambient temperature data were collected from one weather monitoring station located at the Toronto International airport, details of which are described elsewhere [34], and the values were stratified as potential confounders.

2.3 Statistical analyses

A time-stratified case-crossover analytical design was used for all measured and unmeasured time-invariant factors and confounders, such as socioeconomic factors or comorbidities [35]. The analyzed data on associations between air pollutants concentrations and the number of ED visits for nervous system outcomes were organized as time-series values with a day as the time unit. These data consist of daily counts of ED visits, daily concentrations of air pollutants, and daily values of ambient temperature and relative humidity. A time-stratified approach was applied to cluster the data using a calendar hierarchical structure, where days were nested in day of weeks, then further nested in months and then in years [36]. Such clusters grouped as four or five days and were separated from each other. Controlling for time variables (i.e., any trends or fluctuations, etc.) was undertaken as described previously [37]. In the fitted statistical models, temperature and relative humidity were added in the form of natural splines of three degrees of freedom. The concentrations and their lagged values were assigned for the corresponding days. A two-tailed test at the 0.05 significance level was applied. The coefficients (slope, $Beta$) related to air pollutants and their standard errors ($SEBeta$) were estimated by the applied statistical models. Using these values (i.e., $Beta$, $SEBeta$), relative risks (RR) could be calculated. The statistical analyses were performed as conditional quasi-Poisson regression models [37, 38]. The numerical calculations were done employing R statistical software using the procedure for *Generalized Non-linear Models* (the package *GNM*) with the option “quasi-poisson” [39]. The realized statistical models have the following form: “ModelFit = gnm(EDVisits~AirPollutant + ns(RelativeHumidity,3) + ns(Temperature,3)”. The options family = quasipoisson, eliminate = factor (Cluster) were included in the specification. The conditional Poisson model avoids estimating cluster parameters. It is conditional upon the total counts in each constructed cluster.

The conditional Poisson model is represented as a multinomial model and it is given by the formula

$$N_{i,C} | Y_{N_{i,C}} \sim Multinomial \left(\pi_i = \frac{\exp\{\beta^T x_i\}}{\sum_{j \in C} \exp\{\beta^T x_j\}} \right),$$

where $N_{i,C}$ is the number of events (counts) on the cluster and $N_{i,C} = \sum_i N_{i,C}$ is the sum of events in each cluster. The parameters related to cluster are eliminated by conditioning on the sum of events on each cluster [37]. Here i is a day when ED visits occurs, β is a row vector of the coefficients, and T denotes transposition. A vector x contains variables: air pollutant concentrations and weather factors.

In total, 2,160 statistical models {15 (time lags expressed as days) x 18 (strata) x 8 (air pollutants and air quality health indices)} were applied. Strata covered patients' demographic characteristics, including age group (0–10, 11–60 and >60 years old) and sex group (all, males and females), as well as seasonality expressed dichotomously. The numerical results from all models are listed in the Supplementary Materials and at the online location <https://github.com/szyszkowicz/NERVEToronto>. This location also contains histograms (air pollutants, temperature, and relative humidity) and the map of Toronto.

2.4 Research ethics

All datasets are publicly available and de-identified, as such this study was deemed exempt from review by the Health Canada Research Ethics Board.

3. Results

Between April 1, 2004 and December 31, 2015, there were a total of 140,511 ED visits related to diseases of the nervous system (all G00-G99 codes) (Table 1). Overall, a majority of ED visits were by females (59.5%), particularly in the age ranges of 11–60 years old and >60 years. A total of 89,708 visits (64% of all ED visits) for the nervous system diseases were for episodic and paroxysmal disorders (G40-G47), among which nearly 40% were related to migraine attacks (G43) (Table 2).

Table 3 summarizes the descriptive statistics on five major air pollutants, two air quality health indices and weather variables for the entire study period and by warm (April to September) and cold (October to March) season. During cold season, the mean ambient temperature was captured at 1.6°C (-22.2–23.5°C), while during warm season, the mean ambient temperature was 17.0°C (-4.2–31.2°C) (Table 2). The mean levels of some air pollutants, NO₂, PM_{2.5}, and SO₂, were slightly higher in warm season than in cold season; however, none of the annual mean values for the studied air pollutants was above the respective Canadian Ambient Air Quality Standards (CAAQS) [40]. Descriptive statistics for all major pollutants and weather factors are also available in S2 Table in S1 File.

Nervous system diseases (G00-G99): When examining ED data for all G00-G99 codes, a total of 115 consistent positive associations were found, especially for exposure to ambient NO₂ (24 positive associations) and ambient CO (25 positive associations) (S1 Fig in S1 File), with higher number of associations on lag days 0 and 7 with corresponding 27 associations and 33 associations, respectively (S2 Fig in S1 File). More ED visits were noted during the warm season (S3 Fig in S1 File).

Episodic and paroxysmal disorders (G40-G47): ED visits for episodic and paroxysmal disorders were more frequent at higher levels of CO and PM_{2.5}, with 29 and 27 positive associations, respectively (Fig 1). The majority of cases (39%) were due to all classes of migraines (G43), which was associated with higher ambient CO and PM_{2.5} exposure, especially on lag days 6 (54 associations) and 7 (47 associations) (Fig 2) and during colder season for females

Table 1. Descriptive statistics on the number of ED visits for diseases of the nervous system (ICD-10 codes: G00–G99) collected in the city of Toronto between April 1, 2004 and December 31, 2015.

	ED visits for nervous system diseases	%
All individuals	140,511	100
Sex:		
Male (M)	56,909	40.5
Female (F)	83,602	59.5
Ages (in years):		
0–10	2,314 (M) and 1,978 (F)	53.9 (M) and 46.1 (F)
11–60	36,177 (M) and 57,456 (F)	38.6 (M) and 61.4 (F)
60+	18,418 (M) and 24,168 (F)	43.2 (M) and 56.8 (F)
Seasons (months) ^a :		
Cold	67,631	48.1
Warm	72,880	51.9

^a—cold season (October–March), warm season (April–September), F- female, M- male.

<https://doi.org/10.1371/journal.pone.0270459.t001>

Table 2. Comparison of frequencies for each diagnostic sub-code (G00-G99).

Code	Meaning	ED visits (%)
G00-09	Inflammatory diseases of the central nervous system	1,913 (1.4)
G10-14	Systemic atrophies primarily affecting the central nervous system	472 (0.3)
G20-26	Extrapyramidal and movement disorders	4,075 (2.9)
G30-32	Other degenerative diseases of the nervous system	1,704 (1.2)
G35-37	Demyelinating diseases of the central nervous system	1,816 (1.3)
G40-47	Episodic and paroxysmal disorders:	89,708 (63.8):
	• Epilepsy/seizures (G40 and G41)	18,670 (20.8)
	• Migraine (G43)	34,864 (38.9)
	• Other headaches syndromes (G44)	8,159 (9.1)
	• Cerebral ischaemic attacks and other related syndromes (G45)	21,914 (24.4)
	• Sleep disorders (G47)	6,101 (6.8)
G50-59	Nerve, nerve root and plexus disorders	30,607 (21.8)
G60-64	Polyneuropathies and other disorders of the peripheral nervous system	3,436 (2.5)
G70-73	Diseases of myoneural junction and muscle	889 (0.6)
G80-83	Cerebral palsy and other paralytic syndromes	1,244 (0.9)
G90-99	Other disorders of the nervous system	4,650 (3.3)
	TOTAL	140,514 (100)

<https://doi.org/10.1371/journal.pone.0270459.t002>

especially (Fig 3). Ambient SO₂ exposure (20 positive associations) and the AQHI (27 positive associations) were found to be related to ED visits for episodic and paroxysmal disorders (Fig 1). The RR and associated 95% confidence intervals (CI) were then calculated based on the interquartile range (IQR), which was 0.1 ppm for CO. For episodic and paroxysmal disorders, the RR and associated 95% CIs were determined for females exposed to CO on lag day 1 (1.019 (1.004–1.033), lag day 6 (1.024 (1.010–1.039)), and lag day 7 (1.022 (1.007–1.036) (Fig 4).

There was no apparent relationship between seasonality, however, older age contributed to the exposure-effect association, especially among women ≥ 60 years old exposed to CO on the concurrent day (0 lag day) (Fig 4).

4. Discussion

4.1 Summary of findings

The present study found a significant association of air pollutants and ED visits for nervous system diseases. In particular, episodic and paroxysmal disorders were most frequent, and visits for migraines accounted for nearly 40% of all paroxysmal disorders. The results revealed that females were more prone to visit the ED after being acutely (lags of 0–1 or 6–7 days) exposed to ambient CO, SO₂, and PM_{2.5} pollutants. There was a consistent effect of ambient CO exposure on the number of ED visits for all nervous system diseases and episodic and paroxysmal disorders.

4.2 Potential mechanisms underlying the identified associations

Previous studies have shown a positive association between ambient air pollution and recurring headaches and migraine attacks [7–9, 32, 33, 41], especially when individuals are exposed to CO [42–44], PM_{2.5} [8, 41], and SO₂ [7, 9]. Headache pathophysiology, including migraines and vascular headaches, may be due to nociceptive stimuli, which may trigger changes in the vasodilation of the cranial blood vessels and/or sensory nerve fibers [45, 46], and air pollution

Table 3. Descriptive data on ambient air pollutants and meteorological conditions on a daily basis, collected in Toronto, Canada between April 1, 2004 and December 31, 2015.

Factors (units)	Seasons ^a				All months		CAAQS ^d
	Cold		Warm		Mean	IQR ^c	
Air pollutants	Mean	Min/Max ^b	Mean	Min/Max ^b			
PM _{2.5} (µg/m ³)	8.5	0.9/35.6	9.2	0.1/65.5	8.9	6.5	8.8
NO ₂ (ppb)	13.8	3.2/46.5	18.0	4.3/59.8	16.1	8.8	17.0
O ₃ (ppb)	23.9	1.7/56.6	23.2	2.4/62.2	23.5	12.8	N/A
O ₃ H8 (ppb)	42.3	11.0/94.0	44.9	9.0/107.0	43.7	19.0	62.0 (8-hour)
SO ₂ (ppb)	0.9	-0.5/5.3	1.8	0/ 12.0	1.4	1.2	5.0
CO (ppm)	0.3	0/0.7	0.3	0/1.1	0.3	0.1	N/A
Air quality indices							
AQHI	2.8	1.1/5.8	3.2	1.1/7.6	3.0	1.0	N/A
AQHI-x	4.0	1.6/8.0	4.7	1.7/10.3	4.4	1.5	N/A
Weather							
Temperature (° C)	1.6	-22.2/23.5	17.0	-4.2/31.2	9.5	16.7	-
Relative Humidity (%)	72.7	31.7/98.8	68.8	35.4/96.7	70.7	14.3	-

Notes:

^a- cold season (October to March) and warm season (April to September),

^b- Min is minimum and Max is maximum,

^c- IQR is difference between the third (75th percentile) and first (25th percentile) quartiles,

^d- Canadian Ambient Air Quality Standards for annual values of PM_{2.5}, NO₂, and SO₂ (CAAQS) (available on <https://www.ccme.ca/en/air-quality-report#slide-7>) are from 2020 for comparison purposes solely,

N/A-not applicable.

<https://doi.org/10.1371/journal.pone.0270459.t003>

is known to cause changes to vascular and neural activity [47, 48]. Ambient air pollutants may be related to systemic inflammation, oxidative stress, apoptosis [49, 50] or brain oxygenation and metabolism [45].

Results suggest a difference in response to air pollutants between males and females, and among various age groups. There were more ED encounters for CNS disorders by females (between 11 and 60 years of age compared to males, overall (Table 1) and for episodic and paroxysmal disorders (Fig 3)). Observed differences may be attributable to biological, physiological, and behavioral differences among sexes. Population-based studies suggest that air pollution is associated with headaches and migraine attacks, especially in women [7–9, 41, 51].

Previous environmental health studies showed stronger effects from acute air pollutants exposure on human health. In a European study with 12 volunteers of mean age of 24 years, nine volunteers developed prolonged headaches following CO inhalation during the two study days [51]. Additionally, significant increases in heart rate and facial skin blood flow were observed among the exposed volunteers [51]. In a cross-sectional study with 4,073 patients (mean age: 40 years), increased incidence of headaches was detected after CO poisoning, with further symptoms of nausea, dizziness, and shortness of breath [52]. In an American study, higher hospitalization rates and mortality due to CO poisoning were found, especially among older individuals of 75 years of age or above [53]. However, symptoms from prolonged CO exposure normally occurs when levels exceed 70 ppm [43], which is very high compared to levels found in the current study (0.3 ppm).

Several studies have demonstrated a negative impact of air pollution exposure on development and progression of neurodegenerative disorders, including PD [54] and multiple sclerosis [55, 56], as well as potential for violent behavior resulting in 3.13% increase in risk for

Air Pollution:	AQHIX		NO2	O3H8	SO2		Total		
	AQHI	CO	O3	PM2.5					
All	2	2	2	0	0	2	2	2	12
Female	3	2	3	2	0	0	3	2	15
Male	0	0	0	0	0	1	0	0	1
Warm All	2	1	3	2	0	1	2	1	12
Warm Female	2	2	3	2	0	0	3	1	13
Warm Male	1	1	1	0	0	1	2	0	6
Cold All	3	2	2	0	0	3	3	2	15
Cold Female	6	3	5	4	0	6	5	4	33
Cold Male	0	0	0	0	1	2	0	0	3
Age 0-10 All	0	0	0	0	0	0	0	0	0
Age 0-10 Female	0	0	0	0	0	0	0	0	0
Age 0-10 Male	1	1	0	2	0	0	2	1	7
Age 11-60 All	1	1	1	0	0	0	2	0	5
Age 11-60 Female	4	1	3	1	0	0	2	2	13
Age 11-60 Male	0	0	0	0	0	0	0	0	0
Age 60+All	1	2	3	1	0	1	0	2	10
Age 60+ Female	1	2	3	3	0	0	1	3	13
Age 60+ Male	0	0	0	0	1	2	0	0	3
Total	27	20	29	17	2	19	27	20	161

Fig 1. Total frequencies of all associations: 18 strata (rows), five air pollutants and two air quality health indices (columns) between ambient air pollutants levels and the number of ED visits for episodic and paroxysmal disorders (G40-G47) in Toronto, Canada from April 1, 2004 to December 31, 2015. For visual representation: 0 (green) colour represents others than positive statistically significant associations.

<https://doi.org/10.1371/journal.pone.0270459.g001>

homicide and inflicted injury [57]. Such increase in ED visits was observed especially among males, children and Hispanics exposed to ambient CO during warm season in California [57]. Possible explanation can be because of combination effect of the air pollutant and heat stress [57]. Such results are in agreement with another Canadian study showing that even small increases in ambient CO levels may be related to an increases in substance abuse visits [58]. There might be a clear hormesis associated with CO exposure, as such higher concentrations and prolonged exposure may result in neuroinflammation and disruption to blood-brain barrier and transfer of oxygen to the brain [42, 45, 50], but very small quantities may be used for therapeutic benefit in some cases [59].

Lag	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	Total
AQHI	0	0	0	0	0	1	8	9	3	2	3	0	0	1	0	27
AQHI-X	0	0	0	0	0	2	6	8	0	1	1	0	1	1	0	20
CO	2	4	1	0	0	4	9	7	1	0	0	1	0	0	0	29
NO ₂	1	1	2	0	0	2	4	5	0	0	1	1	0	0	0	17
O ₃	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	2
O ₃ H8	0	0	0	0	0	2	7	4	2	1	1	0	2	0	0	19
PM _{2.5}	0	0	0	0	0	0	10	9	3	3	1	0	0	1	0	27
SO ₂	0	0	0	0	0	0	8	5	3	2	2	0	0	0	0	20
Total	3	5	3	0	0	11	54	47	12	9	9	2	3	3	0	161

Fig 2. Total frequencies for all associations: Five air pollutants + two air quality health indices (rows) and 15 time lags of 0–14 days (columns), between exposure to urban air pollutants levels and the number of ED visits for episodic and paroxysmal disorders (G40-G47) in Toronto, Canada between April 1, 2004 and December 31, 2015. For visual representation: 0 (green) colour represents others than positive statistically significant associations.

<https://doi.org/10.1371/journal.pone.0270459.g002>

Exposure to other pollutants, such as PM_{2.5} and SO₂ may also contribute to central nervous system disorders. Exposure to both pollutants may result in higher prevalence of multiple sclerosis [55, 56] and higher risk of progression of PD [54] and AD [4]. In a Canadian study, exposure to ambient SO₂ and PM_{2.5} was associated with increased risks of ED visits for depression

Lag	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	Total
All	0	0	0	0	0	0	6	6	0	0	0	0	0	0	0	12
Female	0	1	0	0	0	0	6	6	0	2	0	0	0	0	0	15
Male	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1
Warm All	1	0	0	0	0	0	4	6	1	0	0	0	0	0	0	12
Warm Female	0	1	0	0	0	0	4	5	0	0	0	0	0	3	0	13
Warm Male	0	0	0	0	0	0	1	4	0	0	0	1	0	0	0	6
Cold All	0	0	0	0	0	1	5	3	4	0	0	0	2	0	0	15
Cold Female	0	2	2	0	0	2	7	7	4	5	3	0	1	0	0	33
Cold Male	0	0	0	0	0	1	2	0	0	0	0	0	0	0	0	3
Age 0-10 All	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Age 0-10 Female	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Age 0-10 Male	0	0	0	0	0	0	0	0	2	1	3	1	0	0	0	7
Age 11-60 All	0	0	0	0	0	0	1	4	0	0	0	0	0	0	0	5
Age 11-60 Female	0	1	0	0	0	0	4	6	1	0	1	0	0	0	0	13
Age 11-60 Male	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Age 60+All	1	0	0	0	0	3	5	0	0	0	1	0	0	0	0	10
Age 60+ Female	1	0	1	0	0	3	6	0	0	1	1	0	0	0	0	13
Age 60+ Male	0	0	0	0	0	1	2	0	0	0	0	0	0	0	0	3
Total	3	5	3	0	0	11	54	47	12	9	9	2	3	3	0	161

Fig 3. Total frequencies of all associations for ambient air pollutants and the number of ED visits for related episodic and paroxysmal disorders (G40-G47). Eighteen strata (classified by patients' sex and age, as well as cold vs. warm seasons) examined and arranged in the rows, and 15 lags (expressed as days) are arranged in columns. For visual representation: 0 (green) colour represents other than positive statistically significant associations.

<https://doi.org/10.1371/journal.pone.0270459.g003>




Fig 4. Relative risks (RR) and 95% confidence intervals (95%CI) for an increase in a one interquartile range (CO, IQR = 0.1 ppm). ED visits diagnosed with the ICD-10 codes G40–G47 for the entire study period in Toronto between 2004 and 2015.

<https://doi.org/10.1371/journal.pone.0270459.g004>

[60]. Short-term exposure to ambient SO₂ resulted in increased 1.12 odds ratio of suicide risks, especially among males and well-educated individuals [61].

4.3 Strengths and limitations of the current study

The present study has several strengths. First, this study used ten years of longitudinal data on exposure and incorporated daily fluctuations in ambient air pollution concentrations into models estimating ED visits for nervous system diseases. To estimate the relative risks of a particular health outcome, the frequency of exposure to pollution prior to the health outcome is compared to control times (for example, a health outcome taken on a Monday is compared to those on other Mondays of the month). In addition, a case-crossover approach controls for all measured and unmeasured confounding factors, including socioeconomic and co-morbidities. The study considered the individual five major ambient air pollutants, as well as multi-pollutants expressed in AQHI (and AQHI-x) indices. The current study also considered both sexes across the lifespan (from newborn to ≥60 years of age) for patients attending the hospitals' ED.

There are some limitations to the present study. This study was unable to control for individuals' exposure duration, as there are disparities in individuals' lifestyles and habits. The study was also unable to control for personal exposure, including indoor and in-vehicle exposure, as well as proximity to major and local roads [62], as the current study was solely dependent on air pollution concentrations captured by seven fixed-site automated monitoring

stations. Secondly, it is possible that some admission data reported to the national NACRS database by facilities may be incomplete or inaccurate due to human error [29]. Additionally, previous patients' health information was not available, and thus, the study assessed the daily changes in air pollution concentrations and the number of ED visits for nervous system diseases in the given study period. Lastly, the multiplicity of comparisons among all five studied air pollutants, time lags of 14 days and 18 strata may introduce a possibility of erroneous associations (type 1 error). However, the study found consistent positive association between exposure to ambient air pollutants and nervous system diseases, as well as for the episodic and paroxysmal disorders alone.

5. Summary and future perspectives

In the present study, consistent positive exposure-effect associations were found between acute (i.e., concurrent day or a weeklong) exposure to ambient CO, PM_{2.5} and SO₂ concentrations and more frequent utilization of ED for CNS diseases (i.e., episodic and paroxysmal disorders), and perhaps, attributable to migraine attacks among patients residing in such a large city as Toronto. Although the biological plausibility of the observed association between ambient air pollution and episodic and paroxysmal disorders, including headaches and migraine attacks, is still remaining unclear. Additionally, further studies should focus on other geographical regions with similar environmental exposures to either confirm or refute the consistency of the findings presented here.

Supporting information

S1 File.
(DOCX)

Acknowledgments

The authors acknowledge Environment and Climate Change Canada for providing the air pollution data from the National Air Pollution Surveillance (NAPS) network.

Parts of this material are based on data and information compiled and provided by the Canadian Institute for Health Information (CIHI). However, the analyses, conclusions, opinions and statements expressed herein are not necessarily those of CIHI.

Author Contributions

Conceptualization: Mieczysław Szyszkowicz.

Data curation: Mieczysław Szyszkowicz.

Formal analysis: Anna O. Lukina, Brett Burstein.

Investigation: Mieczysław Szyszkowicz.

Methodology: Mieczysław Szyszkowicz.

Project administration: Mieczysław Szyszkowicz.

Software: Mieczysław Szyszkowicz.

Supervision: Mieczysław Szyszkowicz.

Validation: Anna O. Lukina, Brett Burstein.

Visualization: Anna O. Lukina.

Writing – original draft: Mieczysław Szyszkowicz.

Writing – review & editing: Anna O. Lukina, Brett Burstein.

References

1. IHME and HEI. State of Global Air/2019. Institute for Health Metrics and Evaluation, and Health Effects Institute. www.stateofglobalair.org/ (accessed 27-11-2020).
2. Health Canada. Health impacts of air pollution in Canada. Estimates of morbidity and premature mortality outcomes. 2019. http://publications.gc.ca/collections/collection_2019/sc-hc/H144-51-2019-eng.pdf (accessed 11-09-2020).
3. WHO (World Health Organization). Ambient (outdoor) air pollution. 2018. [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health) (accessed 09-09-2020).
4. Kioumourtzoglou M-A, Schwartz JD, Weisskopf MG, Melly SJ, Wang Y, Dominici F, et al. Long-term PM2.5 exposure and neurological hospital admissions in the Northeastern United States. *Environ. Health Perspect.* 2016; 124: 23–29. <http://dx.doi.org/10.1289/ehp.1408973> PMID: 25978701
5. Chen T-M, Gokhale J, Shofer S, Kuschner WG. Outdoor air pollution: ozone health effects. *Am. J. Med. Sci.* 2007; 333(4): 244–248. <https://doi.org/10.1097/MAJ.0b013e31803b8e8c> PMID: 17435419
6. Smargiassi A, Sidi EAL, Robert L-E, Plante C, Haddad M, Gamache P, et al. Exposure to ambient air pollutants and the onset of dementia in Quebec, Canada. *Environ. Res.* 2020; 190: 109870. <https://doi.org/10.1016/j.envres.2020.109870> PMID: 32739624
7. Szyszkowicz M, Kaplan GG, Grafstein E, Rowe BH. Emergency department visits for migraine and headache: a multi-city study. *Int. J. Occup. Med. Env.* 2009a; 22(3): 235–242. <https://doi.org/10.2478/v10001-009-0024-5> PMID: 19819836
8. Szyszkowicz M, Stieb DM, Rowe BH. Air pollution and daily ED visits for migraine and headache in Edmonton, Canada. *Am. J. Emerg. Med.* 2009b; 27: 391–396.
9. Szyszkowicz M, Rowe BH, Kaplan GG. Ambient sulfur dioxide exposure and emergency department visits for migraine in Vancouver, Canada. *Int. J. Occup. Med. Env.* 2009c; 22(1):7–12.
10. Bao X, Tian X, Yang C, Li Y, Hu Y. Association between ambient air pollution and hospital admission for epilepsy in Eastern China. *Epilepsy Res.* 2019; 152: 52–58. <https://doi.org/10.1016/j.eplepsyres.2019.02.012> PMID: 30909052
11. Cakmak S, Dales RE, Vidal CB. Air pollution and hospitalization for epilepsy in Chile. *Environ. Int.* 2010; 36: 501–505. <https://doi.org/10.1016/j.envint.2010.03.008> PMID: 20452673
12. Schikowski T, Altuğ H. The role of air pollution in cognitive impairment and decline. *Neurochem. Int.* 2020; 136: 104708. <https://doi.org/10.1016/j.neuint.2020.104708> PMID: 32092328
13. Wei F, Wu M, Qian S, Li D, Jin M, Wang J, et al. Association between short-term exposure to ambient air pollution and hospital visits for depression in China. *Sci. Total Environ.* 2020; 724: 138207. <https://doi.org/10.1016/j.scitotenv.2020.138207> PMID: 32268289
14. Weuve J, Puett RC, Schwartz J, Yanosky JD, Laden F, Grodstein F. Exposure to particulate air pollution and cognitive decline in older women. *Arch. Intern. Med.* 2012; 172(3): 219–227. <https://doi.org/10.1001/archinternmed.2011.683> PMID: 22332151
15. Casanova R, Wang X, Reyes J, Akita Y, Serre ML, Vizuete W, et al. A voxel-based morphometry study reveals local brain structural alterations associated with ambient fine particles in older women. *Front. Hum. Neurosci.* 2016; 10:495. <https://doi.org/10.3389/fnhum.2016.00495> PMID: 27790103
16. Xu C, Fan Y-N, Kan H-D, Chen R-J, Liu J-H, Li Y-F, et al. The novel relationship between urban air pollution and epilepsy: a time series study. *PLoS ONE* 2016; 11(8): e0161992. <https://doi.org/10.1371/journal.pone.0161992> PMID: 27571507
17. Chen H, Kwong JC, Copes R, et al. Exposure to ambient air pollution and the incidence of dementia: a population-based cohort study. *Environ. Int.* 2017; 108: 271–277. <https://doi.org/10.1016/j.envint.2017.08.020> PMID: 28917207
18. Clifford A, Lang L, Chen R, et al. Exposure to air pollution and cognitive functioning across the life course—a systematic literature review. *Environ. Res.* 2016; 147: 383–398. <https://doi.org/10.1016/j.envres.2016.01.018> PMID: 26945620
19. Villeneuve PJ, Chen L, Stieb D, Rowe BH. Associations between outdoor air pollution and emergency department visits for stroke in Edmonton, Canada. *Eur. J. Epidemiol.* 2006; 21: 689–700. <https://doi.org/10.1007/s10654-006-9050-9> PMID: 17048082

20. Calderón-Garcidueñas L, Torres-Jardón R, Kulesza RJ, Mansour Y, González-González LO, González-Maciele A, et al. Alzheimer disease starts in childhood in polluted Metropolitan Mexico City. A major health crisis in progress. *Environ. Res.* 2020; 183: 109137. <https://doi.org/10.1016/j.envres.2020.109137> PMID: 32006765
21. Lee K-S, Lim Y-H, Choi Y-J, Kim S, Bae HJ, Han C, et al. Prenatal exposure to traffic-related air pollution and risk of congenital diseases in South Korea. *Environ. Res.* 2020; 191: 110060. <https://doi.org/10.1016/j.envres.2020.110060> PMID: 32805245
22. Zhu X, Ji X, Shou Y, Huang Y, Hu Y, Wang H. Recent advances in understanding the mechanisms of PM2.5-mediated neurodegenerative diseases. *Toxicol. Lett.* 2020; 329: 31–37. <https://doi.org/10.1016/j.toxlet.2020.04.017> PMID: 32360789
23. Sørensen M, Daneshvar B, Hansen M, Dragsted LO, Hertel O, Knudsen L, et al. Personal PM2.5 exposure and markers of oxidative stress in blood. *Environ. Health Persp.* 2003; 111(2): 161–165. <https://doi.org/10.1289/ehp.111-1241344> PMID: 12573899
24. Yu Y, Yao S, Dong H, Wang L, Wang C, Ji X, et al. Association between short-term exposure to particulate matter air pollution and cause-specific mortality in Changzhou, China. *Environ. Res.* 2019; 170: 7–15. <https://doi.org/10.1016/j.envres.2018.11.041> PMID: 30554054
25. Andersen ZJ, Kristiansen LC, Andersen KK, Olsen TS, Hvidberg M, Jensen SS, et al. Stroke and long-term exposure to outdoor air pollution from nitrogen dioxide. *Stroke* 2012; 43: 320–325. <https://doi.org/10.1161/STROKEAHA.111.629246> PMID: 22052517
26. Li H, Xin X. Nitrogen dioxide (NO₂) pollution as a potential risk factor for developing vascular dementia and its synaptic mechanisms. *Chemosphere* 2013; 92: 52–58. <https://doi.org/10.1016/j.chemosphere.2013.02.061> PMID: 23535468
27. WHO (World Health Organization). International Statistical Classification of Diseases and Related Health Problems 10th Revision. 2016. <https://icd.who.int/browse10/2016/en> (accessed 09-09-2020).
28. The National Ambulatory Care Reporting System (NACRS), CIHI, Canada; 2020. <https://www.cihi.ca/en/national-ambulatory-care-reporting-system-metadata> (accessed 14-05-2020).
29. Gibson D, Richards H, Chapman A. The national ambulatory care reporting system: factors that affect the quality of its emergency data. *Int. J. Information Quality* 2008; 2(2): 97.
30. NAPS. Web site: <http://maps-cartes.ec.gc.ca/rmspa-naps/data.aspx>, Canada. 2020. <https://open.canada.ca/data/en/dataset/1b36a356-defd-4813-acea-47bc3abd859b> (accessed 14-05-2020).
31. Stieb DM, Burnett RT, Smith-Doiron M, Brion O, Shin HH, Economou V. A new multipollutant, no-threshold air quality health index based on short-term associations observed in daily time-series analyses. *J. Air Waste Manag. Assoc.* 2008; 58(3):435–450. <https://doi.org/10.3155/1047-3289.58.3.435> PMID: 18376646
32. Li W, Bertisch SM, Mostofsky E, Buettner C, Mittleman MA. Weather, ambient air pollution, and risk of migraine headaches onset among patients with migraine. *Environ. Int.* 2019; 132: 105100. <https://doi.org/10.1016/j.envint.2019.105100> PMID: 31446321
33. Lee H, Myung W, Cheong H-K, Yi S-M, Hong Y-C, Cho S-I, et al. Ambient air pollution exposure and risk of migraine: synergistic effect with high temperature. *Environ. Int.* 2018; 121: 383–391. <https://doi.org/10.1016/j.envint.2018.09.022> PMID: 30245361
34. Vicedo-Cabrera AM, Sera F, Liu C. et al. Short term association between ozone and mortality: global two stage time series study in 406 locations in 20 countries. *BMJ* 2020, 368: m108. <https://doi.org/10.1136/bmj.m108> PMID: 32041707
35. Maclure M. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am. J. Epidemiol.* 1991; 133:144–153. <https://doi.org/10.1093/oxfordjournals.aje.a115853> PMID: 1985444
36. Janes H, Sheppard L, Lumley T. Case-crossover analyses of air pollution exposure data: referent selection strategies and their implications for bias. *Epidemiology* 2005; 16:717–726. <https://doi.org/10.1097/01.ede.0000181315.18836.9d> PMID: 16222160
37. Armstrong BG, Gasparri A, Tobias A. Conditional Poisson models: a flexible alternative to conditional logistic case cross-over analysis. *BMC Med. Res. Methodol.* 2014; 14:122. <https://doi.org/10.1186/1471-2288-14-122> PMID: 25417555
38. Szyszkowicz M. Use of generalized linear mixed models to examine the association between air pollution and health outcomes. *Int. J. Occup. Med. Environ. Health* 2006; 19:224–227. <https://doi.org/10.2478/v10001-006-0032-7> PMID: 17402217
39. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2018. URL <https://www.R-project.org/>.

40. CCME. Canadian Ambient Air Quality Standards (CAAQS. 2020. <https://www.ccme.ca/en/air-quality-report#slide-7> (accessed 31-03-2021).
41. Dales RE, Cakmak S, Vidal CB. Air pollution and hospitalization for headache in Chile. 2009; *Am. J. Epidemiol.* 170(8): 1057–1066. <https://doi.org/10.1093/aje/kwp217> PMID: 19741041
42. Arnggrim N, Schytz HW, Hauge MK, Ashina M. Carbon monoxide may be an important molecule in migraine and other headaches. *Cephalalgia* 2014; 34(14): 1169–80. <https://doi.org/10.1177/0333102414534085> PMID: 24816616
43. Kanburoglu MK, Cizmeci MN, Akelma AZ. A rare cause of chronic headache that may be misdiagnosed as migraine: chronic carbon monoxide poisoning. *Turkiye Acil. Tip. Dergisi.* 2014; 14(3): 132–134. <https://doi.org/10.5505/1304.7361.2014.00868> PMID: 27331183
44. Szyszkowicz M. Ambient air pollution and daily emergency department visits for headache in Ottawa, Canada. *Headache* 2008; 48:1076–1081. <https://doi.org/10.1111/j.1526-4610.2007.01039.x> PMID: 18218009
45. Moskowitz MA. The neurobiology of vascular head pain. *Ann. Neurol.* 1984; 16(2): 157–68. <https://doi.org/10.1002/ana.410160202> PMID: 6206779
46. Peroutka SJ. Neurogenic inflammation and migraines: implications for therapeutics. *Mol. Interv.* 2005; 5(5): 304–11. <https://doi.org/10.1124/mi.5.5.10> PMID: 16249526
47. Calderón-Garcidueñas L, Franco-Lira M, Hanriquez-Roldán C, Osnaya N, González-Maciél A, Reynoso-Robles R, et al. Urban air pollution: influences on olfactory function and pathology in exposed children and young adults. *Exp. Toxicol. Pathol.* 2010; 62: 91–102. <https://doi.org/10.1016/j.etp.2009.02.117> PMID: 19297138
48. Ehsanifar M, Montazeri Z, Taheri MA, Rafati M, Behjati M, Karimian M. Hippocampal inflammation and oxidative stress following exposure to diesel exhaust nanoparticles in male and female mice. *Neurochem. Int.* 2021; 145: 104989. <https://doi.org/10.1016/j.neuint.2021.104989> PMID: 33582162
49. Kim H, Kim W-H, Kim Y-Y, Park H-Y. Air pollution and central nervous system disease: a review of the impact of fine particulate matter on neurological disorders. *Frontier Publ. Health* 2020; 8:57533. <https://doi.org/10.3389/fpubh.2020.575330> PMID: 33392129
50. Hanafy KA, Oh J, Otterbein LE. Carbon monoxide and the brain: time to rethink the dogma. *Curr. Pharm. Des.* 2013; 19(15): 2771–2775. <https://doi.org/10.2174/1381612811319150013> PMID: 23092321
51. Ghanizada H, Arnggrim N, Schytz HW, Olesen J, Ashina M. Carbon monoxide inhalation induces headache but no migraine in patients with migraine without aura. *Cephalalgia* 2018; 38(13): 1940–1949. <https://doi.org/10.1177/0333102418765771> PMID: 29540069
52. Koyuncu S, Bol O, Ertan T, Gunay N, Akdogan H.I. The detection of occult CO poisoning through noninvasive measurement of carboxyhemoglobin: a cross-sectional study. *Am J. Emerg. Med.* 2020; 38(6): 1110–1114. <https://doi.org/10.1016/j.ajem.2019.158383> PMID: 31416641
53. Harduar-Morano L, Watkins S. Review of unintentional non-fire-related carbon monoxide poisoning morbidity and mortality in Florida, 1999–2007. *Public Health Rep.* 2011; 126: 240–250. <https://doi.org/10.1177/003335491112600215> PMID: 21387954
54. Hu C, Fang Y, Li F, Dong B, Hua X, Jiang W, et al. Association between ambient air pollution and Parkinson's disease: systematic review and meta-analysis. *Environ. Res.* 2019; 168: 448–459. <https://doi.org/10.1016/j.envres.2018.10.008> PMID: 30391837
55. Lavery AM, Waubant E, Casper TC, Roalstad S, Candee M, Rose J, et al. Urban air quality and associations with pediatric multiple sclerosis. *Ann. Clin. Transl. Neur.* 2018; 5(10): 1146–1153. <https://doi.org/10.1002/acn3.616> PMID: 30349849
56. Abbaszadeh S, Tabary M, Aryannejad A, Abolhasani R, Araghi F, Khareshi I, et al. Air pollution and multiple sclerosis: a comprehensive review. *Neurol. Sci.* 2021; 42: 4063–4072. <https://doi.org/10.1007/s10072-021-05508-4> PMID: 34341860
57. Thilakaratne RA, Malig BJ, Basu R. Examining the relationship between ambient carbon monoxide, nitrogen dioxide, and mental health-related emergency department visits in California, USA. *Sci. Total Environ.* 2020; 746: 140915. <https://doi.org/10.1016/j.scitotenv.2020.140915> PMID: 32745847
58. Szyszkowicz M, Thompson EM, Colman I, Rowe BH. Ambient air pollution exposure and emergency department visits for substance abuse. *PLoS ONE* 2018; 13(6): e0199826. <https://doi.org/10.1371/journal.pone.0199826> PMID: 29958279
59. Goebel U, Wollborn J. Carbon monoxide in intensive care medicine-time to start the therapeutic application. *Intens. Care Med. Exp.* 2020; 8:2. <https://doi.org/10.1186/s40635-020-0292-8> PMID: 31919605
60. Szyszkowicz M, Kousha T, Kingsbury M, Colman I. Air pollution and emergency department visits for depression: a multicity case-crossover study. *Environ. Health Insights* 2016; 10: 155–161. <https://doi.org/10.4137/EHI.S40493> PMID: 27597809

61. Lin G, Li L, Song Y, Zhou Y, Shen S, Ou C. The impact of ambient air pollution on suicide mortality: a case-crossover study in Guangzhou, China. *Environ. Health* 2016; 15:90. <https://doi.org/10.1186/s12940-016-0177-1> PMID: 27576574
62. Özkaynak H, Baxter LK, Burke J. Evaluation and application of alternative air pollution exposure metrics in air pollution epidemiology studies. *J. Expo. Sci. Env. Epid.* 2013; 23: 565. <https://doi.org/10.1038/jes.2013.50> PMID: 24145199