

## CASE REPORT

# Successful surgical treatment approach for mitral valve vegetation of infective endocarditis after severe soft tissue infection with mediastinitis

Kenichiro Uchida<sup>1</sup>  | Yosuke Takahashi<sup>2</sup> | Toshihiko Shibata<sup>2</sup> | Yasumitsu Mizobata<sup>1</sup>

<sup>1</sup>Department of Traumatology and Critical Care Medicine, Graduate School of Medicine, Osaka City University, Osaka, Japan

<sup>2</sup>Department of Cardiovascular Surgery, Graduate School of Medicine, Osaka City University, Osaka, Japan

**Correspondence**

Kenichiro Uchida, 1-5-7 Asahimachi, Abeno-ku, Osaka-City, Osaka, Japan.  
Email: cvs.uchida@gmail.com

**Abstract**

Transesophageal echocardiography is mandatory if you do suspect infective endocarditis. By approaching via a small right thoracotomy, vegetectomy and mitral valvuloplasty following severe mediastinitis were successfully accomplished without any complications.

**KEYWORDS**

infective endocarditis, mediastinitis, mitral valve, sepsis, small right thoracotomy

## 1 | INTRODUCTION

Infective endocarditis is commonly lethal, and currently, there are several guidelines recommending indications for surgical interventions.<sup>1-4</sup> However, there are many unsolved and debatable problems related to the appropriate timing of surgery, the surgical approach, and minimizing the risk of comorbidities. Usually, there is evidence of mobile vegetation of over 10 mm in size, for which the guidelines recommend immediate surgical vegetectomy,<sup>1</sup> however, sometimes the patient's condition does not allow surgical intervention to proceed.

## 2 | CASE REPORT

A 57-year-old Asian man with a medical history of hypertension and type II diabetes mellitus was hospitalized because of fever, left shoulder pain, and mild unconsciousness. He also had frozen shoulder syndrome and had undergone brachial plexus block anesthesia administered by a local physician about one week before admission. Physical examination showed a significant skin rash and warmth around his neck,

anterior thoracic wall, and left upper arm with tenderness. No cardiac murmur or abnormal respiratory sounds were heard on auscultation. His blood pressure was 142/108 mm Hg, heart rate was regular at 122 beats/min, respiratory rate was 30 breaths/min, and his body temperature was 38.3°C. Serial hematologic workup is showed in Table 1.

Contrast-enhanced computed tomography (CECT) on admission revealed expansive fluid collection dorsal to the sternal notch, between the pectoralis major and minor muscles, and around the axillary nerve (Figure 1). No valve regurgitation or signs of vegetation were observed on transthoracic echocardiography (TTE) at this time.


After patient admission, we immediately performed surgical debridement with an adequate incision from the suprasternal notch to the axilla and brachial lesion. After washing out pus-like fluid collection, we irrigated the site for the next 4 days. As the targeted bacteria was revealed to be methicillin-sensitive *Staphylococcus aureus* (MSSA) and surgically debrided site became uncontaminated, we applied negative pressure wound therapy (NPWT). We also deescalated the antibiotic therapy to cefazolin from initial ticoplanin, doripenem hydrate, and clindamycin and was continued 8 weeks after admission.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

**TABLE 1** Blood analysis data on admission

Analyzed items	Value	Reference normal range (Lower limit-Upper limit)
White blood cell count ( $\mu\text{L}$ )	31,000	4300-8000
platelet count ( $\times 10^4/\mu\text{L}$ )	25.5	18.0-34.0
C-reactive protein (mg/dL)	40.27	0-0.4
Creatinine (mg/dL)	3.04	0.5-1.1
Blood urea nitrogen (mg/dL)	65	8-20
Creatine kinase (IU/L)	937	59-248
lactate dehydrogenase (IU/L)	505	124-222
Hemoglobin A1c (%)	8.0	4.6-6.2
Total bilirubin (mg/dL)	0.6	0.2-1.0
Fibrinogen (mg/dL)	953	200-400
Fibrin degradation products ( $\mu\text{g/mL}$ )	14.6	0-10.0
Antithrombin III (%)	65	70-120
Prothrombin time-international normalized ratio	1.17	0.90-1.10
Lactate level (mmol/L)	3.8	0.5-1.6

**FIGURE 1** Preoperative contrast-enhanced computed tomography image shows massive intrasoft tissue fluid collection from the neck to pectoralis major and minor muscles that extends to the axillary line, dorsally to the sternal notch (Yellow arrow)

At 4th day after admission, as Osler's node on the patient's left thumb and Janeway spot on the dorsal side of his left hand were detected (Figure 2), we performed the TTE again and found only mild mitral regurgitation. However, subsequent TEE performed the next day revealed a vegetation of  $12 \times 15$  mm in size at the border of the left ventricular posterior wall and posterior mitral leaflet annulus (Figure 3).

The site and size of the vegetation were clear indications for surgery, however as the patient had no signs of heart

failure or major thrombotic event, we decided to prioritize treatment of mediastinitis prior to the planned cardiac surgery. The original surgical wound was sutured on day 36, after treatment with interval irrigation and NPWT. The cerebral magnetic resonance image performed prior to surgery had no signs of mycotic aneurysm or infarctions.

The vegetectomy and additional mitral valvuloplasty were performed 40 days after it was detected and diagnosed as infective endocarditis.

The operation began with the patient in a left half-lateral position, and femoral arteriovenous cardiopulmonary bypass was established. We approached from the right 4th intercostal space with an 8-cm incision. After opening the left atrium via right-side left arteriotomy, we detected a huge vegetation between the mitral annulus and left ventricular posterior wall that surrounded the papillary muscle and its P1 segment of the posterior mitral valve leaflet (Figure 4). The vegetation was resected and the P1 was concomitantly repaired with quadrangular resection using interrupted sutures. Because mitral regurgitation was still present, we performed an annuloplasty with a 28-mm Physio-ring II (Edwards Life Sciences, USA) along with A1-P1 edge-to-edge suturing. The resected vegetation specimen measured  $12 \times 15$  mm, and the culture of the specimen revealed the same MSSA.

The patient was continued on an additional 3 weeks of antibiotic therapy and was discharged from hospital 20 days after the surgery without any complications.

### 3 | DISCUSSION


Infective endocarditis is a severely infectious disease with high rates of mortality and morbidity.<sup>5-7</sup> Although the recommended timing of surgery for IE is documented in guidelines for prevention and treatment of infective endocarditis,<sup>1,8</sup> we usually have to decide appropriate timing for surgery on the basis of each patient's characteristics, conditions, and comorbidities. The greater than 10 mm size and site of the vegetation were clear indications for urgent surgical vegetectomy,<sup>1,9</sup> however, it was difficult to proceed directly to surgical vegetectomy for three reasons. First was the timing of bleeding complications that were a contraindication for cardiac surgery. Second, since the infected site was concomitant with the mediastinitis, we wanted to prevent worsening of the mediastinitis after open-heart surgery. Third, the patient was in septic shock and also suffered acute kidney injury following the septic shock.

Generally, a right thoracotomy approach offers better exposure to the mitral apparatus in patients with a small left atrium, allowing easy repair or replacement of the mitral valve.<sup>10-12</sup> This advantage results in reducing the volume of blood transfused and the length of stay in the intensive care unit.<sup>11,12</sup>

**FIGURE 2** Photographs showing Osler's node on the patient's left thumb and Janeway spot on the dorsal side of the left hand found at admission (Yellow arrow)




**FIGURE 3** Images of the vegetation detected on transesophageal echocardiography. The vegetation was present at the border of the posterior wall of the left ventricle and surrounded the papillary muscle and chordae of the posterior mitral leaflet (Yellow arrow)



Several current papers discuss the sensitivity and specificity of TTE and TEE.<sup>13-15</sup> The sensitivity of TTE for native valve vegetation seems to be approximately 70%.<sup>15</sup> As we could not detect vegetation on TTE initially, therefore, we herein re-emphasize the necessity of performing TEE in patients highly suspected of having infective endocarditis and especially in those with positive blood cultures for *Staphylococcus*. The patient had no caries on his mouth and no current episode of dental treatment. The only cause we could determine for this soft tissue infection was brachial plexus block anesthesia performed by a local physician performed to control the patient's frozen shoulder syndrome.

#### 4 | CONCLUSION

We successfully treated a patient with infective endocarditis following severe mediastinitis. By considering appropriate timing for cardiac surgery and approaching via a right



**FIGURE 4** Image of the intraoperative findings. The huge vegetation was attached between the mitral annulus and left ventricular posterior wall that surrounded the papillary muscle and its P1 segment of the posterior mitral valve leaflet (Yellow arrow)

thoracotomy, vegetectomy, and mitral valvuloplasty were safely performed and the patient was discharged without any complications.

## ACKNOWLEDGMENTS

Published with written consent of the patient.

## CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. There was no external funding in the preparation of this manuscript.

## AUTHOR CONTRIBUTIONS

KU: involved in patient care, decision making of the treatment as well as the drafting, review, and revision of the initial manuscript. YT: involved in patient's care and decision making of the treatment. TS and YM: involved in patient care as well as the review and revision of the initial manuscript. All authors approved the final manuscript submission and agree to be accountable for all aspects of the study.

## ETHICAL APPROVAL

Written informed consent was appropriately obtained from the patient.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

## ORCID

Kenichiro Uchida  <https://orcid.org/0000-0002-3673-6007>

## REFERENCES

- Habib G, Lancellotti P, Antunes MJ, et al. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J*. 2015;36(44):3075-3128.
- Kiefer T, Park L, Tribouilloy C, et al. Association between valvular surgery and mortality among patients with infective endocarditis complicated by heart failure. *JAMA*. 2011;306:2239-2247.
- Thuny F, Beurtheret S, Mancini J, et al. The timing of surgery influences mortality and morbidity in adults with severe complicated infective endocarditis: a propensity analysis. *Eur Heart J*. 2011;32:2027-2033.
- Lalani T, Chu VH, Park LP, et al. In-hospital and 1-year mortality in patients undergoing early surgery for prosthetic valve endocarditis. *JAMA Intern Med*. 2013;173:1495-1504.
- Olmos C, Vilacosta I, Fernández C, et al. Contemporary epidemiology and prognosis of septic shock in infective endocarditis. *Eur Heart J*. 2013;34:1999-2006.
- García-Cabrera E, Fernández-Hidalgo N, Almirante B, et al. Neurological complications of infective endocarditis risk factors, outcome, and impact of cardiac surgery: a multicenter observational study. *Circulation*. 2013;127:2272-2284.
- Kang DH, Kim YJ, Kim SH, et al. Early surgery versus conventional treatment for infective endocarditis. *N Engl J Med*. 2012;366:2466-2473.
- Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC Focused Update of the 2014 AHA/ACC Guideline for the Management of Patients With Valvular Heart Disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2017;70:252-289.
- Wang D, Wang Q, Yang X, Wu Q, Li Q. Mitral valve replacement through a minimal right vertical infra-axillary thoracotomy versus standard median sternotomy. *Ann Thorac Surg*. 2009;87:704-708.
- Aklog L, Adams DH, Couper GS, Gobeze R, Sears S, Cohn LH. Techniques and results of direct-access minimally invasive mitral valve surgery: a paradigm for the future. *J Thorac Cardiovasc Surg*. 1998;116:705-715.
- Glower DD, Landolfo KP, Clements F, et al. Mitral valve operation via Port Access versus median sternotomy. *Eur J Cardiothorac Surg*. 1998;14(suppl 1):S143-S147.
- Suri RM, Antiel RM, Burkhart HM, et al. Quality of life after early mitral valve repair using conventional and robotic approaches. *Ann Thorac Surg*. 2012;93:761-769.
- Hill EE, Herijgers P, Claus P, Vanderschueren S, Peetermans WE, Herregods MC. Abscess in infective endocarditis: the value of transesophageal echocardiography and outcome: a 5-year study. *Am Heart J*. 2007;154:923-928.
- Habib G, Badano L, Tribouilloy C, Vilacosta I, Zamorano JL. Recommendations for the practice of echocardiography in infective endocarditis. *Eur J Echocardiogr*. 2010;11:202-219.
- Ansari A, Rigolin VH. Infective endocarditis: an update on the role of echocardiography. *Curr Cardiol Rep*. 2010;12:265-271.

**How to cite this article:** Uchida K, Takahashi Y, Shibata T, Mizobata Y. Successful surgical treatment approach for mitral valve vegetation of infective endocarditis after severe soft tissue infection with mediastinitis. *Clin Case Rep*. 2021;9:e04209. <https://doi.org/10.1002/ccr3.4209>