

Tris(2-benzamidoethyl)ammonium tetrafluoroborate

Marcy L. Pilate,^a Henry Blount,^a Frank R. Fronczek^b and Md. Alamgir Hossain^{a*}

^aDepartment of Chemistry and Biochemistry, Jackson State University, Jackson, MS 39217, USA, and ^bDepartment of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA

Correspondence e-mail: alamgir@chem.jsums.edu


Received 20 June 2010; accepted 22 June 2010

Key indicators: single-crystal X-ray study; $T = 90\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.048; wR factor = 0.114; data-to-parameter ratio = 16.2.

In the title compound, $\text{C}_{27}\text{H}_{31}\text{N}_4\text{O}_3^+\cdot\text{BF}_4^-$, the central N atom is protonated. The three arms form a pocket and one amidic O atom accepts an intermolecular hydrogen bond with the protonated amine. The tetrafluoroborate anion is outside the cavity and is hydrogen bonded to one amide N atom. Adjacent organic cations are connected by a pair of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a chain.

Related literature

For general background to tris(aminoethyl)-amine and its binding of anions, see: Bianchi *et al.* (1997); Kang *et al.* (2005); Hossain, (2008); For related structures, see: Bazzicalupi *et al.* (2009); Hossain *et al.* (2004); Burgess *et al.* (1991), Lo & Ng (2008); Saeed *et al.* (2010).



Experimental

Crystal data

$\text{C}_{27}\text{H}_{31}\text{N}_4\text{O}_3^+\cdot\text{BF}_4^-$
 $M_r = 546.37$
Monoclinic, $P2_1/n$
 $a = 13.325 (2)\text{ \AA}$
 $b = 9.572 (2)\text{ \AA}$
 $c = 21.118 (3)\text{ \AA}$
 $\beta = 94.546 (10)^\circ$

$V = 2685.1 (8)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$
 $T = 90\text{ K}$
 $0.27 \times 0.25 \times 0.10\text{ mm}$

Data collection

Nonius KappaCCD diffractometer
with Oxford Cryostream
31844 measured reflections
 $R_{\text{int}} = 0.068$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.114$
 $S = 1.00$
5914 reflections
365 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.26\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N \cdots O1	1.003 (19)	1.70 (2)	2.646 (2)	155.2 (16)
N2—H2N \cdots O3 ⁱ	0.87 (2)	2.01 (2)	2.861 (2)	168.8 (19)
N3—H3N \cdots F3	0.92 (2)	2.00 (2)	2.901 (2)	170.0 (18)
N4—H4N \cdots O2 ⁱⁱ	0.87 (2)	2.12 (2)	2.894 (2)	147.5 (19)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *COLLECT* (Nonius, 1999); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

This work was supported by National Institutes of Health, Division of National Center for Research Resources, under grant No. G12RR013459. This material is based upon work supported by the National Science Foundation under CHE-0821357. HB was supported by the National Institutes of Health (NIH) Science Education Partnership Award (SEPA) Program "Recovery Act Administrative Supplements Providing Summer Research Research Experiences for Students and Science Educators" under contract 5R25RR020405-04S1. Purchase of the diffractometer was made possible by grant No. LEQSF (1999–2000)–ENH-TR-13, administered by the Louisiana Board of Regents.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2214).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Molterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Bazzicalupi, C., Bencini, A., Bianchi, A., Danesi, A., Giorgi, C. & Valtancoli, B. (2009). *Inorg. Chem.* **48**, 2391–2398.
- Bianchi, A., García-España, E. & Bowman-James, K. (1997). *Supramolecular Chemistry of Anions*. New York: Wiley–VCH.
- Burgess, J., Al-Alousy, A., Fawcett, J. & Russell, D. R. (1991). *Acta Cryst.* **C47**, 2506–2508.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Hossain, M. A. (2008). *Curr. Org. Chem.* **12**, 1231–1256.
- Hossain, M. A., Liljegren, J. A., Powell, R. D. & Bowman-James, K. (2004). *Inorg. Chem.* **43**, 3751–3755.

organic compounds

- Kang, S. O., Hossain, M. A., Powell, D. & Bowman-James, K. (2005). *Chem. Commun.* pp. 328–330.
- Lo, K. M. & Ng, S. W. (2008). *Acta Cryst. E* **64**, o922.
- Nonius (1999). COLLECT. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Saeed, M. A., Fronczek, F. R. & Hossain, M. A. (2010). *Acta Cryst. E* **66**, o656–o657.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, o1833–o1834 [doi:10.1107/S1600536810024323]

Tris(2-benzamidoethyl)ammonium tetrafluoroborate

M. L. Pilate, H. Blount, F. R. Fronczek and M. A. Hossain

Comment

Tris(aminoethyl)–amine, trend with *C*3 symmetry is known to bind an anion (Burgess *et al.*, 1991; Hossain *et al.*, 2004; Bazzicalupi *et al.*, 2009) and is often used to build azacryptands with amine or amide functionality (Hossain, 2008). These molecules are of particularly interest for binding anions with three fold rotation axes, such as nitrate, phosphate, perchlorate and sulfate (Bianchi *et al.*, 1997). The binding is predominantly occurred through hydrogen bonding and electrostatic interactions of protonated amines or amidic protons with negatively charged species. For example, tetrahedral sulfate was seen to be encapsulated in amine cryptand or amide cryptand where the same trend units are as used as building blocks (Kang *et al.*, 2005). Herein, we report the molecular and crystal structures of the title compound in which a tetrafluoroborate anion is held outside the cavity with one hydrogen bond and remains outside the cavity. Single crystal analysis of the title compound reveals that the molecule crystallizes in its monoclinic space group forming a cavity with the three arms that are connected with one central amine (Fig. 1). The central amine is protonated and the charge is balanced with one tetrafluoroborate that is located outside the cavity and is bonded with one amide nitrogen. In the single-crystal, one amide H points toward the cavity and makes an intermolecular hydrogen bond with the proton on the central amine with N···O distance of 2.864 (2) Å (Table 1). Two adjacent molecules are connected each other forming a centrosymmetric dimer with two hydrogen bonds with N···O bonds distances of 2.861 (2) Å and 2.894 (2) Å (Fig. 2). Similar intramolecular bonding was previously reported for mono-functional (Lo & Ng, 2008) or di-functional (Saeed *et al.*, 2010).

Experimental

To a solution of tris(2-benzamidoethyl) amine (50 mg) in methanol (1 ml) was added a few drops fluoroboric acid at room temperature. The white powder formed was collected by filtration. Yield: 80%. M.P. 423 K. ^1H NMR (500 MHz, D_2O , *TSP*): δ 8.78 (b, 3H, NHCO), 7.81(d, 3H, ArH), 7.72 (b, 1H, NHCH₂), 7.56 (t, 3H, ArH), 7.47 (t, 3H, ArH), 3.72 (t, 6H, CH₂NHCO), 3.35 (t, 6H, NHCH₂). ^{13}C NMR (125 MHz, D_2O , *TSP*): δ 167.8 (CO), 138.9 (ArC), 131.9 (ArC), 128.8 (ArC), 127.6 (ArC), 52.4 (NHCH₂), 34.8 (CH₂CO). The salt was redissolved in water and crystals suitable for X-ray analysis were grown from slow evaporation of the solvent at room temperature.

Refinement

H atoms on C were placed in idealized positions with C—H distances 0.95–0.99 Å and thereafter treated as riding. The coordinates of those on N were refined. The U_{iso} for H was assigned as 1.2 times U_{eq} of the attached atom. The largest residual density peak was 1.50 Å from O2.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as a small spheres of arbitrary radius. Hydrogen bonds are presented by dashed lines.




Fig. 2. A view of the hydrogen bonded centrosymmetric dimer.

Tris(2-benzamidoethyl)ammonium tetrafluoroborate

Crystal data

$C_{27}H_{31}N_4O_3^+ \cdot BF_4^-$	$F(000) = 1144$
$M_r = 546.37$	$D_x = 1.352 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Melting point: 423 K
Hall symbol: -P 2yn	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 13.325 (2) \text{ \AA}$	Cell parameters from 6170 reflections
$b = 9.572 (2) \text{ \AA}$	$\theta = 2.5\text{--}27.1^\circ$
$c = 21.118 (3) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$\beta = 94.546 (10)^\circ$	$T = 90 \text{ K}$
$V = 2685.1 (8) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.27 \times 0.25 \times 0.10 \text{ mm}$

Data collection

Nonius KappaCCD	3310 reflections with $I > 2\sigma(I)$
diffractometer with Oxford Cryostream	
Radiation source: fine-focus sealed tube	
graphite	$R_{\text{int}} = 0.068$
ω -scans with κ -offsets	$\theta_{\text{max}} = 27.1^\circ, \theta_{\text{min}} = 2.6^\circ$
31844 measured reflections	$h = -17 \rightarrow 17$
5914 independent reflections	$k = -12 \rightarrow 12$
	$l = -27 \rightarrow 26$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.114$	$w = 1/[\sigma^2(F_o^2) + (0.0492P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\max} = 0.001$
5914 reflections	$\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$
365 parameters	$\Delta\rho_{\min} = -0.27 \text{ e \AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.0047 (7)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.69598 (10)	0.66708 (15)	0.51248 (6)	0.0235 (3)
O2	0.97004 (10)	0.65535 (14)	0.51217 (6)	0.0237 (3)
O3	0.66033 (11)	0.35955 (15)	0.42976 (6)	0.0269 (4)
N1	0.77743 (12)	0.52786 (18)	0.61191 (7)	0.0190 (4)
H1N	0.7663 (14)	0.584 (2)	0.5718 (9)	0.023*
N2	0.55317 (13)	0.65036 (19)	0.56356 (7)	0.0227 (4)
H2N	0.4881 (16)	0.655 (2)	0.5611 (9)	0.027*
N3	0.89693 (13)	0.78188 (18)	0.58685 (7)	0.0207 (4)
H3N	0.8721 (15)	0.867 (2)	0.5972 (9)	0.025*
N4	0.81500 (13)	0.39353 (18)	0.47972 (7)	0.0204 (4)
H4N	0.8780 (16)	0.412 (2)	0.4753 (9)	0.024*
C1	0.67293 (15)	0.4962 (2)	0.63108 (9)	0.0212 (5)
H1A	0.6782	0.4611	0.6753	0.025*
H1B	0.6432	0.4208	0.6035	0.025*
C2	0.60230 (16)	0.6219 (2)	0.62667 (8)	0.0238 (5)

supplementary materials

H2A	0.5495	0.6073	0.6564	0.029*
H2B	0.6412	0.7058	0.6411	0.029*
C3	0.60327 (16)	0.6812 (2)	0.51281 (9)	0.0208 (5)
C4	0.54443 (15)	0.7377 (2)	0.45532 (9)	0.0203 (5)
C5	0.46519 (16)	0.8306 (2)	0.45962 (9)	0.0243 (5)
H5	0.4442	0.8555	0.5001	0.029*
C6	0.41677 (16)	0.8870 (2)	0.40503 (10)	0.0293 (5)
H6	0.3632	0.9516	0.4081	0.035*
C7	0.44618 (16)	0.8496 (2)	0.34635 (10)	0.0310 (6)
H7	0.4121	0.8875	0.3090	0.037*
C8	0.52504 (17)	0.7572 (2)	0.34138 (10)	0.0312 (6)
H8	0.5447	0.7313	0.3008	0.037*
C9	0.57503 (16)	0.7029 (2)	0.39571 (9)	0.0269 (5)
H9	0.6305	0.6417	0.3924	0.032*
C10	0.83608 (15)	0.6116 (2)	0.66272 (9)	0.0218 (5)
H10A	0.8600	0.5483	0.6978	0.026*
H10B	0.7908	0.6814	0.6801	0.026*
C11	0.92603 (15)	0.6867 (2)	0.63859 (9)	0.0224 (5)
H11A	0.9612	0.7398	0.6740	0.027*
H11B	0.9737	0.6168	0.6239	0.027*
C12	0.92266 (15)	0.7599 (2)	0.52720 (9)	0.0197 (5)
C13	0.88929 (15)	0.8684 (2)	0.47918 (9)	0.0191 (5)
C14	0.80176 (16)	0.9456 (2)	0.48281 (9)	0.0237 (5)
H14	0.7618	0.9325	0.5176	0.028*
C15	0.77232 (17)	1.0423 (2)	0.43572 (9)	0.0270 (5)
H15	0.7120	1.0942	0.4382	0.032*
C16	0.83094 (17)	1.0627 (2)	0.38534 (9)	0.0267 (5)
H16	0.8112	1.1296	0.3535	0.032*
C17	0.91824 (17)	0.9861 (2)	0.38106 (9)	0.0272 (5)
H17	0.9583	1.0000	0.3463	0.033*
C18	0.94687 (16)	0.8890 (2)	0.42784 (9)	0.0231 (5)
H18	1.0065	0.8359	0.4248	0.028*
C19	0.83279 (16)	0.3959 (2)	0.59740 (8)	0.0211 (5)
H19A	0.8299	0.3309	0.6337	0.025*
H19B	0.9045	0.4192	0.5935	0.025*
C20	0.79230 (16)	0.3222 (2)	0.53764 (8)	0.0223 (5)
H20A	0.8210	0.2268	0.5374	0.027*
H20B	0.7184	0.3131	0.5381	0.027*
C21	0.74826 (15)	0.4008 (2)	0.42910 (9)	0.0199 (5)
C22	0.78649 (15)	0.4566 (2)	0.36920 (9)	0.0211 (5)
C23	0.86024 (15)	0.5588 (2)	0.36906 (9)	0.0228 (5)
H23	0.8870	0.5989	0.4080	0.027*
C24	0.89490 (16)	0.6025 (2)	0.31219 (9)	0.0280 (5)
H24	0.9440	0.6746	0.3122	0.034*
C25	0.85853 (18)	0.5420 (2)	0.25530 (9)	0.0308 (6)
H25	0.8844	0.5699	0.2166	0.037*
C26	0.78424 (18)	0.4403 (2)	0.25513 (10)	0.0338 (6)
H26	0.7589	0.3986	0.2163	0.041*
C27	0.74720 (17)	0.4001 (2)	0.31173 (9)	0.0274 (5)

H27	0.6945	0.3332	0.3114	0.033*
B1	0.8258 (2)	1.0373 (3)	0.69733 (11)	0.0262 (6)
F1	0.74417 (10)	0.95062 (13)	0.70473 (5)	0.0393 (4)
F2	0.81014 (10)	1.16644 (13)	0.72363 (5)	0.0396 (4)
F3	0.83701 (9)	1.05339 (12)	0.63193 (5)	0.0297 (3)
F4	0.91193 (10)	0.97574 (15)	0.72626 (5)	0.0444 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0192 (9)	0.0295 (9)	0.0217 (7)	0.0012 (7)	0.0004 (6)	0.0045 (6)
O2	0.0247 (9)	0.0212 (9)	0.0252 (8)	0.0040 (7)	0.0008 (6)	-0.0025 (6)
O3	0.0196 (9)	0.0279 (9)	0.0329 (8)	0.0000 (7)	0.0000 (6)	0.0030 (7)
N1	0.0182 (10)	0.0217 (10)	0.0169 (9)	-0.0006 (8)	-0.0003 (7)	0.0011 (7)
N2	0.0147 (10)	0.0320 (11)	0.0213 (9)	0.0018 (9)	0.0014 (8)	0.0021 (8)
N3	0.0235 (11)	0.0182 (10)	0.0201 (9)	0.0020 (8)	0.0010 (7)	0.0013 (8)
N4	0.0170 (10)	0.0250 (10)	0.0192 (9)	-0.0013 (9)	0.0013 (8)	-0.0006 (8)
C1	0.0182 (12)	0.0258 (13)	0.0199 (11)	-0.0029 (10)	0.0034 (8)	0.0007 (9)
C2	0.0230 (12)	0.0304 (13)	0.0181 (11)	0.0016 (10)	0.0027 (9)	0.0003 (9)
C3	0.0210 (13)	0.0188 (12)	0.0223 (11)	-0.0010 (10)	0.0007 (9)	-0.0017 (9)
C4	0.0183 (12)	0.0212 (12)	0.0209 (11)	-0.0012 (10)	-0.0010 (9)	0.0017 (9)
C5	0.0226 (13)	0.0241 (12)	0.0263 (12)	-0.0013 (10)	0.0036 (9)	0.0007 (10)
C6	0.0250 (13)	0.0293 (14)	0.0333 (13)	0.0044 (11)	0.0010 (10)	0.0069 (11)
C7	0.0247 (13)	0.0398 (15)	0.0275 (12)	0.0034 (12)	-0.0040 (10)	0.0123 (11)
C8	0.0321 (14)	0.0391 (15)	0.0224 (12)	0.0044 (12)	0.0024 (10)	0.0050 (10)
C9	0.0231 (13)	0.0292 (14)	0.0283 (12)	0.0022 (10)	0.0021 (9)	0.0023 (10)
C10	0.0237 (12)	0.0236 (12)	0.0173 (10)	-0.0001 (10)	-0.0028 (9)	-0.0019 (9)
C11	0.0212 (12)	0.0236 (12)	0.0218 (11)	0.0013 (10)	-0.0010 (9)	0.0010 (9)
C12	0.0172 (12)	0.0195 (12)	0.0221 (11)	-0.0051 (10)	-0.0012 (9)	-0.0019 (9)
C13	0.0185 (12)	0.0173 (12)	0.0212 (11)	-0.0019 (10)	-0.0004 (8)	-0.0035 (9)
C14	0.0274 (13)	0.0228 (12)	0.0206 (11)	-0.0009 (11)	0.0010 (9)	-0.0039 (10)
C15	0.0284 (13)	0.0226 (13)	0.0292 (12)	0.0035 (11)	-0.0024 (10)	-0.0026 (10)
C16	0.0325 (14)	0.0207 (12)	0.0253 (12)	-0.0028 (11)	-0.0065 (10)	0.0017 (10)
C17	0.0302 (14)	0.0286 (14)	0.0225 (12)	-0.0068 (11)	0.0009 (9)	-0.0002 (10)
C18	0.0206 (12)	0.0225 (12)	0.0261 (11)	-0.0021 (10)	0.0016 (9)	-0.0034 (10)
C19	0.0239 (12)	0.0181 (12)	0.0213 (11)	0.0051 (10)	0.0019 (9)	0.0012 (9)
C20	0.0248 (13)	0.0215 (12)	0.0210 (11)	0.0003 (10)	0.0048 (9)	0.0006 (9)
C21	0.0159 (12)	0.0168 (11)	0.0268 (12)	0.0024 (10)	0.0004 (9)	-0.0039 (9)
C22	0.0184 (12)	0.0216 (12)	0.0225 (11)	0.0055 (10)	-0.0029 (9)	0.0014 (9)
C23	0.0243 (13)	0.0236 (13)	0.0203 (11)	0.0040 (10)	0.0012 (9)	-0.0021 (9)
C24	0.0264 (13)	0.0264 (13)	0.0312 (13)	0.0020 (11)	0.0021 (10)	0.0031 (10)
C25	0.0428 (16)	0.0291 (14)	0.0209 (12)	0.0080 (12)	0.0050 (10)	0.0042 (10)
C26	0.0472 (16)	0.0296 (14)	0.0231 (12)	0.0057 (13)	-0.0079 (11)	0.0007 (10)
C27	0.0294 (14)	0.0230 (13)	0.0285 (12)	-0.0003 (11)	-0.0058 (10)	0.0036 (10)
B1	0.0297 (16)	0.0295 (16)	0.0199 (13)	0.0049 (13)	0.0043 (11)	0.0011 (11)
F1	0.0437 (9)	0.0353 (8)	0.0392 (7)	-0.0023 (7)	0.0056 (6)	0.0090 (6)
F2	0.0628 (10)	0.0305 (8)	0.0269 (7)	0.0042 (7)	0.0127 (6)	-0.0040 (6)
F3	0.0397 (8)	0.0307 (8)	0.0194 (6)	0.0108 (6)	0.0058 (5)	0.0022 (5)

supplementary materials

F4	0.0383 (9)	0.0703 (11)	0.0249 (7)	0.0223 (8)	0.0048 (6)	0.0119 (7)
----	------------	-------------	------------	------------	------------	------------

Geometric parameters (\AA , $^{\circ}$)

O1—C3	1.243 (2)	C11—H11A	0.9900
O2—C12	1.238 (2)	C11—H11B	0.9900
O3—C21	1.238 (2)	C12—C13	1.495 (3)
N1—C19	1.506 (2)	C13—C14	1.388 (3)
N1—C10	1.507 (2)	C13—C18	1.391 (3)
N1—C1	1.511 (2)	C14—C15	1.392 (3)
N1—H1N	1.003 (19)	C14—H14	0.9500
N2—C3	1.340 (2)	C15—C16	1.383 (3)
N2—C2	1.463 (2)	C15—H15	0.9500
N2—H2N	0.87 (2)	C16—C17	1.384 (3)
N3—C12	1.348 (2)	C16—H16	0.9500
N3—C11	1.452 (2)	C17—C18	1.388 (3)
N3—H3N	0.92 (2)	C17—H17	0.9500
N4—C21	1.337 (2)	C18—H18	0.9500
N4—C20	1.453 (2)	C19—C20	1.508 (3)
N4—H4N	0.87 (2)	C19—H19A	0.9900
C1—C2	1.526 (3)	C19—H19B	0.9900
C1—H1A	0.9900	C20—H20A	0.9900
C1—H1B	0.9900	C20—H20B	0.9900
C2—H2A	0.9900	C21—C22	1.499 (3)
C2—H2B	0.9900	C22—C23	1.387 (3)
C3—C4	1.493 (3)	C22—C27	1.392 (3)
C4—C5	1.389 (3)	C23—C24	1.385 (3)
C4—C9	1.394 (3)	C23—H23	0.9500
C5—C6	1.385 (3)	C24—C25	1.386 (3)
C5—H5	0.9500	C24—H24	0.9500
C6—C7	1.376 (3)	C25—C26	1.388 (3)
C6—H6	0.9500	C25—H25	0.9500
C7—C8	1.384 (3)	C26—C27	1.384 (3)
C7—H7	0.9500	C26—H26	0.9500
C8—C9	1.382 (3)	C27—H27	0.9500
C8—H8	0.9500	B1—F2	1.378 (3)
C9—H9	0.9500	B1—F1	1.387 (3)
C10—C11	1.520 (3)	B1—F4	1.388 (3)
C10—H10A	0.9900	B1—F3	1.409 (2)
C10—H10B	0.9900		
C19—N1—C10	110.86 (15)	O2—C12—N3	122.60 (18)
C19—N1—C1	111.27 (16)	O2—C12—C13	121.20 (17)
C10—N1—C1	110.66 (15)	N3—C12—C13	116.19 (18)
C19—N1—H1N	108.6 (11)	C14—C13—C18	119.04 (19)
C10—N1—H1N	110.5 (11)	C14—C13—C12	122.60 (17)
C1—N1—H1N	104.8 (11)	C18—C13—C12	118.33 (18)
C3—N2—C2	123.66 (18)	C13—C14—C15	120.34 (19)
C3—N2—H2N	120.1 (13)	C13—C14—H14	119.8
C2—N2—H2N	115.9 (13)	C15—C14—H14	119.8

C12—N3—C11	122.19 (18)	C16—C15—C14	119.9 (2)
C12—N3—H3N	119.1 (12)	C16—C15—H15	120.0
C11—N3—H3N	117.5 (12)	C14—C15—H15	120.0
C21—N4—C20	121.86 (18)	C15—C16—C17	120.3 (2)
C21—N4—H4N	119.3 (13)	C15—C16—H16	119.9
C20—N4—H4N	116.9 (13)	C17—C16—H16	119.9
N1—C1—C2	113.71 (17)	C16—C17—C18	119.6 (2)
N1—C1—H1A	108.8	C16—C17—H17	120.2
C2—C1—H1A	108.8	C18—C17—H17	120.2
N1—C1—H1B	108.8	C17—C18—C13	120.8 (2)
C2—C1—H1B	108.8	C17—C18—H18	119.6
H1A—C1—H1B	107.7	C13—C18—H18	119.6
N2—C2—C1	115.56 (16)	N1—C19—C20	114.47 (16)
N2—C2—H2A	108.4	N1—C19—H19A	108.6
C1—C2—H2A	108.4	C20—C19—H19A	108.6
N2—C2—H2B	108.4	N1—C19—H19B	108.6
C1—C2—H2B	108.4	C20—C19—H19B	108.6
H2A—C2—H2B	107.5	H19A—C19—H19B	107.6
O1—C3—N2	122.62 (18)	N4—C20—C19	113.57 (17)
O1—C3—C4	119.51 (17)	N4—C20—H20A	108.9
N2—C3—C4	117.86 (18)	C19—C20—H20A	108.9
C5—C4—C9	119.34 (18)	N4—C20—H20B	108.9
C5—C4—C3	122.10 (17)	C19—C20—H20B	108.9
C9—C4—C3	118.41 (19)	H20A—C20—H20B	107.7
C6—C5—C4	120.12 (19)	O3—C21—N4	122.99 (18)
C6—C5—H5	119.9	O3—C21—C22	120.58 (17)
C4—C5—H5	119.9	N4—C21—C22	116.38 (18)
C7—C6—C5	120.0 (2)	C23—C22—C27	119.33 (19)
C7—C6—H6	120.0	C23—C22—C21	122.77 (17)
C5—C6—H6	120.0	C27—C22—C21	117.88 (19)
C6—C7—C8	120.5 (2)	C24—C23—C22	119.99 (19)
C6—C7—H7	119.8	C24—C23—H23	120.0
C8—C7—H7	119.8	C22—C23—H23	120.0
C9—C8—C7	119.8 (2)	C23—C24—C25	120.5 (2)
C9—C8—H8	120.1	C23—C24—H24	119.8
C7—C8—H8	120.1	C25—C24—H24	119.8
C8—C9—C4	120.3 (2)	C24—C25—C26	119.7 (2)
C8—C9—H9	119.9	C24—C25—H25	120.1
C4—C9—H9	119.9	C26—C25—H25	120.1
N1—C10—C11	113.03 (15)	C27—C26—C25	119.7 (2)
N1—C10—H10A	109.0	C27—C26—H26	120.1
C11—C10—H10A	109.0	C25—C26—H26	120.1
N1—C10—H10B	109.0	C26—C27—C22	120.6 (2)
C11—C10—H10B	109.0	C26—C27—H27	119.7
H10A—C10—H10B	107.8	C22—C27—H27	119.7
N3—C11—C10	112.24 (16)	F2—B1—F1	110.24 (19)
N3—C11—H11A	109.2	F2—B1—F4	110.49 (19)
C10—C11—H11A	109.2	F1—B1—F4	108.96 (19)
N3—C11—H11B	109.2	F2—B1—F3	109.16 (19)

supplementary materials


C10—C11—H11B	109.2	F1—B1—F3	108.70 (18)
H11A—C11—H11B	107.9	F4—B1—F3	109.26 (18)
C19—N1—C1—C2	-162.24 (15)	N3—C12—C13—C18	151.42 (18)
C10—N1—C1—C2	74.0 (2)	C18—C13—C14—C15	-0.1 (3)
C3—N2—C2—C1	-61.3 (3)	C12—C13—C14—C15	-177.89 (18)
N1—C1—C2—N2	83.1 (2)	C13—C14—C15—C16	-0.7 (3)
C2—N2—C3—O1	11.6 (3)	C14—C15—C16—C17	0.8 (3)
C2—N2—C3—C4	-167.04 (19)	C15—C16—C17—C18	-0.3 (3)
O1—C3—C4—C5	-140.5 (2)	C16—C17—C18—C13	-0.5 (3)
N2—C3—C4—C5	38.1 (3)	C14—C13—C18—C17	0.6 (3)
O1—C3—C4—C9	35.0 (3)	C12—C13—C18—C17	178.56 (18)
N2—C3—C4—C9	-146.4 (2)	C10—N1—C19—C20	-166.83 (16)
C9—C4—C5—C6	0.5 (3)	C1—N1—C19—C20	69.6 (2)
C3—C4—C5—C6	175.96 (19)	C21—N4—C20—C19	-140.97 (19)
C4—C5—C6—C7	0.9 (3)	N1—C19—C20—N4	72.5 (2)
C5—C6—C7—C8	-1.0 (3)	C20—N4—C21—O3	6.9 (3)
C6—C7—C8—C9	-0.4 (4)	C20—N4—C21—C22	-170.42 (17)
C7—C8—C9—C4	1.9 (3)	O3—C21—C22—C23	149.1 (2)
C5—C4—C9—C8	-1.9 (3)	N4—C21—C22—C23	-33.5 (3)
C3—C4—C9—C8	-177.53 (19)	O3—C21—C22—C27	-32.3 (3)
C19—N1—C10—C11	75.0 (2)	N4—C21—C22—C27	145.04 (19)
C1—N1—C10—C11	-161.06 (17)	C27—C22—C23—C24	-0.8 (3)
C12—N3—C11—C10	-112.8 (2)	C21—C22—C23—C24	177.74 (19)
N1—C10—C11—N3	58.7 (2)	C22—C23—C24—C25	-1.9 (3)
C11—N3—C12—O2	2.2 (3)	C23—C24—C25—C26	2.3 (3)
C11—N3—C12—C13	-178.72 (17)	C24—C25—C26—C27	-0.1 (3)
O2—C12—C13—C14	148.4 (2)	C25—C26—C27—C22	-2.5 (3)
N3—C12—C13—C14	-30.7 (3)	C23—C22—C27—C26	3.0 (3)
O2—C12—C13—C18	-29.5 (3)	C21—C22—C27—C26	-175.61 (19)

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N1—H1N···O1	1.003 (19)	1.70 (2)	2.646 (2)	155.2 (16)
N2—H2N···O3 ⁱ	0.87 (2)	2.01 (2)	2.861 (2)	168.8 (19)
N3—H3N···F3	0.92 (2)	2.00 (2)	2.901 (2)	170.0 (18)
N4—H4N···O2 ⁱⁱ	0.87 (2)	2.12 (2)	2.894 (2)	147.5 (19)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

