

## CORRECTION

# Correction: Care management intervention to strengthen self-care of multimorbid patients with type 2 diabetes in a German primary care network: A randomized controlled trial

**Dominik Ose, Martina Kamradt, Marion Kiel, Tobias Freund, Werner Besier, Manfred Mayer, Johannes Krisam, Michel Wensing, Hans-Joachim Salize, Joachim Szecsenyi**

There is an error in [Table 1](#). The letter “c” in the “Number of comorbidities” row should not be capitalized. Please see the correct [Table 1](#) here.



## OPEN ACCESS

**Citation:** Ose D, Kamradt M, Kiel M, Freund T, Besier W, Mayer M, et al. (2019) Correction: Care management intervention to strengthen self-care of multimorbid patients with type 2 diabetes in a German primary care network: A randomized controlled trial. PLoS ONE 14(8): e0221229. <https://doi.org/10.1371/journal.pone.0221229>

**Published:** August 9, 2019

**Copyright:** © 2019 Ose et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 1. Patient characteristics—Comparison between groups at baseline.

	Intervention	N	Control	N
Number		252		243
Age years (SD)	68.40 (11.33)	249	68.31 (10.80)	238
Female	47.22%	119	48.56%	118
Married / cohabited	58.73%	148	53.09%	129
Up to 9 years in school	30.16%	76	26.75%	65
Professional education*	66.67%	168	67.07%	163
Diabetes duration in years (SD)	14.13 (9.50)	187	15.06 (12.07)	173
Treatment with insulin	33.7%	85	33.3%	81
Enrolled in an additional DMP <sup>§</sup>	23.8%	60	28.0%	68
Number of comorbidities	3.79 (1.16)	252	3.74 (1.22)	243
Mobility restrictions	18.7%	47	17.3%	42
Hospital stays per patients <sup>#</sup>	0.19 (0.51)	252	0.23 (0.61)	243
Systolic blood pressure (mmHg)	135.45 (14.65)	252	133.88 (14.25)	243
Diastolic blood pressure (mmHg)	80.27 (8.84)	252	79.63 (8.59)	243
Blood glucose (mg/dl)	148.11 (47.56)	252	151.76 (50.13)	243
Blood glucose (HbA1C)	7.13 (1.23)	252	7.25 (1.19)	243
BMI calculated	31.52 (6.73)	252	31.62 (5.97)	243

\*minimum vocational training;

<sup>§</sup>Asthma, CHD, COPD;

<sup>#</sup>last 9 month;

<https://doi.org/10.1371/journal.pone.0221229.t001>

There is an error in Table 3. The subheadings “T0” and “T1” are missing from the “Control group” section. Please see the correct Table 3 here.

Table 3. Description of SDSCA scores within the intervention and control group alongside p-values for differences in scores over time.

SDSCA-G	Intervention							Control group						
	T0			T1			P**	T0			T1			P**
	N	Mean	SD	N	Mean	SD		N	Mean	SD	N	Mean	SD	
sum	186	3.31	1.11	181	3.63	1.22	0.012	182	3.50	1.23	164	3.58	1.22	0.197
diet	208	4.30	1.58	199	4.54	1.50	0.180	201	4.38	1.76	186	4.40	1.72	0.959
exercise	231	3.13	1.84	209	3.12	1.83	0.876	220	3.01	1.85	194	3.10	2.01	0.731
BST*	78	5.48	2.28	73	5.58	2.01	0.781	75	5.45	2.26	68	5.93	1.90	0.051
foot care	222	2.47	2.19	211	2.85	2.28	0.207	223	3.06	2.25	195	3.15	2.46	0.322

\*blood sugar test;

\*\*p-value for testing whether least square means are equal to 0, based on linear mixed model with repeated measurements

<https://doi.org/10.1371/journal.pone.0221229.t002>

There is an error in reference 32. The correct reference is: Sheaff R, Gené-Badia J, Marshall M, S`vab I. The evolving public-private mix. In: Saltman RB, Rico A, Boerma W, Hrsg. Primary care in the driver's seat? Organizational reform in European primary care. New York City, NY: Open University Press; 2006. S. 129–46.

Within references 1, 5, 15, 20, 26, 29, 31, 34, 40, 47, 49, and 54, the “View Article” links do not lead to the cited articles and some PMIDs are not listed. Please view the complete references here.

1. Korff M von, Gruman J, Schaefer J, Curry SJ, Wagner EH. Collaborative management of chronic illness. *Ann Intern Med* 1997; 127(12):1097–102. <https://doi.org/10.7326/0003-4819-127-12-199712150-00008> PMID:9412313
5. Busse R. Disease management programs in Germany's statutory health insurance system. *Health Aff (Millwood)* 2004; 23(3):56–67. <https://doi.org/10.1377/hlthaff.23.3.56> PMID:15160803
15. Vogeli C, Shields AE, Lee TA, Gibson TB, Marder WD, Weiss KB et al. Multiple chronic conditions: prevalence, health consequences, and implications for quality, care management, and costs. *J Gen Intern Med* 2007; 22 Suppl 3:391–5. <https://doi.org/10.1007/s11606-007-0322-1> PMID:18026807
20. Wagner EH, Austin BT, Davis C, Hindmarsh M, Schaefer J, Bonomi A. Improving chronic illness care: translating evidence into action. *Health Aff (Millwood)* 2001; 20(6):64–78. <https://doi.org/10.1377/hlthaff.20.6.64> PMID:11816692
26. Gensichen J, Korff M von, Peitz M, Muth C, Beyer M, Guthlin C et al. Case management for depression by health care assistants in small primary care practices: a cluster randomized trial. *Ann Intern Med* 2009; 151(6):369–78. <https://doi.org/10.7326/0003-4819-151-6-200909150-00001> PMID:19755362
29. Kamradt M, Bozorgmehr K, Krisam J, Freund T, Kiel M, Qreini M et al. Assessing self-management in patients with diabetes mellitus type 2 in Germany: validation of a German version of the Summary of Diabetes Self-Care Activities measure (SDSCA-G). *Health Qual Life Outcomes* 2014; 12(1):185. <https://doi.org/10.1186/s12955-014-0185-1> PMID:25519204
31. Amelung V, Hildebrandt H, Wolf S. Integrated care in Germany—a stony but necessary road! *Int J Integr Care* 2012; 12:e16. <https://doi.org/10.5334/ijic.853> PMID:22977429
34. Eisenstat SA, Chang Y, Porneala BC, Geagan E, Wilkins G, Chase B et al. Development and Implementation of a Collaborative Team Care Model for Effective Insulin Use in an Academic Medical Center Primary Care Network. *Am J Med Qual* 2016. <https://doi.org/10.1177/1062860616651715> PMID:27259871
40. Lee H, Ahn S, Kim Y. Self-care, Self-efficacy, and Glycemic Control of Koreans With Diabetes Mellitus. *Asian Nurs Res (Korean Soc Nurs Sci)* 2009; 3(3):139–46. [https://doi.org/10.1016/S1976-1317\(09\)60025-6](https://doi.org/10.1016/S1976-1317(09)60025-6) PMID:25030472
47. Collins AE, Niles BL, Mori DL, Silberbogen AK, Seligowski AV. A telephone-based intervention to promote diabetes management in veterans with posttraumatic stress symptoms. *Prof Psychol Res Pr* 2014; 45(1):20–6. <https://doi.org/10.1037/a0032604>
49. Sarayani A, Mashayekhi M, Nosrati M, Jahangard-Rafsanjani Z, Javadi M, Saadat N et al. Efficacy of a telephone-based intervention among patients with type-2 diabetes; a randomized controlled trial in pharmacy practice. *Int J Clin Pharm* 2018. <https://doi.org/10.1007/s11096-018-0593-0> PMID:29435911
54. Ose D, Miksch A, Urban E, Natanzon I, Szecsenyi J, Kunz CU et al. Health related quality of life and comorbidity. A descriptive analysis comparing EQ-5D dimensions of patients in the German disease management program for type 2 diabetes and patients in routine care. *BMC Health Serv Res* 2011; 11(1):179. <https://doi.org/10.1186/1472-6963-11-179> PMID:21810241

## Reference

1. Ose D, Kamradt M, Kiel M, Freund T, Besier W, Mayer M, et al. (2019) Care management intervention to strengthen self-care of multimorbid patients with type 2 diabetes in a German primary care network: A randomized controlled trial. PLoS ONE 14(6): e0214056. <https://doi.org/10.1371/journal.pone.0214056> PMID: [31188825](https://pubmed.ncbi.nlm.nih.gov/31188825/)