

POSTER PRESENTATION

Open Access

# Genetic Variation in Intercellular Adhesion Molecule-1 (*ICAM-1*): candidate gene in susceptibility to malaria in the Indian population

Anuroopa Gupta\*, Harish Padh

From International Conference on Human Genetics and 39th Annual Meeting of the Indian Society of Human Genetics (ISHG)  
Ahmadabad, India. 23-25 January 2013

## Background

*Plasmodium falciparum* malaria is the most severe form of malaria causing morbidity and mortality worldwide. The key event in *P. falciparum* pathogenesis is the sequestration of the infected erythrocytes to the capillary endothelium, mediated by the receptor, *Plasmodium falciparum* erythrocyte membrane protein-1 (PfEMP1). PfEMP1 has distinct characteristics of binding to varied host ligands including Inter Cellular Adhesion Molecule (*ICAM-1*). Polymorphisms in the gene encoding *ICAM-1* influence the rate of progression or confer protection from or susceptibility to malaria parasite infection. The objective of our study was to evaluate the frequency of *ICAM-1* SNP in exon 6 (rs5498, K469E) in the Indian population and compare it with the global population influenced by the parasitic infection.

## Materials and methods

The genotyping for *ICAM-1* SNP rs5498 was performed by Restriction Fragment Length Polymorphism (RFLP) PCR. Analysis of *ICAM-1* variants was performed in 200 healthy subjects.

## Results

The allelic frequency for the variant *ICAM1*-Aallele was 59% in healthy control subjects. Chi square analysis reveals that it does not follow the Hardy Weinberg Equilibrium. Comparison with literature data indicates that frequency distribution in Indian population is similar to German, Swedish and Danish populations while it is significantly different from Finnish and Japanese

populations. The genotypic frequency of AA was 35%, AG was 49% and GG was 16% in healthy control subjects.

## Conclusions

The frequency data from the above study will help in understanding the role of genetic variations in *ICAM-1* adhesion molecule in the *falciparum* malaria pathogenesis in the Indian population. Upon comparison, a noteworthy difference in the genotype frequency of Nigerian children ( $p < 0.001$ ) was evinced. This data can be useful for predicting the potential risk of malaria in world population.

Published: 21 January 2014

doi:10.1186/1755-8166-7-S1-P106

**Cite this article as:** Gupta and Padh: Genetic Variation in Intercellular Adhesion Molecule-1 (*ICAM-1*): candidate gene in susceptibility to malaria in the Indian population. *Molecular Cytogenetics* 2014 **7**(Suppl 1): P106.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



Department of Cell and Molecular Biology, B.V. Patel Pharmaceutical Education and Research Development (PERD) Centre, Ahmedabad-380 054, Gujarat, India



© 2014 Gupta and Padh; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.