



Artesunate potentiates antibiotics by inactivating heme-harboured bacterial nitric oxide synthase and catalase

Zeng *et al.*

SHORT REPORT

Open Access

Artesunate potentiates antibiotics by inactivating heme-harboured bacterial nitric oxide synthase and catalase

Qing-Ping Zeng^{1*}, Na Xiao¹, Pei Wu¹, Xue-Qin Yang², Li-Xiang Zeng¹, Xiao-Xia Guo¹, Ping-Zu Zhang¹ and Frank Qiu³

Abstract

Background: A current challenge of coping with bacterial infection is that bacterial pathogens are becoming less susceptible to or more tolerant of commonly used antibiotics. It is urgent to work out a practical solution to combat the multidrug resistant bacterial pathogens.

Findings: Oxidative stress-acclimatized bacteria thrive in rifampicin by generating antibiotic-detoxifying nitric oxide (NO), which can be repressed by artesunate or an inhibitor of nitric oxide synthase (NOS). Suppressed bacterial proliferation correlates with mitigated NO production upon the combined treatment of bacteria by artesunate with antibiotics. Detection of the heme-artesunate conjugate and accordingly declined activities of heme-harboured bacterial NOS and catalase indicates that artesunate renders bacteria susceptible to antibiotics by alkylating the prosthetic heme group of hemo-enzymes.

Conclusions: By compromising NO-mediated protection from antibiotics and triggering harmful hydrogen peroxide burst, artesunate may serve as a promising antibiotic synergist for killing the multidrug resistant pathogenic bacteria.

Findings

Artesunate is a semi-synthetic soluble derivative of artemisinin, a plant sesquiterpene endoperoxide lactone with pleiotropic functions of anti-malaria, anti-tumour and anti-inflammation [1]. Here, we report that artesunate can also exert an accelerated anti-bacterial activity in combination with antibiotics. Initially, we found that artesunate mitigated the oxidative stress-induced generation of nitric oxide (NO) from the Gram-positive bacterium, *Bacillus licheniformis*. A gradual increase of nitrate/nitrite, the oxidation product of NO, was detected in the bacterial culture standing without agitation either at room temperature (the hypoxia group) or in a 4°C refrigerator (the hypoxia + cold group) although no significant difference was observed between the hypoxia + cold group and the hypoxia group (Figure 1). In similar, it has been demonstrated that NO is not


consumed and accumulates in the microenvironment of human tissue at lower oxygen concentrations [2]. It is also manifested from Figure 1 that oxidative stress-triggered NO generation was repressed by either artesunate or a specific inhibitor of nitric oxide synthase (NOS), NG-monomethyl-L-arginine monoacetate (L-NMMA), indicating that artesunate may represent a novel NOS inhibitor that acts in an unknown manner.

Knowing that bacterial NO can detoxify antibiotics through direct or indirect mechanisms [3], we tested whether NO that was triggered by acclimatization to hypoxia confers *B. licheniformis* tolerance to the antibiotic rifampicin. Consequently, when bacteria propagated to a logarithmic phase in the absence of rifampicin, acclimatized bacteria started to proliferate at a higher rate than non-acclimatized bacteria in the presence of rifampicin (Figure 2a). Similarly, acclimatized bacteria also showed somewhat accelerated proliferation in the presence of either cefotaxime (Figure 2b) or ampicillin (Figure 2c).

* Correspondence: qpzeng@gzhtcm.edu.cn

¹Tropical Medicine Institute, Guangzhou University of Chinese Medicine, Guangzhou, China


Full list of author information is available at the end of the article



Due to attenuating protective NO production in bacteria, artesunate was anticipated to reverse NO-mediated protection of bacteria from antibiotics. Indeed, artesunate in combination with rifampicin led to more stringent growth inhibition of *B. licheniformis* than rifampicin alone. As illustrated in Figure 3a, much slower growth of bacteria was monitored in artesunate + rifampicin than in rifampicin, hence addressing that artesunate potentiates rifampicin perhaps by attenuating NO production from bacteria. In the present study, we observed that enhanced NO burst occurred in bacteria that were exposed to rifampicin, whereas less NO was measured in bacteria that were co-treated by artesunate with rifampicin (Figure 3b).

In the NOS-free Gram-negative bacterium, *Escherichia coli*, a complex compassing nitrite reductase, flavohemoglobin and a NO-sensing regulator is harnessed to form NO [4]. Indeed, we detected the generation of NO in *E. coli*, either in the presence or absence of cefotaxime, but no correlation of bacterial proliferation with NO production was established (Figure 4a and 4b). From the extremely low level of NO in *E. coli*, we inferred that such a trace amount of NO might insufficient to sensitize cefotaxime.

The non-pathogenic *Bacillus subtilis* and pathogenic *Bacillus anthracis* as well as many other Gram-positive bacteria generate NO by their own NOS that associates with a prosthetic heme group for the reduction of ferric heme [Fe^{3+}] to ferrous heme [Fe^{2+}] [5]. Given that heme alkylation by artemisinin has been verified by identifying






Figure 3 Bacterial propagation and NO generation in the presence of antibiotics or artesunate + antibiotics for *B. licheniformis*. (a) Propagation of *B. licheniformis* in the presence of rifampicin or artesunate + rifampicin. (b) Generation of NO from *B. licheniformis* in the presence of rifampicin or artesunate + rifampicin. Rif: rifampicin; ART: artesunate.




Figure 4 Bacterial propagation and NO generation in the presence of antibiotics or artesunate + antibiotics for *E. coli*. (a) Propagation of *E. coli* in the presence of cefotaxime or artesunate + cefotaxime. (b) Generation of NO from *E. coli* in the presence of cefotaxime or artesunate + cefotaxime. Cef: cefotaxime; ART: artesunate.

the heme-artemisinin adduct in malaria-infected mice [6], and the interaction between heme and artesunate has been also recognised by monitoring the dynamic shift of one peak specific to heme and another peak unique to the heme-artesunate conjugate in tumour cells [7], we assumed that artesunate would also bind to the heme group of bacterial NOS and hence prohibit the inter-conversion of Fe^{3+} with Fe^{2+} within the hemoprotein. Indeed, A_{415} that reflects the absorbance of heme and A_{476} that represents the absorbance of the heme-artesunate conjugate reached high values after the bacterial culture was supplemented with artesunate for 3 h (Figure 5), suggesting that the A_{415} peak was likely resulted from the increase of NOS, while the A_{476} peak was probably derived from the conjugation of heme with artesunate. Nevertheless, the increase of NOS needs the overexpression of a corresponding gene that encodes NOS in *B. licheniformis*, whether NOS gene is inducible by artesunate-bound inactivation of NOS awaits further elucidation.

Until recently, there has no documented evidence regarding the impact of artesunate on bacterial catalase. Considering that catalase is also a heme-harboring enzyme among *Bacillus* [8], it can be deduced that a similar mechanism exists in *B. licheniformis*, by which artesunate binds to the prosthetic heme group of catalase and abrogates the conversion of hydrogen peroxide. Therefore, we further measured catalase activity after incubating bacteria with rifampicin or rifampicin + artesunate. The results showed that enzyme activity was considerably reduced once artesunate was included in the culture (Figure 6). In other observations, NO activates catalase in bacteria by diminishing the rate of cystine reduction to cysteine, which drives the Fenton




Figure 5 Time-course monitoring of the heme-artesunate conjugate in *B. licheniformis* incubated with artesunate. ART: artesunate. A_{415} represents the absorbance of heme at the wavelength of 415 nm; A_{476} represents the absorbance of the heme-artesunate conjugate at the wavelength of 476 nm.




Figure 6 Determination of catalase activity in *B. licheniformis* upon exposure to rifampicin or artesunate + rifampicin. ART: artesunate; Rif: rifampicin. A single asterisk (*) represents significant difference of the artesunate + rifampicin group from the rifampicin group ($P < 0.05$); Double asterisks (**) represent very significant difference of the artesunate + rifampicin group from the rifampicin group ($P < 0.01$).

reaction and simultaneously inhibits catalase [9]. Due to catalase inhibition, hydroxyl radicals that were derived from excess hydrogen peroxide exhibit the extreme toxicity to bacterial DNA through base modifications and strand breaks [10]. It can be concluded that artesunate facilitates the action of antibiotics against bacteria by synchronously inactivating NOS and catalase. Interestingly, artesunate also covalently binds to hemoproteins of tumour cells and potentiates the cytotoxicity of 5-fluorouracil *in vitro* and *in vivo* [11], implying that a common mechanism exists to interfere with the fate of bacterial and tumour cells. However, inhibition of a bacterial multidrug efflux pump system should represent an alternative manner of artesunate in sensitizing antibiotics [12].

A current challenge of coping with bacterial infection is that bacterial pathogens are becoming less susceptible to or more tolerant of commonly used antibiotics. For example, globally endemic tuberculosis caused by multidrug resistant strains of *Mycobacterium tuberculosis* remains a formidable threat to human health. Although the development of more potent antibiotics may prohibit the lethal pathogens from worldwide transmission, it has proved to be costly, time-consuming and technically difficult [13]. Otherwise, if an antibiotic synergist like artesunate can benefit to fight against the antibiotic resistant bacteria, it will minimise the dosage of antibiotics in antibacterial therapy and diminish the heavy incidence and rapid transmission of multidrug resistant bacterial pathogens.

Methods

Bacterial culture and NO content estimation

A single colony of *B. licheniformis* BL20386 or *E. coli* DH5 α was inoculated in LB broth and cultured overnight at 37°C. A 1% aliquot of pre-cultured bacteria was inoculated in LB broth and cultured at 37°C until absorbance at 600 nm (A_{600}) to 0.6. For oxidative stress acclimatization, triangle bottles with the bacterial culture were placed without agitation for 3 d at room temperature (the hypoxia group) or in a 4°C refrigerator (the hypoxia + cold group). For antibiotic exposure, a 1% aliquot of the overnight bacterial culture was inoculated in LB broth supplementing with different concentrations of rifampicin, cefotaxime, or ampicillin. For inhibitor treatment, a 1% aliquot of the overnight bacterial culture was inoculated in LB broth supplementing with 60 μ g/ml artesunate or 1 mM NG-monomethyl-L-arginine monoacetate (L-NMMA). On each day, 1 ml of the bacterial culture was taken out for estimation of NO content using a commercially available kit and according to the manufacturer's instruction. The content of NO is represented by the amount of nitrate (NO_3^-)/nitrite (NO_2^-). For plotting a standard curve of nitrate/nitrite, 1 M NaNO_3 was dissolved to a series of dilutions (1, 2, 5, 10, 20, 30, 40, 60 μ M) by LB broth for measurement of A_{540} , from which a regression equation and a determinant coefficient were calculated.

Bacterial growth assay

For monitoring the growth rate of oxidative stress-acclimatized bacteria, a 1% aliquot of the bacterial culture standing at room temperature for 3 d or that without oxidative stress treatment was inoculated in LB broth supplementing with rifampicin and cultured at 37°C with agitation. After cultured for 3, 6, 9 and 12 h, 1 ml of the bacterial culture was taken out and diluted with 9 ml of fresh LB broth to measure A_{600} for plotting the growth rate curve. For assaying the bacterial growth rate following treatment by rifampicin or artesunate + rifampicin, a 1% aliquot of the overnight bacterial culture was inoculated in LB broth supplementing with 60 μ g/ml artesunate and different concentration of rifampicin, and cultured at 37°C. After cultured for 3, 6, 9, 12 and 24 h, 1 ml of the bacterial culture was diluted with 9 ml of fresh LB broth to measure A_{600} for plotting the growth rate curve.

Detection of heme and heme-artesunate conjugate

A 1% overnight culture of *B. licheniformis* BL20386 was inoculated in LB broth supplementing with 60 μ g/ml artesunate and cultured at 37°C for 2 d. The triangle bottle with a pre-culture was first placed at ambient temperature for 1 d and subsequently at a 4°C refrigerator for 1 d. After supplementing with fresh LB broth

and 60 μ g/ml artesunate, the bacterial strain was continuously cultured overnight at 37°C. The culture was taken out on 3, 6 and 9 h for collecting cells by centrifugation and lysing them through repeated cycles of frozen-thaw, and finally the lysate was applied to measure A_{415} and A_{476} .

Determination of catalase activity

A 1% overnight culture of *B. licheniformis* BL20386 was inoculated in LB broth supplementing with 60 μ g/ml artesunate in the treatment group or without artesunate in the control group, and cultured at 37°C for 2 d. The triangle bottle with a pre-culture was first placed at ambient temperature for 1 d and subsequently at a 4°C refrigerator for 1 d. After supplementing with fresh LB broth containing 60 μ g/ml artesunate or 60 μ g/ml artesunate + different concentrations of rifampicin, the bacterial strain was continuously cultured overnight at 37°C. The culture was taken out from each group on 9, 12 and 24 h for measurement of A_{405} using a commercially available kit. The activity of catalase (U/ml) was calculated according the formula of $(A_{405} \text{ control group} - A_{405} \text{ treatment group}) \times 271 \times 1/6$.

Acknowledgements

This work was supported by Natural Science Foundation of Guangdong China (No. 9145624536-4000003) and Scientific Development Project of Guangdong China (No. 2007B031404008).

Author details

¹Tropical Medicine Institute, Guangzhou University of Chinese Medicine, Guangzhou, China. ²Artemisinin Research Centre, Guangzhou University of Chinese Medicine, Guangzhou, China. ³Simplex Biotechnologies, LLC, Clinton, NJ08809, USA.

Authors' contributions

NX, PW and XQY were responsible for bacterial culture and drug evaluation. LXZ and XXG carried out the spectrophotometric assay. PZZ performed the statistical analysis. QPZ designed and coordinated the experiment. FQ helped to draft the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 20 February 2011 Accepted: 30 June 2011

Published: 30 June 2011

References

1. Krishna S, Bustamante L, Haynes RK, Staines HM: **Artemisinins: their growing importance in medicine.** *Trends Pharmacol Sci* 2008, **29**:520-527.
2. Taylor CT, Moncada S: **Nitric oxide, cytochrome C oxidase, and the cellular response to hypoxia.** *Arterioscler Thromb Vasc Biol* 2010, **30**:643-647.
3. Gusarov I, Shatalin K, Starodubtseva M, Nudler E: **Endogenous nitric oxide protects bacteria against a wide spectrum of antibiotics.** *Science* 2009, **325**:1380-1384.
4. Corker H, Poole RK: **Nitric oxide formation by *Escherichia coli*.** *J Biol Chem* 2003, **278**:31584-31592.
5. Gusarov I, Starodubtseva M, Wang ZQ, McQuade L, Lippard SJ, Stuehr DJ, Nudler E: **Bacterial nitric-oxide synthases operate without a dedicated redox partner.** *J Biol Chem* 2008, **283**:13140-13147.

6. Robert A, Benoit-Vical F, Claparols C, Meunier B: **The antimalarial drug artemisinin alkylates heme in infected mice.** *Proc Natl Acad Sci USA* 2005, **102**:13676-13680.
7. Zhang SM, Gerhard GS: **Heme mediates cytotoxicity from artemisinin and serves as a general anti-proliferation target.** *PLoS ONE* 2009, **4**:e7472.
8. Bol DK, Yasbin RE: **Analysis of the dual regulatory mechanisms controlling expression of the vegetative catalase gene of *Bacillus subtilis*.** *J Bacteriol* 1994, **176**:6744-6748.
9. Gusarov I, Nudler E: **NO-mediated cytoprotection: instant adaptation to oxidative stress in bacteria.** *Proc Natl Acad Sci USA* 2005, **102**:13855-13860.
10. Woodmansee AN, Imlay JA: **Reduced flavins promote oxidative DNA damage in non-respiring *Escherichia coli* by delivering electrons to intracellular free iron.** *J Biol Chem* 2002, **277**:34055-34066.
11. Zeng QP, Zhang PZ: **Artesunate mitigates proliferation of tumor cells by alkylating heme-harboring nitric oxide synthase.** *Nitric Oxide* 2011, **24**:110-112.
12. Li B, Yao Q, Pan XC, Wang N, Zhang R, Li J, Ding G, Liu X, Wu C, Ran D, Zheng J, Zhou H: **Artesunate enhances the antibacterial effect of β -lactam antibiotics against *Escherichia coli* by increasing antibiotic accumulation via inhibition of the multidrug efflux pump system AcrAB-TolC.** *J Antimicrob Chemother* 2011, **66**:769-777.
13. Patel BA, Crane B: **When it comes to antibiotics, bacteria show some NO-how.** *J Mol Cell Biol* 2010, **2**:234-236.

doi:10.1186/1756-0500-4-223

Cite this article as: Zeng et al.: Artesunate potentiates antibiotics by inactivating heme-harboring bacterial nitric oxide synthase and catalase. *BMC Research Notes* 2011 **4**:223.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

