

A Surprising Cause of Contained Aortic Rupture: Perforation from a Lumbar Osteophyte

Nikolaos Kontopodis, MD, MSc, PhD¹ Christos V. Ioannou, MD, PhD¹

¹Vascular Surgery Unit, Department of Cardiothoracic and Vascular Surgery, University Hospital of Heraklion, Heraklion, Greece

Address for correspondence Nikolaos Kontopodis, MD, PhD, Vascular Surgery Unit, Department of Cardiothoracic and Vascular Surgery, University Hospital of Heraklion, PO Box 1352, 711 10, Heraklion, Greece (e-mail: kontopodisn@yahoo.gr).

AORTA 2020;8:184–186.

Abstract

Keywords

- ▶ contained aortic rupture
- ▶ pseudoaneurysm
- ▶ osteophyte

We describe a patient with contained aortic rupture due to perforation from a protruding lumbar osteophyte, who was treated by open surgery. This case underlines that less common aortic pathologies are possible, which require a high suspicion index to be diagnosed.

Degenerative abdominal aortic aneurysm (AAA) is the most common type of AAA. Infection, dissection, and trauma are much less frequent causes of AAA development. Traumatic AAAs are in fact contained ruptures due to aortic perforation resulting in local containment of hemorrhage rather than uncontrolled bleeding. We describe a patient who presented with a traumatic aortic pseudoaneurysm due to perforation from a lumbar osteophyte.

A 75-year-old male patient was referred because of an incidentally discovered 5-cm AAA, which was identified on




Fig. 1 Axial computed tomography image. AAA, abdominal aortic aneurysm.




Fig. 2 Intraoperative appearance of the lesion.

received
January 24, 2020
accepted
June 20, 2020
published online
March 24, 2021

DOI <https://doi.org/10.1055/s-0040-1715468>.
ISSN 2325-4637.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical Publishers, Inc., 333 Seventh Avenue, 18th Floor, New York, NY 10001, USA




Fig. 3 Intraoperative photo after incising the infrarenal aorta, showing the protrusion of the osteophyte into the posterior aortic wall.




Fig. 4 Sagittal image after multiplanar reconstruction.




Fig. 5 Three-dimensional reconstruction showing the long osteophyte (L2 vertebra) in direct contact with the posterior aortic wall at the level of contained rupture.

an ultrasound done for other reasons. The patient subsequently underwent computed tomography (CT) imaging, where a 5.7-cm infrarenal AAA with irregular wall boundaries was seen. A very distinctive appearance compared with the typical fusiform AAAs was found (► **Fig. 1**).

The patient was admitted and operated via open surgery. Endovascular options were not considered for this case because we were concerned that the irregular appearance of the lesion could be related to an infectious process. Intraoperatively, the unusual appearance of the lesion was obvious (► **Fig. 2**). After sac incision, a posterior aortic wall perforation became apparent from a lumbar osteophyte protruding into the lumen (► **Fig. 3**). The arterial lesion was resected, while the osteophyte was trimmed to avoid compression of the new interposition graft (expanded polytetrafluoroethylene, 16 mm) that was implanted. The postoperative course was uncomplicated.

Retrospective examination of the CT angiography identified the bony structure responsible for aortic perforation (L2 vertebra), which had not been apparent initially (► **Figs. 4** and **5**). The patient reported chronic back pain and an incident of pain exaggeration 20 days ago which was treated with anti-inflammatory medication.

To the best of our knowledge, this is the first reported case of unprovoked aortic perforation from a lumbar osteophyte, while there have been a few reports of a similar lesion after blunt trauma.^{1,2}

Funding

None.

Conflict of Interest

The authors declare no conflict of interest related to this article.

Acknowledgments

None.

References

- 1 Myers PO, Hachulla-Lemaire AL, Murith N. Traumatic thoracic aortic rupture: caught between a thoracic vertebral osteophyte and a hard place. *J Thorac Cardiovasc Surg* 2015;150(06): 1661–1662
- 2 Vernon SA, Murphy WR, Murphy TW, Haan JM. Abdominal aortic rupture from an impaling osteophyte following blunt trauma. *J Vasc Surg* 2014;59(04):1112–1115