

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N'-(*E*)-1-(2-Hydroxyphenyl)ethylidene]pyrazine-2-carbohydrazide

 Shahid Hameed,^a Mushtaq Ahmad,^{a,b} M. Nawaz Tahir,^{c*} Muhammad Israr^d and Muhammad Anwar^d
^aDepartment of Chemistry, Quaid-i-Azam University, Islamabad, Pakistan,

^bMedicinal Botanic Centre, PCSIR Laboratories Complex, Peshawar, Pakistan,

^cUniversity of Sargodha, Department of Physics, Sargodha, Pakistan, and

^dDepartment of Chemistry, Kohat University of Science and Technology, Kohat, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 7 August 2013; accepted 7 August 2013

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.050; wR factor = 0.138; data-to-parameter ratio = 13.4.

The title compound, $\text{C}_{13}\text{H}_{12}\text{N}_4\text{O}_2$, crystallized with two independent molecules (*A* and *B*) in the asymmetric unit. Molecule *B* is planar to within 0.044 (3) Å for all non-H atoms, while molecule *A* is slightly twisted, with a dihedral angle of 6.29 (4)° between the mean planes of the pyrazine-2-carbohydrazide and 1-(2-hydroxyphenyl)ethanone moieties (r.m.s. deviations = 0.0348 and 0.0428 Å, respectively). *S*(5) and *S*(6) ring motifs are formed in both molecules due to the presence of intramolecular O—H···N and N—H···N hydrogen bonds. In the crystal, molecules *A* and *B* are linked by a C—H···O hydrogen bond. They stack along the *a*-axis direction, forming columns with π - π interactions involving inversion-related pyrazine and benzene rings [centroid-centroid distances = 3.5489 (13)–3.8513 (16) Å].

Related literature

For a related crystal structure and other studies, see: Hameed *et al.* (2013). For graph-set notation, see: Bernstein *et al.* (1995).



Experimental

Crystal data

 $\text{C}_{13}\text{H}_{12}\text{N}_4\text{O}_2$
 $M_r = 256.27$

 Triclinic, $P\bar{1}$
 $a = 7.1767$ (7) Å

 $b = 10.1743$ (10) Å
 $c = 17.1150$ (17) Å
 $\alpha = 86.172$ (3)°
 $\beta = 85.275$ (2)°
 $\gamma = 80.963$ (4)°
 $V = 1228.2$ (2) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.10$ mm⁻¹
 $T = 296$ K

 $0.28 \times 0.23 \times 0.20$ mm

Data collection

 Bruker Kappa APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.973$, $T_{\max} = 0.981$

 19042 measured reflections
 4821 independent reflections
 2608 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.138$
 $S = 1.00$
 4821 reflections
 361 parameters

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.16$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1···N1	0.82	1.82	2.537 (2)	145
N2—H2A···N3	0.80 (2)	2.26 (2)	2.654 (3)	111.5 (17)
O3—H3A···N5	0.82	1.82	2.534 (3)	145
N6—H6···N7	0.80 (3)	2.21 (3)	2.628 (3)	113 (3)
C3—H3···O3 ⁱ	0.93	2.59	3.403 (3)	146

 Symmetry code: (i) $-x + 1, -y + 1, -z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

The authors acknowledge the provision of funds for the purchase of the diffractometer and encouragement by Dr Muhammad Akram Chaudhary, Vice Chancellor, University of Sargodha, Pakistan, and also thank the Higher Education Commission (HEC) of Pakistan for financial support. MA is grateful to the Pakistan Council of Scientific and Industrial Research (PCSIR) Laboratories of Pakistan for financial support through out his study leave.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2634).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
 Hameed, S., Ahmad, M., Tahir, M. N., Shah, M. A. & Shad, H. A. (2013). *Acta Cryst. E* **69**, o1141.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2013). E69, o1419 [doi:10.1107/S1600536813022137]

***N'*-(*E*)-1-(2-Hydroxyphenyl)ethylidene]pyrazine-2-carbohydrazide**

Shahid Hameed, Mushtaq Ahmad, M. Nawaz Tahir, Muhammad Israr and Muhammad Anwar

1. Comment

The title compound was prepared in continuation of our interest in the synthesis of compounds containing the moiety pyrazine-2-carbohydrazide (Hameed *et al.*, 2013).

The title compound crystallized with two independent molecules (A and B) in the asymmetric unit, Fig. 1. In molecule A, the 1-(2-hydroxyphenyl)ethanone (C1—C8/O1) moiety and the pyrazine-2-carbohydrazide moiety (C9—C13/N1—N4/O2) are almost planar with r.m.s. deviations of 0.0348 Å and 0.0428 Å, respectively. They are inclined to one another by 6.289 (44)°. In molecule B, similar groups (C14—C21/O3) and (C22—C26/N5—N8/O4) are planar with r.m.s. deviations of 0.0111 and 0.019 Å, respectively, and are inclined to one another by only 0.305 (21)° *i.e.* almost coplanar.

There exist strong intramolecular N—H···N and O—H···N hydrogen bonds in each molecule (Table 1 and Fig. 1) forming S(5) and S(6) ring motifs (Bernstein *et al.*, 1995).

In the crystal, molecules A and B are linked by a C—H···O hydrogen bond (Table 1 and Fig. 2). They stack along the *a* axis direction forming columns with π – π interactions involving inversion related pyrazine and benzene rings. The centroid-to-centroid distances are 3.5489 (13) Å [Cg1—Cg2ⁱ], 3.6289 (13) Å [Cg1—Cg2ⁱⁱ], 3.7738 (16) Å [Cg3—Cg4ⁱⁱⁱ], and 3.8513 (16) Å [Cg3—Cg4^{iv}], where Cg1, Cg2, Cg3 and Cg4 are the centroids of rings (C10/N3/C11/C12/N4/C13), (C1—C6), (C23/N7/C24/C25/N8/C26) and (C14—C19), respectively [symmetry codes: (i) = $-x, -y, -z$; (ii) = $-x+1, -y, -z$; (iii) = $-x, -y+1, -z+1$; (iv) = $-x+1, -y+1, -z+1$].

2. Experimental


The title compound was prepared by the condensation of an equimolar ratio of pyrazine-2-carbohydrazide (0.50 g, 3.6 mmol) and 1-(2-hydroxyphenyl)ethanone (0.45 ml, 3.6 mmol) in methanol by stirring well and then refluxing of 5 h. The resulting reaction mixture was allowed to cool over night. The precipitated solid was filtered, washed with petroleum ether and recrystallized from chloroform in petroleum ether and then dried under reduced pressure over CaCl₂ to give colourless prisms.

3. Refinement


The H-atom of the amide and one of the methyl groups were refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ and $= 1.5U_{\text{eq}}(\text{C-methyl})$. The other H-atoms were positioned geometrically (C—H = 0.93 - 0.96 Å, O—H = 0.82 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = x \times U_{\text{eq}}(\text{C,N,O})$, where $x = 1.5$ for hydroxy and methyl H atoms and $= 1.2$ for other H atoms.

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 2012) and *PLATON* (Spek, 2009).

**Figure 1**

View of the molecular structure of the two independent molecules (A and B) of the title compound, with atom labelling. Displacement ellipsoids are drawn at the 50% probability level. The intramolecular hydrogen bonds are shown as dashed lines (see Table 1 for details).

**Figure 2**

The crystal packing diagram of the title compound, viewed along the *a* axis. The various hydrogen bonds are shown as dashed lines (see Table 1 for details).

N'-[(*E*)-1-(2-Hydroxyphenyl)ethylidene]pyrazine-2-carbohydrazide

Crystal data

$C_{13}H_{12}N_4O_2$

$M_r = 256.27$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.1767 (7) \text{ \AA}$

$b = 10.1743 (10) \text{ \AA}$

$c = 17.1150 (17) \text{ \AA}$

$\alpha = 86.172 (3)^\circ$

$\beta = 85.275 (2)^\circ$
 $\gamma = 80.963 (4)^\circ$
 $V = 1228.2 (2) \text{ \AA}^3$
 $Z = 4$
 $F(000) = 536$
 $D_x = 1.386 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2608 reflections
 $\theta = 1.2\text{--}26.0^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Prism, colourless
 $0.28 \times 0.23 \times 0.20 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: $8.00 \text{ pixels mm}^{-1}$
 ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.973$, $T_{\max} = 0.981$

19042 measured reflections
 4821 independent reflections
 2608 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.048$
 $\theta_{\max} = 26.0^\circ$, $\theta_{\min} = 1.2^\circ$
 $h = -8 \rightarrow 8$
 $k = -12 \rightarrow 12$
 $l = -21 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.138$
 $S = 1.00$
 4821 reflections
 361 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0537P)^2 + 0.2363P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0083 (12)

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.4150 (3)	0.07549 (16)	-0.16151 (9)	0.0602 (7)
O2	0.2080 (2)	-0.19271 (16)	-0.05633 (10)	0.0582 (6)
N1	0.2781 (2)	0.04641 (18)	-0.02111 (10)	0.0372 (6)
N2	0.2155 (3)	-0.04221 (18)	0.03446 (11)	0.0398 (7)
N3	0.0878 (2)	-0.20255 (18)	0.14825 (12)	0.0456 (7)
N4	0.0423 (3)	-0.46305 (19)	0.11955 (14)	0.0545 (8)
C1	0.4198 (3)	0.2026 (2)	-0.14404 (14)	0.0421 (8)
C2	0.4732 (3)	0.2888 (3)	-0.20396 (15)	0.0561 (10)
C3	0.4803 (3)	0.4192 (3)	-0.19149 (17)	0.0604 (11)

C4	0.4368 (3)	0.4658 (2)	-0.11752 (17)	0.0560 (10)
C5	0.3875 (3)	0.3803 (2)	-0.05687 (14)	0.0448 (8)
C6	0.3735 (3)	0.2475 (2)	-0.06741 (13)	0.0358 (7)
C7	0.3094 (3)	0.1610 (2)	-0.00190 (12)	0.0352 (7)
C8	0.2827 (3)	0.2063 (2)	0.08019 (13)	0.0515 (9)
C9	0.1834 (3)	-0.1606 (2)	0.01156 (14)	0.0382 (8)
C10	0.1186 (3)	-0.2501 (2)	0.07671 (13)	0.0371 (8)
C11	0.0343 (3)	-0.2871 (3)	0.20495 (16)	0.0561 (10)
C12	0.0136 (3)	-0.4158 (3)	0.19042 (17)	0.0573 (10)
C13	0.0928 (3)	-0.3772 (2)	0.06276 (15)	0.0469 (9)
O3	0.2432 (3)	0.48626 (18)	0.34993 (10)	0.0754 (8)
O4	0.1729 (3)	0.7739 (2)	0.48091 (12)	0.0870 (9)
N5	0.2460 (3)	0.5031 (2)	0.49687 (11)	0.0491 (7)
N6	0.2206 (3)	0.5847 (2)	0.55841 (12)	0.0557 (8)
N7	0.1834 (3)	0.7165 (2)	0.68710 (12)	0.0550 (8)
N8	0.0968 (3)	0.9941 (2)	0.68391 (16)	0.0754 (10)
C14	0.2858 (3)	0.3535 (3)	0.36491 (14)	0.0500 (9)
C15	0.3115 (4)	0.2738 (3)	0.30127 (15)	0.0617 (11)
C16	0.3510 (4)	0.1389 (3)	0.31072 (17)	0.0671 (11)
C17	0.3662 (4)	0.0795 (3)	0.38440 (19)	0.0747 (11)
C18	0.3421 (4)	0.1569 (3)	0.44851 (16)	0.0682 (11)
C19	0.3018 (3)	0.2958 (2)	0.44095 (13)	0.0454 (8)
C20	0.2773 (3)	0.3760 (3)	0.51033 (13)	0.0496 (9)
C21	0.2897 (7)	0.3111 (4)	0.5907 (2)	0.104 (2)
C22	0.1843 (4)	0.7184 (3)	0.54528 (16)	0.0556 (10)
C23	0.1605 (3)	0.7897 (3)	0.62001 (15)	0.0502 (9)
C24	0.1623 (4)	0.7838 (3)	0.75194 (15)	0.0633 (10)
C25	0.1216 (4)	0.9200 (3)	0.74989 (18)	0.0689 (11)
C26	0.1158 (4)	0.9267 (3)	0.61878 (17)	0.0693 (11)
H1	0.36890	0.03550	-0.12340	0.0900*
H2	0.50520	0.25780	-0.25390	0.0670*
H2A	0.202 (3)	-0.031 (2)	0.0805 (13)	0.0480*
H3	0.51450	0.47650	-0.23300	0.0720*
H4	0.44080	0.55450	-0.10880	0.0670*
H5	0.36250	0.41190	-0.00680	0.0540*
H8A	0.27220	0.13110	0.11640	0.0770*
H8B	0.16950	0.27020	0.08570	0.0770*
H8C	0.38940	0.24660	0.09110	0.0770*
H11	0.01000	-0.25890	0.25590	0.0670*
H12	-0.02200	-0.47180	0.23230	0.0690*
H13	0.11150	-0.40420	0.01150	0.0560*
H3A	0.23690	0.52440	0.39100	0.1130*
H6	0.234 (4)	0.557 (3)	0.6029 (15)	0.0670*
H15	0.30140	0.31340	0.25090	0.0740*
H16	0.36760	0.08720	0.26720	0.0810*
H17	0.39280	-0.01280	0.39120	0.0900*
H18	0.35300	0.11540	0.49840	0.0820*
H21A	0.250 (6)	0.367 (4)	0.628 (3)	0.1560*
H21B	0.409 (6)	0.269 (4)	0.598 (2)	0.1560*

H21C	0.210 (6)	0.246 (4)	0.597 (2)	0.1560*
H24	0.17570	0.73700	0.80020	0.0760*
H25	0.11100	0.96230	0.79690	0.0820*
H26	0.09830	0.97380	0.57080	0.0830*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0827 (13)	0.0529 (11)	0.0456 (11)	-0.0117 (9)	0.0008 (9)	-0.0108 (9)
O2	0.0818 (12)	0.0506 (10)	0.0459 (11)	-0.0196 (9)	-0.0032 (9)	-0.0089 (9)
N1	0.0369 (10)	0.0324 (10)	0.0419 (11)	-0.0046 (8)	-0.0035 (8)	0.0004 (9)
N2	0.0458 (11)	0.0354 (11)	0.0388 (12)	-0.0082 (8)	-0.0029 (9)	-0.0019 (10)
N3	0.0471 (11)	0.0425 (12)	0.0472 (13)	-0.0096 (9)	-0.0004 (9)	-0.0001 (10)
N4	0.0479 (12)	0.0366 (12)	0.0775 (17)	-0.0067 (9)	-0.0032 (11)	0.0063 (12)
C1	0.0404 (13)	0.0423 (14)	0.0440 (15)	-0.0061 (10)	-0.0073 (10)	-0.0003 (12)
C2	0.0533 (16)	0.0713 (19)	0.0426 (16)	-0.0109 (13)	-0.0040 (12)	0.0079 (14)
C3	0.0508 (16)	0.066 (2)	0.066 (2)	-0.0227 (13)	-0.0113 (14)	0.0253 (16)
C4	0.0527 (15)	0.0438 (15)	0.075 (2)	-0.0178 (12)	-0.0159 (14)	0.0102 (15)
C5	0.0430 (13)	0.0397 (14)	0.0530 (16)	-0.0082 (10)	-0.0080 (11)	-0.0020 (12)
C6	0.0310 (11)	0.0341 (13)	0.0418 (14)	-0.0024 (9)	-0.0072 (9)	0.0012 (10)
C7	0.0321 (11)	0.0322 (13)	0.0409 (14)	-0.0008 (9)	-0.0061 (9)	-0.0049 (10)
C8	0.0696 (16)	0.0409 (14)	0.0450 (15)	-0.0124 (12)	-0.0007 (12)	-0.0045 (12)
C9	0.0372 (12)	0.0332 (13)	0.0449 (15)	-0.0032 (10)	-0.0080 (10)	-0.0054 (11)
C10	0.0305 (11)	0.0331 (13)	0.0468 (15)	-0.0021 (9)	-0.0048 (10)	-0.0005 (11)
C11	0.0584 (16)	0.0580 (17)	0.0517 (17)	-0.0135 (13)	0.0022 (12)	0.0015 (14)
C12	0.0470 (15)	0.0517 (17)	0.071 (2)	-0.0116 (12)	0.0003 (13)	0.0166 (15)
C13	0.0451 (14)	0.0353 (14)	0.0600 (17)	-0.0040 (11)	-0.0044 (12)	-0.0044 (13)
O3	0.1333 (18)	0.0556 (12)	0.0381 (11)	-0.0212 (11)	0.0008 (11)	-0.0001 (9)
O4	0.148 (2)	0.0666 (13)	0.0462 (13)	-0.0209 (13)	-0.0017 (12)	0.0016 (11)
N5	0.0622 (13)	0.0506 (13)	0.0357 (12)	-0.0122 (10)	0.0009 (9)	-0.0085 (10)
N6	0.0763 (15)	0.0575 (15)	0.0338 (12)	-0.0110 (11)	-0.0004 (11)	-0.0090 (12)
N7	0.0584 (13)	0.0605 (14)	0.0467 (14)	-0.0059 (10)	-0.0033 (10)	-0.0146 (12)
N8	0.0934 (19)	0.0602 (16)	0.0776 (19)	-0.0186 (13)	-0.0074 (15)	-0.0238 (15)
C14	0.0542 (15)	0.0568 (17)	0.0399 (15)	-0.0139 (12)	0.0040 (11)	-0.0066 (13)
C15	0.0750 (19)	0.071 (2)	0.0413 (16)	-0.0182 (15)	0.0036 (13)	-0.0133 (14)
C16	0.0650 (18)	0.080 (2)	0.058 (2)	-0.0068 (15)	-0.0027 (14)	-0.0279 (17)
C17	0.092 (2)	0.0533 (17)	0.077 (2)	0.0073 (15)	-0.0161 (17)	-0.0223 (17)
C18	0.087 (2)	0.0600 (19)	0.0557 (18)	-0.0001 (15)	-0.0147 (15)	-0.0041 (15)
C19	0.0470 (14)	0.0497 (15)	0.0397 (15)	-0.0068 (11)	-0.0023 (11)	-0.0055 (12)
C20	0.0567 (15)	0.0554 (17)	0.0361 (14)	-0.0074 (12)	-0.0025 (11)	-0.0026 (12)
C21	0.194 (5)	0.072 (3)	0.0387 (19)	0.002 (3)	-0.011 (2)	0.0014 (17)
C22	0.0675 (17)	0.0532 (18)	0.0481 (18)	-0.0152 (13)	-0.0005 (13)	-0.0078 (14)
C23	0.0532 (15)	0.0532 (16)	0.0468 (17)	-0.0140 (12)	-0.0013 (12)	-0.0101 (13)
C24	0.0668 (17)	0.075 (2)	0.0491 (17)	-0.0044 (14)	-0.0086 (13)	-0.0206 (15)
C25	0.0684 (19)	0.076 (2)	0.066 (2)	-0.0119 (16)	-0.0047 (15)	-0.0304 (18)
C26	0.091 (2)	0.0570 (19)	0.063 (2)	-0.0200 (15)	-0.0050 (15)	-0.0071 (16)

Geometric parameters (Å, °)

O1—C1	1.353 (3)	C11—C12	1.382 (4)
O2—C9	1.219 (3)	C2—H2	0.9300
O1—H1	0.8200	C3—H3	0.9300
O3—C14	1.349 (3)	C4—H4	0.9300
O4—C22	1.207 (3)	C5—H5	0.9300
O3—H3A	0.8200	C8—H8B	0.9600
N1—N2	1.364 (3)	C8—H8C	0.9600
N1—C7	1.290 (3)	C8—H8A	0.9600
N2—C9	1.350 (3)	C11—H11	0.9300
N3—C11	1.327 (3)	C12—H12	0.9300
N3—C10	1.335 (3)	C13—H13	0.9300
N4—C13	1.329 (3)	C14—C15	1.384 (4)
N4—C12	1.323 (4)	C14—C19	1.398 (3)
N2—H2A	0.80 (2)	C15—C16	1.359 (4)
N5—N6	1.367 (3)	C16—C17	1.367 (4)
N5—C20	1.286 (4)	C17—C18	1.377 (4)
N6—C22	1.351 (4)	C18—C19	1.397 (4)
N7—C24	1.328 (3)	C19—C20	1.469 (3)
N7—C23	1.334 (3)	C20—C21	1.489 (4)
N8—C25	1.324 (4)	C22—C23	1.497 (4)
N8—C26	1.334 (4)	C23—C26	1.380 (4)
N6—H6	0.80 (3)	C24—C25	1.369 (4)
C1—C6	1.414 (3)	C15—H15	0.9300
C1—C2	1.374 (4)	C16—H16	0.9300
C2—C3	1.367 (4)	C17—H17	0.9300
C3—C4	1.376 (4)	C18—H18	0.9300
C4—C5	1.369 (3)	C21—H21A	0.88 (5)
C5—C6	1.395 (3)	C21—H21B	0.91 (4)
C6—C7	1.468 (3)	C21—H21C	0.94 (4)
C7—C8	1.496 (3)	C24—H24	0.9300
C9—C10	1.484 (3)	C25—H25	0.9300
C10—C13	1.374 (3)	C26—H26	0.9300
C1—O1—H1	109.00	C7—C8—H8A	109.00
C14—O3—H3A	110.00	N3—C11—H11	119.00
N2—N1—C7	120.53 (18)	C12—C11—H11	119.00
N1—N2—C9	118.47 (18)	N4—C12—H12	118.00
C10—N3—C11	115.5 (2)	C11—C12—H12	119.00
C12—N4—C13	115.0 (2)	C10—C13—H13	119.00
C9—N2—H2A	116.9 (15)	N4—C13—H13	119.00
N1—N2—H2A	124.6 (15)	C15—C14—C19	120.1 (3)
N6—N5—C20	119.58 (19)	O3—C14—C15	117.3 (2)
N5—N6—C22	120.3 (2)	O3—C14—C19	122.6 (2)
C23—N7—C24	115.9 (2)	C14—C15—C16	121.5 (3)
C25—N8—C26	115.2 (2)	C15—C16—C17	119.8 (3)
C22—N6—H6	117 (2)	C16—C17—C18	119.7 (3)
N5—N6—H6	123 (2)	C17—C18—C19	122.0 (3)
O1—C1—C2	117.5 (2)	C18—C19—C20	120.9 (2)

C2—C1—C6	120.1 (2)	C14—C19—C18	116.9 (2)
O1—C1—C6	122.37 (19)	C14—C19—C20	122.2 (2)
C1—C2—C3	121.3 (2)	N5—C20—C21	123.2 (3)
C2—C3—C4	120.0 (3)	N5—C20—C19	116.0 (2)
C3—C4—C5	119.4 (2)	C19—C20—C21	120.8 (3)
C4—C5—C6	122.5 (2)	O4—C22—N6	124.0 (3)
C1—C6—C5	116.7 (2)	O4—C22—C23	123.9 (3)
C1—C6—C7	122.37 (18)	N6—C22—C23	112.0 (2)
C5—C6—C7	120.9 (2)	C22—C23—C26	120.6 (2)
N1—C7—C6	115.00 (18)	N7—C23—C22	117.8 (3)
C6—C7—C8	121.04 (18)	N7—C23—C26	121.6 (2)
N1—C7—C8	123.96 (19)	N7—C24—C25	122.0 (3)
O2—C9—N2	123.4 (2)	N8—C25—C24	122.9 (3)
O2—C9—C10	122.60 (19)	N8—C26—C23	122.4 (3)
N2—C9—C10	114.0 (2)	C14—C15—H15	119.00
N3—C10—C13	122.1 (2)	C16—C15—H15	119.00
C9—C10—C13	120.3 (2)	C15—C16—H16	120.00
N3—C10—C9	117.61 (18)	C17—C16—H16	120.00
N3—C11—C12	121.8 (3)	C16—C17—H17	120.00
N4—C12—C11	122.9 (3)	C18—C17—H17	120.00
N4—C13—C10	122.7 (2)	C17—C18—H18	119.00
C3—C2—H2	119.00	C19—C18—H18	119.00
C1—C2—H2	119.00	C20—C21—H21A	113 (3)
C2—C3—H3	120.00	C20—C21—H21B	111 (2)
C4—C3—H3	120.00	C20—C21—H21C	109 (2)
C3—C4—H4	120.00	H21A—C21—H21B	111 (4)
C5—C4—H4	120.00	H21A—C21—H21C	106 (4)
C4—C5—H5	119.00	H21B—C21—H21C	107 (4)
C6—C5—H5	119.00	N7—C24—H24	119.00
H8A—C8—H8B	110.00	C25—C24—H24	119.00
H8A—C8—H8C	109.00	N8—C25—H25	119.00
C7—C8—H8C	109.00	C24—C25—H25	119.00
C7—C8—H8B	109.00	N8—C26—H26	119.00
H8B—C8—H8C	110.00	C23—C26—H26	119.00
C7—N1—N2—C9	179.6 (2)	C1—C6—C7—N1	-7.5 (3)
N2—N1—C7—C6	-179.11 (18)	C1—C6—C7—C8	173.0 (2)
N2—N1—C7—C8	0.5 (3)	C5—C6—C7—C8	-8.5 (3)
N1—N2—C9—O2	0.2 (3)	N2—C9—C10—N3	4.2 (3)
N1—N2—C9—C10	179.11 (18)	N2—C9—C10—C13	-175.6 (2)
C11—N3—C10—C9	-178.49 (19)	O2—C9—C10—N3	-177.0 (2)
C11—N3—C10—C13	1.2 (3)	O2—C9—C10—C13	3.3 (3)
C10—N3—C11—C12	0.5 (3)	N3—C10—C13—N4	-2.5 (3)
C13—N4—C12—C11	0.0 (3)	C9—C10—C13—N4	177.2 (2)
C12—N4—C13—C10	1.8 (3)	N3—C11—C12—N4	-1.1 (4)
C20—N5—N6—C22	-178.8 (2)	O3—C14—C15—C16	-178.7 (3)
N6—N5—C20—C19	179.7 (2)	C19—C14—C15—C16	0.5 (4)
N6—N5—C20—C21	-0.7 (4)	O3—C14—C19—C18	178.5 (2)
N5—N6—C22—O4	-0.4 (4)	O3—C14—C19—C20	-1.3 (3)

N5—N6—C22—C23	179.8 (2)	C15—C14—C19—C18	-0.6 (3)
C24—N7—C23—C22	179.6 (2)	C15—C14—C19—C20	179.5 (2)
C24—N7—C23—C26	-0.6 (4)	C14—C15—C16—C17	-0.1 (4)
C23—N7—C24—C25	-0.7 (4)	C15—C16—C17—C18	-0.3 (4)
C26—N8—C25—C24	-0.5 (4)	C16—C17—C18—C19	0.1 (4)
C25—N8—C26—C23	-0.8 (4)	C17—C18—C19—C14	0.3 (4)
O1—C1—C6—C5	178.8 (2)	C17—C18—C19—C20	-179.8 (2)
O1—C1—C6—C7	-2.6 (3)	C14—C19—C20—N5	-2.1 (3)
C2—C1—C6—C7	177.9 (2)	C14—C19—C20—C21	178.2 (3)
O1—C1—C2—C3	179.5 (2)	C18—C19—C20—N5	178.1 (2)
C2—C1—C6—C5	-0.8 (3)	C18—C19—C20—C21	-1.7 (4)
C6—C1—C2—C3	-0.9 (3)	O4—C22—C23—N7	-177.6 (3)
C1—C2—C3—C4	1.2 (4)	O4—C22—C23—C26	2.6 (4)
C2—C3—C4—C5	0.3 (3)	N6—C22—C23—N7	2.2 (3)
C3—C4—C5—C6	-2.1 (3)	N6—C22—C23—C26	-177.6 (2)
C4—C5—C6—C7	-176.4 (2)	N7—C23—C26—N8	1.4 (4)
C4—C5—C6—C1	2.3 (3)	C22—C23—C26—N8	-178.8 (2)
C5—C6—C7—N1	171.1 (2)	N7—C24—C25—N8	1.3 (5)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 \cdots N1	0.82	1.82	2.537 (2)	145
N2—H2A \cdots N3	0.80 (2)	2.26 (2)	2.654 (3)	111.5 (17)
O3—H3A \cdots N5	0.82	1.82	2.534 (3)	145
N6—H6 \cdots N7	0.80 (3)	2.21 (3)	2.628 (3)	113 (3)
C3—H3 \cdots O3 ⁱ	0.93	2.59	3.403 (3)	146

Symmetry code: (i) $-x+1, -y+1, -z$.