

# Transient Receptor Potential Channel Polymorphisms Are Associated with the Somatosensory Function in Neuropathic Pain Patients

Andreas Binder<sup>1\*</sup>, Denisa May<sup>2,9</sup>, Ralf Baron<sup>1</sup>, Christoph Maier<sup>3</sup>, Thomas R. Tölle<sup>4</sup>, Rolf-Detlef Treede<sup>5</sup>, Achim Berthele<sup>4</sup>, Frank Faltraco<sup>2</sup>, Herta Flor<sup>6</sup>, Janne Gierthmühlen<sup>1</sup>, Sierk Haenisch<sup>2</sup>, Volker Hüge<sup>7</sup>, Walter Magerl<sup>5</sup>, Christian Maihöfner<sup>8</sup>, Helmut Richter<sup>3</sup>, Roman Rolke<sup>9</sup>, Andrea Scherens<sup>3</sup>, Nurcan Üçeyler<sup>10</sup>, Mike Ufer<sup>2</sup>, Gunnar Wasner<sup>1</sup>, Jihong Zhu<sup>2</sup>, Ingolf Cascorbi<sup>2</sup>

**1** Division of Neurological Pain Research and Therapy, Department of Neurology, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany, **2** Institute of Experimental and Clinical Pharmacology, University Hospital Schleswig-Holstein, Campus Kiel, Kiel, Germany, **3** Department of Pain Management, BG Kliniken Bergmannsheil, Ruhr University Bochum, Bochum, Germany, **4** Department of Neurology, Technische Universität München, Klinikum rechts der Isar, München, Germany, **5** Center for Biomedicine and Medical Technology Mannheim, Ruprecht-Karls-University Heidelberg, Mannheim, Germany, **6** Institute of Clinical and Cognitive Neuroscience, Central Institute for Mental Health, Ruprecht-Karls-University Heidelberg, Mannheim, Germany, **7** Department of Anaesthesiology, Ludwig-Maximilians-University München, München, Germany, **8** Department of Neurology, University of Erlangen, Erlangen, Germany, **9** Department of Neurology, University Medical Center, Johannes-Gutenberg-University of Mainz, Mainz and Department of Palliative Care, University of Bonn, Bonn, Germany, **10** Department of Neurology, University of Würzburg, Würzburg, Germany

## Abstract

Transient receptor potential channels are important mediators of thermal and mechanical stimuli and play an important role in neuropathic pain. The contribution of hereditary variants in the genes of transient receptor potential channels to neuropathic pain is unknown. We investigated the frequency of transient receptor potential ankyrin 1, transient receptor potential melastin 8 and transient receptor potential vanilloid 1 single nucleotide polymorphisms and their impact on somatosensory abnormalities in neuropathic pain patients. Within the German Research Network on Neuropathic Pain (Deutscher Forschungsverbund Neuropathischer Schmerz) 371 neuropathic pain patients were phenotypically characterized using standardized quantitative sensory testing. Pyrosequencing was employed to determine a total of eleven single nucleotide polymorphisms in transient receptor potential channel genes of the neuropathic pain patients and a cohort of 253 German healthy volunteers. Associations of quantitative sensory testing parameters and single nucleotide polymorphisms between and within groups and subgroups, based on sensory phenotypes, were analyzed. Single nucleotide polymorphisms frequencies did not differ between both the cohorts. However, in neuropathic pain patients transient receptor potential ankyrin 1 710G>A (rs920829, E179K) was associated with the presence of paradoxical heat sensation ( $p = 0.03$ ), and transient receptor potential vanilloid 1 1911A>G (rs8065080, I585V) with cold hypoalgesia ( $p = 0.0035$ ). Two main subgroups characterized by preserved (1) and impaired (2) sensory function were identified. In subgroup 1 transient receptor potential vanilloid 1 1911A>G led to significantly less heat hyperalgesia, pinprick hyperalgesia and mechanical hypaesthesia ( $p = 0.006$ ,  $p = 0.005$  and  $p < 0.001$ ) and transient receptor potential vanilloid 1 1103C>G (rs222747, M315I) to cold hypaesthesia ( $p = 0.002$ ), but there was absence of associations in subgroup 2. In this study we found no evidence that genetic variants of transient receptor potential channels are involved in the expression of neuropathic pain, but transient receptor potential channel polymorphisms contributed significantly to the somatosensory abnormalities of neuropathic pain patients.

**Citation:** Binder A, May D, Baron R, Maier C, Tölle TR, et al. (2011) Transient Receptor Potential Channel Polymorphisms Are Associated with the Somatosensory Function in Neuropathic Pain Patients. PLoS ONE 6(3): e17387. doi:10.1371/journal.pone.0017387

**Editor:** Carlo Gaetano, Istituto Dermopatico dell'Immacolata, Italy

**Received:** November 25, 2010; **Accepted:** February 1, 2011; **Published:** March 29, 2011

**Copyright:** © 2011 Binder et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by the German Ministry for Education and Research, BMBF, an unrestricted educational grant from Pfizer Pharma, the National Health and Medical Research Council of Australia (NHMRC), the EFIC-Grünenthal-Grant (EGG), the Ministry of Science, Commerce and Transportation of Schleswig-Holstein, Germany and the Alexander von Humboldt-Stiftung. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** A. Binder received support for travel from the BMBF, honoraria for lectures from Grünenthal, Allergan and Pfizer, payment for development of educational presentations from Pfizer and travel/accommodation expenses from Pfizer, Allergan and Grünenthal and his institution received financial support from the BMBF. RB received support for travel from the BMBF, has consultancies with and received honoraria and travel/accommodation expenses from Pfizer, Genzyme, Grünenthal, Mundipharma, Allergan, Sanofi Pasteur, Medtronic, Eisai, UCB, Lilly, Astellas, received grants from Pfizer, Grünenthal, Genzyme, received payment for development of educational presentations from Pfizer, Medtronic and his institution received financial support from the BMBF. TT is board member at Pfizer, Lilly, Boehringer Grünenthal, and has consultancies with and received travel/accommodation expenses from Pfizer, Lilly, Boehringer, Grünenthal, Mundipharma, UCB/Schwarz. A. Berthele received honoraria and travel/accommodations expenses from Pfizer and his institution a grant from the BMBF. JG received support for travel from the BMBF and her institution received financial support from the BMBF. VH's institution received financial support from the BMBF. CMh received honoraria from Pfizer and travel/accommodation expenses from Allergan. RR is a board member at Astellas Pharma and received honoraria for lectures from Lilly and Pfizer. GW received honoraria and travel/accommodation expenses from Grünenthal, Medtronic, Pfizer, Mundipharma, his institution received financial support from the BMBF and Pfizer Pharma, and he received financial support for research from the National Health and Medical Research Council of Australia (NHMRC), the EFIC-Grünenthal-Grant (EGG), the Ministry of Science, Commerce and Transportation of Schleswig-Holstein, Germany and the Alexander von Humboldt-Stiftung. IC received honoraria and travel/accommodation expenses from MSD, Bayer-Schering and Sanofi Aventis for lectures. DM, RDT, FF, HF, SH, CM, AS, MU, NÜ, JZ, WM, HR have disclosed no competing interests. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLoS ONE policies on sharing data and materials.

\* E-mail: a.binder@neurologie.uni-kiel.de

† These authors contributed equally to this manuscript.

## Introduction

Neuropathic pain arises after lesions or diseases of the somatosensory nervous system and involves multiple somatosensory phenomena. In addition to spontaneous pain, these include negative and positive symptoms and signs such as hypaesthesia, hypoalgesia, thermal and mechanical hyperalgesia and allodynia [1]. These somatosensory abnormalities and their combination, i.e. the somatosensory profile, may mirror the underlying pain-generating mechanisms [1]. The German Research Network on Neuropathic Pain (DFNS) developed a standardized quantitative sensory testing (QST) battery for somatosensory phenotyping of neuropathic pain patients, including healthy control reference data, facilitating the routine clinical assessment [2,3].

The understanding of the molecular basis of neuropathic pain has been expanded by the cloning and characterization of the transient receptor potential (TRP) family of voltage-gated ion channels. Among these the TRP vanilloid 1 (TRPV1), TRP melastin 8 (TRPM8) and TRP ankyrin 1 (TRPA1) subfamilies have been found to play important roles in transduction and sensitisation in primary afferent somatosensory neurons [4]. TRPV1 is involved in the transduction of noxious heat and a receptor for capsaicin, thus mediating heat hyperalgesia after stimulation with this hot chilli constituent [5]. It is expressed on small diameter nociceptive neurons, likely to be C-fibers [6]. TRPM8 is also expressed on small and medium diameter sensory neurons, likely to be C and A-delta fibers. It is strongly activated by its agonist menthol and serves as a major cold and cooling transduction channel [7,8]. TRPA1, activated by its agonists cinnamon and mustard oil, transduces cold and mechanical stimuli. Moreover, on the peptide level it is 20% homologous to TRPV1 and is co-expressed with TRPV1 in a subpopulation of unmyelinated nociceptive neurons suggesting an important role in nociception [9].

Beside these physiological functions, TRPs are thought to play an important role in the generation of neuropathic pain, as demonstrated by *in vitro* and *in vivo* animal studies and neurophysiological and psychophysical studies in human experimental pain models and pain patients [10]. TRP channels have been shown to be differentially expressed following experimental nerve injury [11,12]. Moreover, enhanced expression of TRP channels on neighbouring undamaged fibres following peripheral nerve injury was previously described to regulate pain behaviour [11,12]. TRPV1 was shown to be involved in the activation and sensitisation of primary sensory neurons to heat, acid and different endogenous inflammatory mediators, leading to heat and mechanical hyperalgesia and spontaneous pain. Likewise TRPM8 contributes to cold pain, cold hyperalgesia and allodynia and TRPA1 leads to activation, sensitisation and facilitation of nociception like TRPV1 [11,12].

However, the contribution of a genetic variability in TRP channels on pain perception and pain generation in neuropathic pain patients is unknown. In this study we describe the results of an association analysis of polymorphisms in TRPV1, TRPM8 and TRPA1 genes with somatosensory signs of neuropathic pain patients. The aims were (1) to determine the frequencies of TRP variants in a large group of neuropathic pain patients of different aetiology and a cohort of healthy volunteers and (2) to determine the relationship of TRP gene variants and somatosensory function in neuropathic pain patients. Within this study we could demonstrate significant associations between the somatosensory function and TRP gene variants in neuropathic pain patients but no differences in the TRP frequencies compared with healthy volunteers.

## Methods

### Ethics statement

All subjects gave their written informed consent and the study was approved by the ethics committee of the Medical Faculty of the Christian-Albrechts-University Kiel and subsequently the Ethic Committees of the other participating DFNS centers and performed according to the Declaration of Helsinki.

### Study populations

A total of 371 patients with chronic neuropathic pain syndromes, all Caucasians, who participated in the German Research Network on Neuropathic Pain (DFNS; [www.neuro.med.tu-muenchen.de/dfns/](http://www.neuro.med.tu-muenchen.de/dfns/)) were included in the study (210 female, 161 male). The patients suffered from complex regional pain syndrome, postherpetic neuralgia, peripheral nerve injury, trigeminal neuropathy, polyneuropathy, central pain and neuropathies that did not fulfill diagnostic criteria of the above mentioned syndromes as diagnosed by expert physicians. Patient characteristics are listed in Table 1. For comparison of allelic frequencies of target polymorphisms, 253 healthy German volunteers (173 female, mean age  $26.8 \pm 6.9$  years; 80 male, mean age  $28.4 \pm 7.0$  years), all Caucasian, were included.

### Phenotypical characterization (quantitative sensory testing)

All patients were phenotypically characterized using standardized quantitative sensory testing (QST) established by the DFNS to determinate somatosensory function and signs of pain [2]. This QST battery consists of seven tests that assess 13 parameters. Briefly, thermal detection thresholds for the perception of cold (CDT) and warm (WDT), thermal pain thresholds for cold (CPT) and heat (HPT), paradoxical heat sensation (PHS), mechanical detection thresholds for touch (MDT) and vibration (VDT), mechanical pain sensitivity including thresholds for pinprick (MPT) and blunt pressure (PPT), a stimulus-response-function for pinprick sensitivity (MPS) and dynamic mechanical allodynia (DMA) as well as pain summation to repetitive pinprick stimuli (wind-up ratio; WUR) were determined in the area of pain. Patient data were included in the study analysis, if at least results of 12 QST tests were available.

### Genetic analysis

Whole genomic DNA was extracted from venous blood samples obtained from patients and volunteers using the Qiagen Genra Puregene Blood Kit (Qiagen, Hilden, Germany). The final concentration of gDNA was determined spectrophotometrically. Two SNPs in TRPV1, six in TRPM8 and three in TRPA1 genes were identified by literature review in the context of neuropathic pain and selected for detailed association analysis (Table 2). Except of TRPM8 787A>T and TRPM8 790G>C (both synonymous polymorphisms detected in parallel in our genotyping assay), all SNPs were carefully selected based on the assumption that missense variants may have functional consequences (detailed information are included in the Discussion section). Genotyping of the TRP polymorphisms was performed by pyrosequencing according to protocols provided by the manufacturer on a PSQ HS96 platform (Biotage AB, Uppsala, Sweden). Primer sequences and conditions are available on request.

### Statistical analysis

Before statistical analysis the QST values that did not show normal distribution (all except CPT, HPT, VDT and PHS) were transformed logarithmically [2]. Subsequently, Z-scores were calculated, except for DMA and PHS that are given with original values (DMA: 0–100

**Table 1.** Demographic data of patients.

	All neuropathic pain patients	Cluster 1	Cluster 2	p-values Cluster 1 vs. Cluster 2
N	371	121	120	
Age (years)	56.4±14.4	52.2±13.7	56.6±13.8	p<0.02
Females (%)	57.7	56.1	49.1	p=0.34
Disease duration (months)	43.9±71.9	39.1±78.6	35.1±47.4	p=0.63
Pain intensity (0–10 Likert scale NRS)	6.0±2.0	6.0±2.0	6.0±2.0	p=1.00
Polyneuropathy (n, %)	101 (27.2)	24 (19.8)	41 (35.0)	p<0.02
Post herpetic neuralgia (n, %)	35 (9.4)	7 (5.8)	14 (11.6)	p=0.16
Painful nerve injury (n, %)	34 (9.2)	13 (10.7)	14 (11.6)	p=0.98
CRPS (n, %)	87 (23.5)	40 (33.1)	29 (24.1)	p=0.17
Trigeminal pain (n, %)	57 (15.4)	28 (23.1)	7 (5.8)	p<0.0001
Central pain (n, %)	24 (6.5)	4 (3.3)	7 (5.8)	p=0.53
Other neuropathy (n, %)	33 (8.9)	5 (4.2)	8 (6.1)	p=0.56

Values are given as means ± single standard deviation, percentages or absolute values respectively. Pain intensity of the last four weeks was assessed using the numeric rating scale (NRS; 0 = no pain, 10 = worst pain imaginable).  
doi:10.1371/journal.pone.0017387.t001

numeric rating scale; PHS: numbers of PHS, 0–3) [2]. Z-scores above “0” indicate a gain of function (hyperalgesia, allodynia), while a Z-score below “0” indicates a loss of function (hypoesthesia, hypoalgesia). Pathological values were determined according normative data (95% confidence interval = Z-score of 0±1.96). Additionally, DMA and PHS were encoded as dichotomized values (1 = pathological, 0 = within normal range).

In order to identify major subpopulations of patients who are characterized by a typical combination of signs, a cluster analysis was performed in those patients only from whom complete QST results were available (241 of 371 patients, Table 1). The commonly recommended WARD-approach with an Euclidian distance measure was used as described elsewhere [13]. Only those clusters determined within the first WARD fusion algorithm calculation were used. The clusters represent significantly different somatosensory QST Z-profiles demonstrating cluster specific sensory profiles.

**Table 2.** Significant associations of TRPV1 variants and somatosensory function.

		QST test			
		CDT	HPT	MDT	MPS
<b>1103C&gt;G</b>	CC/CG/GG	0.0023 <sup>1</sup>			
	CC/CG+GG	0.0019 <sup>2</sup>			
	CC+CG/GG	0.0018 <sup>2</sup>			
<b>1911A&gt;G</b>	AA/AG/GG		0.0059 <sup>1</sup>	0.0004 <sup>1</sup>	n.s. <sup>1</sup>
	AA/AG+GG		n.s. <sup>2</sup>	<0.0001 <sup>2</sup>	0.005 <sup>2</sup>
	AA+AG/GG		0.0006 <sup>2</sup>	n.s. <sup>2</sup>	n.s. <sup>2</sup>

Significant associations of selected TRPV1 variants with three parameters of quantitative sensory testing (CDT, cold detection threshold, HPT, heat pain threshold, MDT, mechanical detection threshold) in 121 patients belonging to cluster 1. Z-score means of each QST parameter were correlated with genotypes applying a dominant or recessive model.

<sup>1</sup>ANOVA test,

<sup>2</sup>t-test.

doi:10.1371/journal.pone.0017387.t002

The statistical software SPSS 17.0 was employed for all statistical evaluations. Genotype, respectively allele distributions, in the population of patients and healthy controls as well as differences of dichotomized QST values related to target polymorphisms were assessed with Chi-square and Yates corrected Chi-square test respectively. Differences in measured QST parameters (means of Z-scores) and for bi-allelic polymorphisms each subtest and genotype were calculated using analysis of variance (ANOVA). For dominant and recessive models the t-test was used, if applicable. To correct for multiple testing each test was performed using 100,000 permutations.

## Results

### Genotype distribution


The frequency distribution of all polymorphisms investigated was in Hardy-Weinberg equilibrium. Moreover, after correcting for multiple comparisons, the frequency of all SNPs did not differ significantly between neuropathic pain patients, healthy controls and data from the HapMap project (Table S1).

### Association of TRP gene variants to QST Z-scores

First, within the group of patients who suffered from paradoxical heat sensation (PHS) heterozygous and homozygous carriers of the TRPA1 710G>A variant were significantly under-represented as compared to neuropathic pain patients without PHS (odds ratio 0.44, 95% confidence interval 0.21–0.90, p=0.03) (Fig. 1). Second, homozygous variant carriers (GG) of the TRPV1 1911A>G polymorphism exhibited significant cold hypoalgesia compared with heterozygous or wild type carriers (CPT Z-scores:  $-0.13 \pm 1.10$ ,  $0.19 \pm 1.20$ , and  $0.18 \pm 1.14$ , respectively; p=0.003).

### Somatosensory profiles of subpopulations

Hierarchical cluster analysis detected two main clusters (Fig. 2). Cluster 1 includes a group of patients with mainly preserved sensory function, while cluster 2 was characterized by a relative loss of small and large fiber function as detected by significant lower Z-scores of CDT, WDT, TSL, MDT (p<0.0001), VDT (p=0.002) and a higher Z-score of WUR (p=0.027) in patients




**Figure 1. Association of TRPA1 710G>A to appearance of paradoxical heat sensation.** Patients with positive paradoxical heat (PHS) showed a lower frequency of variant TRPA1 710 genotypes (GA+AA carriers) compared to patients with negative PHS (odds ratio 0.44, 95% confidence interval 0.21–0.90,  $p=0.03$ ). doi:10.1371/journal.pone.0017387.g001


who belong to cluster 2 as compared to cluster 1. DMA and PHS differed significantly between the two clusters. Pathological PHS occurred significantly more frequent in cluster 2 (odds ratio 3.44, 95% CI 1.78–6.68,  $p<0.0001$ ) and DMA was more frequent in cluster 1 (odds ratio 2.30, 95% CI 1.30–4.07,  $p=0.004$ ).

**Association of TRP gene variants to QST Z-scores of subpopulations**

Z-scores of all QST tests were stratified to all TRP variants investigated in the entire sample and in both clusters. After correcting for multiple comparisons, two TRPV1 polymorphisms,



**Figure 2. Somatosensory profiles of cluster 1 and cluster 2.** Cluster 1 is characterized by a predominant preserved somatosensory function whereas cluster 2 exhibits relative loss of small fiber and large fiber function (lower Z-scores for CDT, WDT, TSL, MDT, VDT and WUR respectively). Values are given as arithmetic mean and standard error of mean. The grey marked area displays the normative range (Z-scores between  $+/-1.96$ ). P-values: \* $<0.05$ , \*\* $<0.01$ , \*\*\* $<0.001$ . doi:10.1371/journal.pone.0017387.g002



**Figure 3. The influence of TRPV1 1911 A>G on QST parameters in cluster 1.** The influence of the TRPV1 1911A>G genotype on the somatosensory profile as detected by QST in the patients of cluster 1 is shown. Homozygote variant carriers expressed significant lower, i.e. normal, Z-scores for HPT ( $p=0.0006$ ). The second genotype-affected QST parameter is MDT. The higher Z-scores for MDT, i.e. less mechanical hypaesthesia to non-painful mechanical stimuli, correlated with the presence of G-variant allele in TRPV1 1911A>G ( $p<0.0001$ ). Third, lower Z-scores of mechanical pain sensitivity (MPS), i.e. less pinprick hyperalgesia, were associated with the presence of at least one G-variant allele in TRPV1 1911A>G ( $p=0.005$ ). Values are given as arithmetic mean and standard error of mean. P-values are shown for significant genotype-phenotype associations:  $**<0.01$ ,  $***<0.001$ .

doi:10.1371/journal.pone.0017387.g003


1911A>G and 1103C>G, were identified that had a significant relationship with the somatosensory function in cluster 1 (Table 2).

The TRPV1 1911A>G polymorphism was significantly associated with altered heat pain thresholds (HPT). Neuropathic pain patients being heterozygote or with wild type TRPV1 1911G genotype (AA or AG) tended to show heat hyperalgesia whereas TRPV1 1911G homozygotes exhibited significantly higher, i.e. normal heat pain thresholds ( $p=0.0006$ ; Fig. 3, 4). TRPV1 1911A>G was also identified to be associated significantly to mechanical pain sensitivity (MPS;  $p=0.005$ ). The presence of at least one G-variant allele was associated with lower, i.e. normalized mechanical pain sensitivity to pinprick stimuli (MPS; Fig. 3, 4).

Moreover, TRPV1 1911A>G wild type carriers (AA) showed higher mechanical detection thresholds (MDT), i.e. mechanical hypaesthesia, than variant subjects ( $p<0.0001$ ; Fig. 3, 4).


In contrast to TRPV1 1911A>G, the 1103C>G SNP was significantly associated with cold detection threshold (CDT,  $p=0.0023$ ). Homozygous variant carriers (GG) exhibited cold hypaesthesia compared with heterozygous or wild type carriers ( $p=0.0018$ ; Fig. 5).

Analyses within cluster 2 did not show any genotype-dependent association with any QST parameter (Fig. 6). Due to the small sample size, no analysis of possible associations of the different neuropathic pain syndromes and QST parameters were performed.



**Figure 4. Associations of TRPV1 1911 A>G with HPT, MPS and MDT in cluster 1.** Z-scores of heat pain threshold (HPT), mechanical pain sensitivity (MPS) and mechanical detection threshold (MDT) stratified to TRPV1 1911G>A. **HPT:** Homozygous variant GG carriers exhibited significantly lower, i.e. normal, Z-scores than carriers of the heterozygote or wild type genotype, which tended to show heat hyperalgesia. **MPS:** Presence of at least one G-variant allele resulted in a decreased Z-score indicating less pinprick hyperalgesia. **MDT:** Presence of at least one G-variant allele was associated with less mechanical hypaesthesia to non-painful mechanical stimuli resulting in higher Z-scores. Z-scores are given as median with quartiles and ranges, p-values were calculated with t-test and corrected for multiple testing by permutation testing.

doi:10.1371/journal.pone.0017387.g004



**Figure 5. The effect of TRPV1 1103 C>G on detection of non-painful cold.** The box plots show the cluster 1 specific association of TRPV1 1103C>G SNP with the cold detection threshold (CDT). Homozygous variant carriers exhibited cold hypaesthesia, i.e. lower Z-score, compared with heterozygous or wild type carriers. Z-scores are given as median with quartiles and range. P-values were calculated with t-test and corrected for multiple testing by permutation testing. doi:10.1371/journal.pone.0017387.g005


## Discussion

The major finding of this study is that TRP channel polymorphisms modulate the somatosensory function in neuropathic pain patients. Since there were no significant differences in the allelic distribution between neuropathic pain patients and healthy volunteers it appears unlikely that the variants play a role as susceptibility factors of chronic neuropathic pain.

## Association of TRPA1 710G>A with paradoxical heat sensation

Different mechanisms contribute to paradoxical heat sensation, i.e. the perception of painful heat when a cold stimulus is applied. First, A-delta fiber impairment, clinically manifest in a disturbed cold sensation, is thought to disinhibit C-nociceptors by mechanisms of gate control in the dorsal horn of the spinal cord leading to PHS [14,15]. This hypothesis is supported by observations in the experimental human pain model of selective A-fiber blockade and in patients with peripheral neuropathy [14,16]. Second, dysfunction of thalamic cells leads to disinhibition of central heat transmission pathways, resulting in PHS e.g. in patients with multiple sclerosis [17,18]. The frequency of TRPA1 710G>A variant carriers was higher in PHS cases. Although the functional significance of this polymorphism has not yet been investigated in vitro so far, it can be hypothesized that the amino acid exchange from hydrophilic glutamate to hydrophobic lysine in codon 237 may result in a conformational change of the protein secondary structure. With respect to the clinical observation, a loss of function appears to be likely and thus result in an impairment of cold nociceptor sensitivity. This might lead to a decreased input to the central nociceptive pathway that is the same pathway for the conduction of heat and PHS. This would, according to the hypotheses described above, lead in turn to a lower likelihood of an effective disinhibition and thereby PHS.

Similar to the TRPV1 receptor, TRPA1 is known to sensitize sensory neurons after nerve injury and contribute to altered pain sensation in a familial episodic pain syndrome [19]. Therefore, wild type carriers of the active TRPA1 710 might have a higher propensity for sensitisation of nociceptors peripherally and/or centrally, increasing the likelihood of PHS compared to carries of the variant genotype. Interestingly, a psychophysical study in CRPS patients demonstrated that PHS is a frequent finding in acute CRPS [20] and inflammation is thought to be a dominant cause of sensitisation in acute CRPS patients [21]. Possibly TRPA1 contributes to this proposed mechanism in CRPS, since TRPA1 is activated in models of inflammatory pain and leads to pain and hyperalgesia [11,12].



**Figure 6. The influence of TRPV1 1911 A>G on QST parameters in cluster 2.** The influence of the TRPV1 1911A>G genotype on the somatosensory profile as detected by QST in the patients of cluster 2 is shown. The hereditary impact on somatosensory functions was detectable in cluster 1 only. No significant associations were detected in cluster 2. Values are given as arithmetic mean and standard error of mean. doi:10.1371/journal.pone.0017387.g006

### Association of TRPV1 1911A>G with cold pain threshold

At first sight the association of TRPV1 variants to CPT may sound paradox, since TRPV1 is known as a heat sensor. However, our observation is in line with earlier findings by Kim and coworkers [22] who demonstrated in healthy volunteers that homozygote TRPV1 1911A>G variant carriers exhibited longer cold withdrawal times to ice water immersion of one hand, i.e. higher cold pain tolerance. In our study, the Z-score of the cold pain threshold in 1911GG carriers was lower than in AG and AA carriers ( $-0.13 \pm 1.10$  vs.  $0.19 \pm 1.20$  and  $0.18 \pm 1.14$  respectively) indicating cold hypoalgesia.

TRPV1 1911A>G causes an isoleucine-valine exchange, both hydrophobic amino acids. Though the variant 1911G was associated with elevated protein levels in a human kidney cell line expression system [23], the investigation of functional responses to capsaicin, pH and temperature in whole-cell patch-clamp and calcium imaging experiments suggests no critical alteration of receptor structure and function with respect to this receptor variant [24]. Another explanation for the effect of TRPV1 on cold pain thresholds could be a possible interaction of TRPV1 and the cold sensor TRPA1 since both are co-expressed in sensory neurons [11,12]. However, functional and linkage studies are currently lacking. Therefore our finding requires further confirmation.

### Associations in phenotypic subpopulations

We were able to allocate the phenotypically characterized patients to two different clusters. Patients in a cluster share sensory symptoms and signs and thus may share similar underlying pathological mechanisms [25]. Interestingly, in a study that investigated nerve and skin preparations of patients who suffered from various neuropathic pain syndromes, TRP channels have been found to be differentially expressed [26], thus supporting the approach of stratification. One may hypothesize that in those patients in whom TRPs are still preserved or increasingly expressed SNP polymorphisms contribute to the somatosensory function, whereas those who show a depletion of e.g. nerve growth factor (NGF), the membrane density of TRP channels is likely to be decreased [27]. Interestingly, Facer and coworkers [26] found TRPV1 to be reduced in diabetic neuropathy patients. Accordingly, in cluster 2 where the number of polyneuropathy patients was significantly higher as in cluster 1, we did not find any association between TRP variants and somatosensory function.

As mentioned above, TRPV1 1911A>G has not been shown to alter TRPV1 function in vitro or in vivo [24]. However, in this study it clearly modified different somatosensory functions in patients allocated to cluster 1.

TRPV1 is a heat sensor and contributes to the sensitisation of peripheral nociceptors in neuropathic pain [11,12]. The corresponding clinical sign is heat hyperalgesia [1]. Peripheral sensitisation of TRPV1 sensitive nociceptors in turn leads to sensitisation of the central nociceptive system and clinically to mechanical hyperalgesia [1]. Due to central adaptive mechanisms tactile hypoesthesia accompanies mechanical hyperalgesia [28]. Thus, each sensory finding is compatible with neurobiological and neuropharmacological mechanisms that can also be studied in human surrogate models of neuropathic pain, e.g. capsaicin model that involve a reversible modulation of the nociceptive system [29,30]. These observations are in line with our findings on the TRPV1 1911 genotype. Wild types or heterozygotes showed heat hyperalgesia, mechanical hyperalgesia and mechanical hypoesthesia in comparison to variant carriers that showed normalised sensitivity to heat pain (HPT), mechanical pain detection (MPT) and tactile sensation (MDT) (Figure 3, 4). Thus, the variant

TRPV1 1911A>G may prevent from peripheral and secondary central sensitisation in patients with chronic neuropathic pain.

The second TRPV1 SNP of interest, namely TRPV1 1103C>G was significantly associated with cold hypoesthesia. The SNP causes a methionine-isoleucine exchange and in-vitro investigation demonstrated an increased channel activity [23]. However since no effect of TRPV1 1103C>G on HPT was observed, the question arises if this observation might be explained by a functional interaction of TRPM8 and TRPV1 that leads to increased functional inhibition of the TRPM8 by this SNP [31,32].

### Study limitations

There are three main limiting factors of this study. First, the number of SNP selected for this study was limited to missense variants. No intronic, promoter, 5'- or 3'-UTR variants were included as no association to pain or functional characterization could be shown so far [33]. The functional significance of the missense variants found to be associated with sensory function in our study, has not yet been characterized in vitro. Thus the findings are only based on a genetic association analysis. However, our statistical analysis was conducted conservatively using - among others - permutation corrections that still led to significant associations and thus makes chance findings less likely. Second, these associations need to be cross-validated in larger samples. However, to our knowledge, there is currently no patient cohort available outside the DFNS that used a comparable phenotyping method. Hopefully, within the European Pain Consortium, a second cohort will evolve, making soon further genetic association studies and cross-validation possible. Third, no conclusions on the role of these TRP polymorphisms in other ethnic groups can be drawn, as it was recently demonstrated that African Americans exhibit differential response to the topical application of the TRPV1 agonist in capsaicin compared to three other ethnic groups [34].

### Research and clinical perspectives

Our results demonstrate the need for further research on the functionality of TRP genetic variants on sensory function and nociception. Further on, our results open new avenues for the treatment of neuropathic pain patients. First, specific TRPV1, TRPM8 and TRPA1 antagonists are under research as new therapeutic options, with the analgesic TRPV1 agonist capsaicin already available [35–37]. Second, it has to be clarified how far genetic variants of TRP channels may account for response, non-response or symptom-specific efficacy. This would suggest the subgrouping of patients along clinically relevant markers when they enter clinical trials and follow-up trials [38]. Moreover, variants identified here may serve as detectable risk factors for the course of the disease and gene tests on specific SNPs might serve as predictors of pain qualities in the future.

### Supporting Information

**Table S1** Frequencies of selected TRP channel gene variants in 371 neuropathic pain patients and 253 healthy volunteers. P-values between genotypes of patients and controls are corrected for multiple testing using 100,000 permutations. (DOC)

### Acknowledgments

The authors are indebted to all patients and volunteers participating in this multicenter study. Additionally we thank all members of the DFNS in building up the basis of this study by patient recruitment and assessment.

## Author Contributions

Conceived and designed the experiments: A. Binder RB C. Maier TT RDT A. Berthele SH WM MU IC. Performed the experiments: A. Binder DM C. Maier HF JG VH C. Maihöfner RR AS NÜ MU JZ. Analyzed the

data: A. Binder DM RB C. Maier TT RDT A. Berthele FF JG SH WM C. Maihöfner HR GW JZ IC. Wrote the manuscript: A. Binder DM RB FF IC. Involved in revising the paper: C. Maier TT RDT A. Berthele HF JG SH VH WM C. Maihöfner HR RR AS NÜ MU GW JZ.

## References

- Baron R, Binder A, Wasner G (2010) Neuropathic pain: assessment, understanding and treatment. *Lancet Neurology* 9: 807–819.
- Rolke R, Baron R, Maier C, Tölle TR, Treede RD, et al. (2006) Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* 123: 231–243.
- Maier C, Baron R, Toelle T, Binder A, Birbaumer N, et al. (2010) Quantitative Sensory Testing in the German Research Network on Neuropathic Pain (DFNS): Somatosensory abnormalities in 1236 patients with different neuropathic pain syndromes. *Pain* 150: 439–450.
- Tominaga M (2007) Nociception and TRP channels. *Handb Exp Pharmacol* 179: 489–505.
- Caterina MJ, Schumacher MA, Tominaga M, Rosen TA, Levine JD, et al. (1997) The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389: 816–824.
- Tominaga M, Caterina MJ, Malmberg AB, Rosen TA, Gilbert H, et al. (1998) The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron* 21: 531–543.
- McKemy DD, Neuhauser WM, Julius D (2002) Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* 416: 52–58.
- Peier AM, Moqrich A, Hergarden AC, Reeve AJ, Andersson DA, et al. (2002) A TRP channel that senses cold stimuli and menthol. *Cell* 108: 705–715.
- Bautista DM, Siemens J, Glazer JM, Tsuruda PR, Basbaum AI, et al. (2007) The menthol receptor TRPM8 is the principal detector of environmental cold. *Nature* 448: 204–208.
- Story GM, Peier AM, Reeve AJ, Eid SR, Mosbacher J, et al. (2003) ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures. *Cell* 112: 819–829.
- Cortright DN, Szallasi A (2009) TRP channels and pain. *Curr Pharm Des* 15: 1739–1749.
- Stucky CL, Dubin AE, Jeske NA, Malin SA, McKemy DD, et al. (2009) Roles of transient receptor potential channels in pain. *Brain Res Rev* 60: 2–23.
- Baron R, Tölle TR, Gockel U, Brosz M, Freynhagen R (2009) A cross-sectional cohort survey in 2100 patients with painful diabetic neuropathy and postherpetic neuralgia: Differences in demographic data and sensory symptoms. *Pain* 146: 34–40.
- Yarnitsky D, Ochoa JL (1990) Release of cold-induced burning pain by block of cold-specific afferent input. *Brain* 113: 893–902.
- Susser E, Sprecher E, Yarnitsky D (1999) Paradoxical heat sensation in healthy subjects: peripherally conducted by A delta or C fibres? *Brain* 122: 239–246.
- Ochoa JL, Yarnitsky D (1994) The triple cold syndrome. Cold hyperalgesia, cold hypoesthesia and cold skin in peripheral nerve disease. *Brain* 117: 185–197.
- Craig AD, Bushnell MC (1994) The thermal grill illusion: unmasking the burn of cold pain. *Science* 265: 252–255.
- Hansen C, Hopf HC, Treede RD (1996) Paradoxical heat sensation in patients with multiple sclerosis: evidence for a supraspinal integration of temperature sensation. *Brain* 119: 1729–1736.
- Kremeyer B, Lopera F, Cox JJ, Momin A, Rugiero F, et al. (2010) A gain-of-function mutation in TRPA1 causes familial episodic pain syndrome. *Neuron* 66: 671–680.
- Huge V, Lauchart M, Förderreuther S, Kaufhold W, Valet M, et al. (2008) Interaction of hyperalgesia and sensory loss in complex regional pain syndrome type I (CRPS I). *PLoS One* 3: e2742.
- Birklein F, Schmelz M (2008) Neuropeptides, neurogenic inflammation and complex regional pain syndrome (CRPS). *Neurosci Lett* 437: 199–202.
- Kim H, Neubert JK, San Miguel A, Xu K, Krishnaraju RK, et al. (2004) Genetic influence on variability in human acute experimental pain sensitivity associated with gender, ethnicity and psychological temperament. *Pain* 109: 488–496.
- Xu H, Tian W, Fu Y, Oyama TT, Anderson S, et al. (2007) Functional effects of nonsynonymous polymorphisms in the human TRPV1 gene. *Am J Physiol Renal Physiol* 293: 1865–1876.
- Hayes P, Meadows HJ, Gunthorpe MJ, Harries MH, Duckworth DM, et al. (2000) Cloning and functional expression of a human orthologue of rat vanilloid receptor-1. *Pain* 88: 205–215.
- Jensen TS, Baron R (2003) Translation of symptoms and signs into mechanisms in neuropathic pain. *Pain* 102: 1–8.
- Facer P, Casula MA, Smith GD, Benham CD, Chessell IP, et al. (2007) Differential expression of the capsaicin receptor TRPV1 and related novel receptors TRPV3, TRPV4 and TRPM8 in normal human tissues and changes in traumatic and diabetic neuropathy. *BMC Neurol* 7: 11.
- Pezet S, McMahon SB (2006) Neurotrophins: mediators and modulators of pain. *Annu Rev Neurosci* 29: 507–538.
- Geber C, Magerl W, Fondel R, Fechir M, Rolke R, et al. (2008) Numbness in clinical and experimental pain—a cross-sectional study exploring the mechanisms of reduced tactile function. *Pain* 139: 73–81.
- Woolf CJ, Bennett GJ, Doherty M, Dubner R, Kidd B, et al. (1998) Towards a mechanism-based classification of pain? *Pain* 77: 227–229.
- Klein T, Magerl W, Rolke R, Treede RD (2005) Human surrogate models of neuropathic pain. *Pain* 115: 227–233.
- Okazawa M, Inoue W, Hori A, Hosokawa H, Matsumura K, et al. (2004) Noxious heat receptors present in cold-sensory cells in rats. *Neurosci Lett* 359: 33–36.
- Kobayashi K, Fukuoka T, Obata K, Yamanaka H, Dai Y, et al. (2005) Distinct expression of TRPM8, TRPA1, and TRPV1 mRNAs in rat primary afferent neurons with adelta/c-fibers and colocalization with trk receptors. *J Comp Neurol* 493: 596–606.
- Kim H, Mittal DP, Iadarola MJ, Dionne RA (2006) Genetic predictors for acute experimental cold and heat pain sensitivity in humans. *J Med Genet* 43: e40.
- Wang H, Papoiu AD, Coghill RC, Patel T, Wang N, et al. (2010) Ethnic differences in pain, itch and thermal detection in response to topical capsaicin: African Americans display a notably limited hyperalgesia and neurogenic inflammation. *Br J Dermatol* 162: 1023–1029.
- Premkumar LS (2010) Targeting TRPV1 as an alternative approach to narcotic analgesics to treat chronic pain conditions. *AAPS J* 12: 361–370.
- Backonja M, Wallace MS, Blonsky ER, Cutler BJ, Malan PJR, et al. (2008) NGX-4010 C116 Study Group. NGX-4010, a high-concentration capsaicin patch, for the treatment of postherpetic neuralgia: a randomised, double-blind study. *Lancet Neurol* 7: 1106–1112.
- Bevan S, Andersson DA (2009) TRP channel antagonists for pain—opportunities beyond TRPV1. *Curr Opin Investig Drugs* 10: 655–663.
- Patapoutian A, Tate S, Woolf CJ (2009) Transient receptor potential channels: targeting pain at the source. *Nat Rev Drug Discov* 8: 55–68.