



## Correlates of objectively measured sedentary time in adults with intellectual disabilities

Leanne Harris<sup>a</sup>, Arlene M. McGarty<sup>a</sup>, Thessa Hilgenkamp<sup>b</sup>, Fiona Mitchell<sup>c</sup>, Craig A. Melville<sup>a,\*</sup>

<sup>a</sup> Institute of Health and Wellbeing, College of Medicine, Veterinary and Life Sciences, University of Glasgow, G12 0XH, United Kingdom

<sup>b</sup> Department of General Practice, Erasmus MC University Medical Center Rotterdam, Netherlands

<sup>c</sup> Physical Activity for Health, School of Psychological Sciences and Health, University of Strathclyde, G1 1QE, United Kingdom

### ARTICLE INFO

#### Keywords:

Sedentary behaviour  
Intellectual disabilities

### ABSTRACT

Sedentary behaviour is an independent risk factor for adverse health conditions. Adults with intellectual disabilities spend a high proportion of their day engaged in sedentary behaviour, however, there is limited evidence on potential correlates of objectively measured sedentary behaviour in this population group. In Glasgow, UK from July to September 2017, a secondary analysis of pooled baseline accelerometer data from two randomised controlled trials of lifestyle behaviour change programmes was conducted. Backwards linear regression was used to investigate the associations between demographic, biological, and environmental correlates and objective measure of sedentary behaviour (percentage of time spent sedentary). One-hundred and forty-three participants provided valid accelerometer data. Mean percentage time spent sedentary (adjusted for wear time) was 72.9% [Standard Deviation (SD) = 8.7] per day. In the final model, physical and mental health problems were significantly ( $p < 0.05$ ) associated with increased percentage time spent sedentary. This is the first study to provide evidence on multi-level, demographic, biological, and environmental correlates of objectively measured sedentary behaviour in adults with intellectual disabilities. To inform the development of interventions to modify sedentary behaviours in adults with intellectual disabilities, further research is required including a wide range of socio-ecological correlates.

### 1. Introduction

Prolonged periods of time spent in sedentary behaviour is an independent risk factor for adverse health conditions, such as obesity, type II diabetes, cardiovascular disease (de Rezende et al., 2014; Thorp et al., 2011; Wilmot et al., 2012), and mental-ill health (Hamer et al., 2014; Teychenne et al., 2010). Sedentary behaviour is defined as any waking behaviour with an energy expenditure  $\leq 1.5$  metabolic equivalent, while in a sitting or reclined position, and is a separate construct from physical inactivity (Tremblay et al., 2017).

Epidemiological studies have shown that adults spend a high proportion of their day engaged in sedentary activities, such as television (TV) viewing, screen time, and sitting for occupation and travel (Matthews et al., 2008; Colley et al., 2011; Healy et al., 2011a). Adults with intellectual disabilities [defined as having impairments in intellectual functioning and adaptive behaviour that are present before the age of 18 (American Association on Intellectual and Developmental

Disabilities, 2010)] are consistently regarded as a highly sedentary population (Melville et al., 2017), and have been shown to engage in more sedentary time in comparison to the general population (Schuna et al., 2013). Therefore, reducing sedentary time in adults with intellectual disabilities may improve the health of this population group.

The knowledge base relating to correlates of sedentary behaviour in adults with intellectual disabilities is limited (Melville et al., 2017). Two previous studies have utilised subjective measures of sedentary behaviour. Exploratory bivariate analysis revealed that having obesity was positively correlated with more hours spent watching TV in a large ( $n = 1450$ ) sample of adults with intellectual disabilities from the USA, using baseline data from the Longitudinal Health and Intellectual Disabilities Study (LHIDS; Hsieh et al., 2014). However, this was not replicated in a study of 570 French adults with intellectual disabilities (Mikulovic et al., 2014). In contrast, exploratory bivariate analysis revealed that mean weekly time spent sedentary (use of TV, computers or video games) was lower in overweight participants in comparison to

Abbreviations: CPM, counts per minute; IQR, interquartile range; LHIDS, Longitudinal Health and Intellectual Disabilities Study; SD, Standard Deviation; TV, television

\* Corresponding author at: Institute of Health & Wellbeing, College of Medical Veterinary and Life Sciences, University of Glasgow, Academic Unit for Mental Health & Wellbeing, 1st floor Administrative Building, Gartnavel Royal Hospital, 1055 Great Western Road, Glasgow G12 0XH, United Kingdom.

E-mail addresses: [Leanne.Harris@glasgow.ac.uk](mailto:Leanne.Harris@glasgow.ac.uk) (L. Harris), [Arlene.McGarty@glasgow.ac.uk](mailto:Arlene.McGarty@glasgow.ac.uk) (A.M. McGarty), [t.hilgenkamp@erasmusmc.nl](mailto:t.hilgenkamp@erasmusmc.nl) (T. Hilgenkamp), [fiona.c.mitchell@strath.ac.uk](mailto:fiona.c.mitchell@strath.ac.uk) (F. Mitchell), [Craig.Melville@glasgow.ac.uk](mailto:Craig.Melville@glasgow.ac.uk) (C.A. Melville).

<https://doi.org/10.1016/j.pmedr.2017.11.010>

Received 15 September 2017; Received in revised form 9 November 2017; Accepted 27 November 2017

Available online 29 November 2017

2211-3355/ © 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

non-overweight participants, although this was not statistically significant (Mikulovic et al., 2014).

Two studies have examined the correlates of objectively measured sedentary behaviour in adults with intellectual disabilities. In a UK sample of 62 adults with mild to moderate intellectual disabilities, gender was correlated with sedentary behaviours, with women more sedentary than men (Finlayson et al., 2011). A Norwegian study with a sample of 96 individuals with Down syndrome, William syndrome, or Prader Willi syndrome found that body weight was a correlate of sedentary behaviour, with participants who were underweight or of normal weight more sedentary than individuals who were overweight or obese (Nordström et al., 2013).

Although these initial studies are of interest, they are limited by their focus on individual factors and provide little information on the complex environmental influences on sedentary behaviour. A socio-ecological perspective has been proposed as a useful framework to understand correlates of sedentary behaviour. This multi-level approach includes intrapersonal (biological), interpersonal (social), organizational, environmental, and policy factors (Bronfenbrenner, 1979; Owen et al., 2011). This model has been widely used to investigate how different factors affect sedentary behaviour and activity in typically developing populations (Sallis et al., 2008).

Due to the limited existing research relating to adults with intellectual disabilities, it is important to increase our understanding of multi-level factors which correlate with sedentary behaviour in this population. This can then inform the development of future interventions to decrease the high levels of sedentary behaviours and reduce the health inequalities experienced by this population group (Owen et al., 2011). Furthermore, it is important that the limitations of previous studies measuring sedentary behaviour in adults with intellectual disabilities are addressed and research is conducted with objective measures of sedentary behaviour. Therefore, the aim of this study is to add to the available evidence by investigating correlates of objectively measured sedentary behaviour in adults with intellectual disabilities.

## 2. Method

### 2.1. Design

A secondary analysis of pooled baseline data from two randomised controlled trials (RCTs) of lifestyle behaviour change programmes was conducted. One RCT was focused on weight management ( $n = 50$ ; Harris et al., 2015; Harris et al., 2017) and the second one on increasing physical activity ( $n = 102$ ; Mitchell et al., 2013; Melville et al., 2015).

### 2.2. Participants

Participants ( $n = 152$ ) were recruited for the multi-component lifestyle behaviour change programmes, which were conducted in Glasgow, Scotland, between 2013 and 2014. The eligibility criteria for both studies were similar. Participants over 18 years of age with any level of intellectual disabilities and who were independently ambulatory were included. Full details of these studies have been published previously.

### 2.3. Measures

#### 2.3.1. Objective measure of sedentary behaviour

The baseline accelerometer data of participants was used in this study, to remove any potential influence of the lifestyle behaviour change programmes. Sedentary behaviour was objectively measured using the ActiGraph GT3X + accelerometer (ActiGraph, LLC, Pensacola, FL, USA). This small, lightweight device ( $46 \times 33 \times 15$  mm, 19 g) was worn on the right hip at the iliac crest, attached using an elastic belt, for seven days, except when showering, bathing or swimming. The minimum data requirements for valid accelerometer data was set at 6 h

of data, on at least three days from seven. Non-wear time was defined by intervals of at least 60 min of zero activity counts (Troiano et al., 2008). Activity counts were recorded over 15 second intervals (epochs) and counts for four consecutive epochs summed to give activity counts per minute (cpm). Sedentary behaviour was defined as  $< 100$  cpm based on cut points in the general population (Atkin et al., 2012). Sedentary behaviour is reported as minutes per day and percentage time spent in sedentary behaviour, adjusted for wear time.

#### 2.3.2. Potential correlates of sedentary behaviour

Descriptive data from the primary studies were included as potential correlates of sedentary behaviour, which were categorised into demographic, biological, and environmental factors based on the framework by Sallis et al. (2000).

#### 2.3.3. Demographics

Demographic factors included age (categorised as  $< 45$  years/ $\geq 45$  years), gender (male/female), and level of intellectual disabilities (mild to moderate/severe to profound). Level of intellectual disabilities was assessed based on questions on ability and development in five key areas of functioning: eating and drinking, intimate care, personal safety, communication, and decision-making (The C21 Health check; [https://www.gla.ac.uk/media/media\\_62785\\_en.pdf](https://www.gla.ac.uk/media/media_62785_en.pdf) pages 63–64). Total scores assessed by the ability and development questionnaire have shown to be highly associated (Melville et al., 2008) with the Vineland's Adaptive Behaviour Scale a validated assessment of functioning and ability level (Sparrow et al., 1984). Ethnicity (White/Asian), marital status (married or live-in partner/separated or divorced/single) were also measured but due to categories with very low numbers of data points these factors were not included in the analysis.

#### 2.3.4. Biological

Biological factors included were physical and mental health problems, problem behaviours, and obesity. Health was assessed by whether participants had physical health problems (yes/no), mental health problems (yes/no), or problem behaviours (yes/no), using self- or family/paid carer-proxy-reports. Physical health was assessed using an open-ended question where participants listed any physical health problems, e.g. type II diabetes or high blood pressure. Mental health was assessed based on a yes/no response to the following definitions: any mental health needs, emotional problems, psychological problems, dementia, or other psychiatric ill-health. Problem behaviours were defined as any problem behaviours, challenging behaviour, or special needs related to behaviour, for example verbal aggressive behaviour toward other people, physical behaviour that lead to injury to the individual or others, or destruction of property.

Prevalence of obesity (yes/no) was assessed based on objective measurements of weight and height. Measurements were made with the participant wearing light clothes without shoes. All measurements were made in duplicate and the final value calculated as the mean of the two measurements. Weight in kg was measured to the nearest 100 g using SECA877 scales (SE approval class III; SECA Germany). Height in m was measured to the nearest 1 mm using the SECA Leicester stadiometer (SECA, Germany). BMI ( $\text{kg}/\text{m}^2$ ) was calculated as weight (kg) / height<sup>2</sup> (m). Obesity was defined as a BMI  $\geq 30$   $\text{kg}/\text{m}^2$ .

#### 2.3.5. Environmental

Environmental factors included were accommodation type and neighbourhood deprivation.

Accommodation type was assessed based on a self- or family/paid carer-proxy-reported question asking where the participant lives, which had the following nine potential responses:

- 1) Parents home
- 2) Other family carers home
- 3) Lives independently  $\pm$  children, without any paid support

- 4) Lives independently with paid support
- 5) Supported group living (shared tenancy, with paid support)
- 6) Supported living - individual (single tenancy, with paid support)
- 7) Residential care (registered home)
- 8) Nursing home
- 9) NHS accommodation.

Based on the answer to the question about where the person lived, accommodation type was categorised as: lives with family support (1–2 from the list above), lives independently (3–5), or lives with paid support (6–9).

Neighbourhood deprivation category was assessed based on the Scottish Index of Multiple Deprivation (SIMD) quintile (<http://www.scotland.gov.uk/Topics/Statistics/SIMD>), calculated based on a participant's postcode. This is categorised into five quintiles of deprivation (most deprived 0–20% to least deprived 80–100%).

### 3. Statistical analysis

Descriptive statistics were used to describe the socio-demographic characteristics of adults with intellectual disabilities and sedentary behaviour (minutes/day). Percentage of time spent sedentary was calculated as the dependent outcome, adjusted for wear time [mean (standard deviation; SD)]. Median and interquartile range (IQR) is reported for non-normally distributed data. Simple linear regression was used to examine the relationship between each correlate and percentage of time spent sedentary. Variables from these exploratory bivariate analyses with  $p < 0.25$  were considered to have a potentially significant relevance to sedentary behaviour and were taken forward to multiple linear regression modelling (Bendal and Afifi, 1977). Multiple linear regression modelling was performed using a backwards linear regression method to remove variables that were non-significant ( $p > 0.05$ ). The final multivariate model fit was tested using  $R^2$  and the model was assessed to ensure it met the assumptions of linear regression. All statistical data were analysed using SPSS 21 IBM statistical package (SPSS IBM, New York, NY, USA).

## 4. Results

### 4.1. Participant characteristics

The socio-demographic characteristics are presented in Table 1. One hundred and forty-three participants provided valid accelerometer data from the total recruited sample of 152 adults with intellectual disabilities. Missing data were due to nine participants not meeting the wear criteria of at least six hours per day on three or more days. The participant health characteristics and deprivation levels in this study are similar to a large population-based sample of adults with intellectual disabilities from the same geographical location (Cooper et al., 2007; Cooper et al., 2011).

### 4.2. Sedentary behaviour

Levels of objectively measured sedentary behaviour were high. The median time spent sedentary was 467.5 min/day (IQR = 411.0–542.2). Mean percentage time spent in sedentary behaviour (adjusted for wear time) was 72.9% (SD = 8.7), with a range of 49.2–96.4% (Table 1).

### 4.3. Correlates of objectively measured sedentary behaviour

Seven variables from the bivariate analyses had a  $p$ -value of  $< 0.25$  and were included in the initial multivariate model (age, level of intellectual disabilities, physical health problems, mental health problems, behavioural problems, obesity, and accommodation type); full results of the bivariate analyses are presented in Table 2. Final multivariate backwards regression models revealed that only physical and

**Table 1**

Socio-demographic characteristics of adults with intellectual disabilities in Glasgow, UK from July to September 2017.

Variables	N	N (%)
<b>Demographic</b>		
<b>Age</b>	140	
< 45 years		54 (38.6)
≥ 45 years		86 (61.4)
<b>Gender</b>	143	
Male		69 (48.3)
Female		74 (51.7)
<b>Marital status</b>	143	
Married or live-in partner		5 (3.5)
Separated or divorced		3 (2.1)
Single		135 (94.4)
<b>Ethnicity</b>		
White		141 (98.6)
Asian		2 (1.4)
<b>Level of intellectual disabilities</b>	142	
Mild		69 (48.3)
Moderate		51 (35.7)
Severe		18 (12.6)
Profound		4 (2.8)
<b>Biological</b>		
<b>Physical health problems</b>	142	
Yes		56 (39.2)
No		86 (60.8)
<b>Mental health problems</b>	128	
Yes		48 (33.6)
No		80 (66.4)
<b>Problem behaviours</b>	133	
Yes		39 (27.3)
No		94 (72.7)
<b>Obesity</b>	141	
Normal (18.5–24.9 kg/m <sup>2</sup> )		17 (12.1)
Overweight (25–29.9 kg/m <sup>2</sup> )		23 (16.3)
Obesity (30–39.9 kg/m <sup>2</sup> )		68 (48.2)
Morbid obesity (> 40.0 kg/m <sup>2</sup> )		33 (23.4)
<b>Environmental</b>		
<b>Type of accommodation</b>	143	
Lives independent		41 (28.7)
Family carers		64 (44.8)
Paid carers		38 (26.6)
<b>SIMD</b>	138	
0–20 % most deprived		68 (49.3)
20–40 %		28 (20.3)
40–60 %		29 (21.0)
60–80 %		9 (6.5)
80–100 % least deprived		4 (2.9)
<b>Sedentary behaviour</b>	143	
Time spent sedentary (min/day) Median (IQR)		467.5 (411.0–542.2)
Percentage of time spent sedentary (%) Mean (SD)		72.9 (8.7)
<b>Physical activity</b>	143	
Time spent in light intensity physical activity (min/day) Median (IQR)		141.9 (114.5–173.7)
Percentage of time spent in light intensity physical activity (%) Mean (SD)		22.6 (6.8)
Time spent in MVPA intensity (min/day) Median (IQR)		26.0 (15.9–38.1)
Percentage of time spent in MVPA (%) Median (IQR)		3.8 (2.4–6.1)

SIMD: Scottish Index of Multiple Deprivation; IQR: Interquartile Range; SD: Standard Deviation; MVPA: Moderate to Vigorous Physical Activity.

mental health problems were significantly associated with sedentary behaviour. The fit of the final model (Table 3) explained a low proportion of the variance (8.8%), leaving a high proportion still to be accounted for.

## 5. Discussion

This is the first study to investigate a wide range of demographic, biological, and environmental correlates of objectively measured

**Table 2**  
Bivariate analysis of correlates associated with percentage time spent sedentary Glasgow, UK from July to September 2017.

Variables	B (SE)	$\beta$	p-Value
<b>Demographic</b>			
<i>Age</i>			
≥ 45 years vs < 45 years <sup>a</sup>	2.46 (1.48)	0.14	0.098
<i>Gender</i>			
Male vs female <sup>a</sup>	0.39 (1.46)	0.02	0.789
<i>Level of intellectual disabilities</i>			
Severe to profound vs mild to moderate <sup>a</sup>	2.73 (2.01)	0.11	0.176
<b>Health</b>			
<i>Physical health problems</i>			
Yes vs no <sup>a</sup>	3.03 (1.48)	0.17	0.042
<i>Mental health problems</i>			
Yes vs no <sup>a</sup>	3.74 (1.57)	0.21	0.019
<i>Problem behaviours</i>			
Yes vs no <sup>a</sup>	1.95 (1.64)	0.10	0.238
<i>Obesity</i>			
Yes vs no <sup>a</sup>	3.48 (1.60)	0.18	0.032
<b>Accommodation type</b>			
Family vs independent <sup>a</sup>	- 2.22 (1.45)	- 0.13	0.129
Paid vs independent <sup>a</sup>	2.05 (1.64)	0.10	0.214
<b>Environmental</b>			
<i>SIMD</i>			
20–40 vs most (0–20) <sup>a</sup>	- 0.94 (1.85)	- 0.04	0.612
40–60 vs most (0–20) <sup>a</sup>	- 0.11 (1.83)	- 0.01	0.951
60–80 vs most (0–20) <sup>a</sup>	0.02 (3.01)	< 0.001	0.996
Least (80–100) vs most (0–20) <sup>a</sup>	3.79 (4.42)	0.07	0.393

SIMD: Scottish Index of Multiple Deprivation; SE: Standard Error.

<sup>a</sup> Reference Category.

**Table 3**  
Final multivariate analysis of correlates associated with percentage time spent sedentary Glasgow, UK from July to September 2017.

Variables	B (SE)	$\beta$	p-Value
<i>Physical health problems</i>			
Yes vs no <sup>a</sup>	3.59 (1.58)	0.20	0.025
<i>Mental health problems</i>			
Yes vs no <sup>a</sup>	3.68 (1.61)	0.20	0.024
R <sup>2</sup> /adjusted R <sup>2</sup>	0.09/0.07		

SE: Standard Error.

<sup>a</sup> Reference Category.

sedentary behaviour in adults with intellectual disabilities. The principal findings of this study illustrate that physical and mental health problems are significantly associated with increased sedentary behaviour in adults with intellectual disabilities.

Understanding the relationship between health and sedentary behaviour is of paramount importance for adults with intellectual disabilities due to the increased health inequalities experienced by this population group (Emerson and Baines, 2011; Krahn and Fox, 2014). There is accumulating evidence in the general population illustrating the negative health effects of sedentary behaviour (de Rezende et al., 2014; Hamer et al., 2014; Teychenne et al., 2010; Thorp et al., 2011; Wilmot et al., 2012). However, due to the paucity of research involving adults with intellectual disabilities the effect of sedentary behaviour on health in this population is unknown. Based on the results of this study it is not possible to assess cause and effect i.e. whether physical and/or mental health problems cause increased sedentary behaviour or whether sedentary behaviour causes physical and/or mental health problems. This is a consistent limitation with cross-sectional sedentary behaviour research (Byun et al., 2011).

The mechanisms of the effect of sedentary behaviour on health have not been fully elucidated, however, evidence in the general population demonstrated potential explanations include both physiological and

psychological factors. Sedentary behaviour has shown to elicit a reduction in metabolic activity which leads to increased cardiovascular risk factors, including high levels of circulating blood glucose and triglycerides (Hamilton et al., 2007; Healy et al., 2008). Metabolic risk factors have also shown to be associated with psychological factors and therefore this may also provide an explanation for mental-ill health (Matthews and Kuller, 2002). Moreover, it is reported that sedentary activities such as TV/computer viewing may also evoke a psychological response due to a lack of social interaction and thus hindering the development of social networks (Kraut et al., 1998; Kubey, 1990). This may be applicable to adults with intellectual disabilities who engage in less social/civic activities in comparison to individuals without intellectual disabilities (Emerson et al., 2014), and spend extended periods of time in their household environment (Verdonschot et al., 2009).

Research on sedentary behaviour in adults with intellectual disabilities is in its infancy and, therefore, there is little evidence to compare with the findings of this study. It is surprising that none of the demographic characteristics were significantly associated with sedentary behaviour, as evidence in adults without intellectual disabilities has demonstrated a number of individual factors correlated with sedentary behaviour (e.g. age, gender, body mass index; O'Donoghue et al., 2016). However, the relationship between individual factors and sedentary behaviour in adults with intellectual disabilities has shown to be unclear, with inconsistent findings reported. Finlayson et al. (2011) found that females were more sedentary than males with mild to moderate intellectual disabilities. Whereas, Nordström et al. (2013) reported no gender difference in objectively measured sedentary behaviour in individuals with Down syndrome, Williams syndrome and Prader Willi syndrome. Therefore, future research is necessary to elucidate the relationship between gender and sedentary behaviour in adults with intellectual disabilities.

In the present study, obesity was not significantly correlated with sedentary behaviour in the final model. Studies involving adults with intellectual disabilities have reported contrasting findings. Nordström et al. (2013) reported weight to be a correlate of objectively measured sedentary time. In contrast, Hsieh et al. (2014) and Mikulovic et al. (2014) did not find a significant relationship between weight status and screen time as a proxy for sedentary behaviour at the multivariate level. However, as screen time only represents a small portion of sedentary time, differences between the present study and previous research could be due to differing sedentary-related outcomes (Tremblay et al., 2017). Further research involving multivariate analysis is required to investigate these socio-demographic characteristics and sedentary behaviour in adults with intellectual disabilities.

The results of this study demonstrate no significant associations between environmental correlates (accommodation type and level of deprivation) and percentage time sedentary. This was an unexpected finding as previous research has highlighted that adults with intellectual disabilities experience their environment differently to the general population; for example, facing barriers to accessing transport (Bodde and Seo, 2009), inaccessibility of fitness centers (Heller et al., 2003), and low rates of employment (Siperstein et al., 2013). Given the important role that environmental factors have in understanding and changing sedentary behaviours (O'Donoghue et al., 2016), it is important that future research investigates a wider range of environmental correlates. This is also important from a theoretical perspective as, in the general population, an ecological model has been widely proposed to categorize sedentary behaviour and inform the development of interventions (Owen et al., 2011). However, this model focuses on sedentary behaviour across four domains: leisure time, transport, household, and occupation, yet the applicability of these domains to the lives of adults with intellectual disabilities is unknown. Therefore, understanding environmental correlates of sedentary behaviour specific to adults with intellectual disabilities, and the development of population-specific theoretical frameworks, is essential to inform evidence- and theory-based interventions for this population group.



### 5.1. Strengths and limitations

This study adds to the limited knowledge base on correlates of objectively measured sedentary behaviour in adults with intellectual disabilities. Objectively measured sedentary behaviour has shown to provide a more valid and reliable measure of sedentary time by reducing the risk of recall bias (Healy et al., 2011b). Furthermore, this study also included a more representative sample of adults with intellectual disabilities by including adults with all levels of intellectual disabilities.

A limitation of this study was the multiple correlates included in the analysis were restricted to those collected during baseline data collection of two multi-component behaviour change interventions (Harris et al., 2017; Melville et al., 2015). As a result, a wider investigation into additional interpersonal, social, and organizational factors was not possible. The cross-sectional design of this study precludes any causality. Furthermore, the direction of the association between physical and mental health problems and sedentary behaviour remains unclear. Therefore, further experimental research is needed to distinguish between cause and effect.

### 5.2. Implications for future research

The results of this study highlight that the factors influencing sedentary behaviour in this population group are complex. The included variables in this study explained a relatively small proportion of the variance in sedentary behaviour. Additional factors which were not included in this study may be influential in affecting sedentary behaviour. To implement significant changes in behaviour, there is a need for studies to move away from addressing individual level factors in isolation, and incorporate a broad range of environmental, social and organizational factors (Sallis et al., 2008). Patterns of sedentary time have been shown to be different during weekdays and weekends in the general population (McVeigh et al., 2016) and therefore may have different influencing factors. It is important that further research examines correlates of patterns of sedentary behaviour (i.e. during weekdays/weekends and time of day), and different types of sedentary behaviour, which may have different influences on the lives of adults with intellectual disabilities. Finally, research involving longitudinal studies is required to distinguish the relationship between correlates and determinants of sedentary behaviour to inform the development of evidence-based interventions tailored to the needs of adults with intellectual disabilities.

The results of the present study should, however, be interpreted with caution as it is not possible to assess cause and effect, i.e. whether physical and/or mental health problems cause increased sedentary behaviour or whether sedentary behaviour causes physical and/or mental health problems. Future research involving longitudinal studies to distinguish the relationship between correlates and determinants of sedentary behaviour in adults with intellectual disabilities, and subsequent experimental research, is therefore necessary.

## 6. Conclusion

This is the first study to provide evidence on multi-level, demographic, biological, and environmental correlates of objectively measured sedentary behaviour in adults with intellectual disabilities. Physical and mental health problems were identified as significantly associated with increased sedentary behaviour. These health problems are considered potentially modifiable. Therefore, knowledge of these correlates may be influential in designing interventions to improve the health of this population group and concurrently reduce sedentary behaviour. Further studies are required to examine the effect of additional intrapersonal, interpersonal, environmental, and organizational factors which may affect sedentary behaviour in adults with intellectual disabilities.

## Conflict of interest

None of the authors have any conflicts of interest to declare.

## Acknowledgments

We would like to thank the participants and family and paid carers who took part in the research projects and provided the data for this secondary analysis.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## References

- American Association on Intellectual and Developmental Disabilities, 2010. Intellectual Disability: Definition, Classification and System of Supports, 11th Edition. AAIDD, Washington.
- Atkin, A.J., Gorely, T., Cledes, S.A., et al., 2012. Methods of measurement in epidemiology: sedentary behaviour. *Int. J. Epidemiol.* 41, 1460–1471.
- Bendal, R.B., Afifi, A.A., 1977. Comparison of stopping rules in forward regression. *J. Am. Stat. Assoc.* 72, 46–53.
- Bodde, A.E., Seo, D.C., 2009. A review of social and environmental barriers to physical activity for adults with intellectual disabilities. *Disabil. Health J.* 2, 57–66.
- Bronfenbrenner, U., 1979. *The Ecology of Human Development: Experiments by Design and Nature*. Harvard University Press, Cambridge, MA.
- Byun, W., Dowda, M., Pate, R.R., 2011. Correlates of objectively measured sedentary behaviour in US preschool children. *Pediatrics* 128, 937–945.
- Colley, R.C., Garriguet, D., Janssen, L., et al., 2011. Physical activity of Canadian adults: accelerometer results from the 2007 to 2009 Canadian Health Measures Survey. *Health Rep.* 22, 7–16.
- Cooper, S.A., Smiley, E., Morrison, J., et al., 2007. Mental ill-health in adults with intellectual disabilities: prevalence and associated factors. *Br. J. Psychiatry* 190, 27–35.
- Cooper, S.A., McConnachie, A., Allan, L.M., et al., 2011. Neighbourhood deprivation, health inequalities and service access by adults with intellectual disabilities: a cross-sectional study. *J. Intellect. Disabil. Res.* 55, 313–323.
- Emerson, E., Baines, S., 2011. Health inequalities and people with learning disabilities in the UK. *Tizard Learn. Disabil. Rev.* 16, 42–48.
- Emerson, E., Hutton, C., Robertson, J., et al., 2014. Perceptions of neighbourhood quality, social and civic participation and the self-rated health of British adults with intellectual disability: cross sectional study. *BMC Public Health* 14, 1252–1266.
- Finlayson, J., Turner, A., Granat, M.H., 2011. Measuring the actual levels and patterns of physical activity/inactivity of adults with intellectual disabilities. *J. Appl. Res. Intellect. Disabil.* 24, 508–517.
- Hamer, M., Coombs, N., Stamatakis, E., 2014. Associations between objectively assessed and self-reported sedentary time with mental health in adults: an analysis of data from the health survey for England. *Br. Med. J. Open.* 4, 1–7.
- Hamilton, M.T., Hamilton, D.G., Zderic, T.W., 2007. Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 56, 2655–2667.
- Harris, L., Melville, C., Jones, N., et al., 2015. A single-blind, pilot randomised trial of a weight management intervention for adults with intellectual disabilities and obesity: study protocol. *BMC Pilot Feasibility Stud.* 1, 5–17.
- Harris, L., Hankey, C., Jones, N., et al., 2017. A cluster randomised control trial of a multi-component weight management programme for adults with intellectual disabilities and obesity. *Br. J. Nutr.* 118, 229–240.
- Healy, G.N., Dunstan, D.W., Salmon, J., et al., 2008. Television time and continuous metabolic risk in physically active adults. *Med. Sci. Sports Exerc.* 40, 639–664.
- Healy, G.N., Matthews, C.E., Dunstan, D.W., et al., 2011a. Sedentary time and cardiometabolic biomarkers in US adults: NHANES 2003–06. *Eur. Heart J.* 32, 590–597.
- Healy, G.N., Clark, B.K., Winkler, E.A., et al., 2011b. Measurement of adults' sedentary time in population-based studies. *Am. J. Prev. Med.* 41, 216–227.
- Heller, T., Hsieh, K., Rimmer, J., 2003. Barriers and supports for exercise participation among adults with Down syndrome. *J. Gerontol. Soc. Work.* 38, 161–178.
- Hsieh, K., Rimmer, J.H., Heller, T., 2014. Obesity and associated factors in adults with intellectual disability. *J. Intellect. Disabil. Res.* 58, 851–863.
- Krahn, G.L., Fox, M.H., 2014. Health disparities of adults with intellectual disabilities: what do we know? What do we do? *J. Appl. Res. Intellect. Disabil.* 27, 431–446.
- Kraut, R., Patterson, M., Lundmark, V., et al., 1998. Internet paradox: a social technology that reduces social involvement and psychological well-being? *Am. Psychol.* 53, 1017–1031.
- Kubey, R., 1990. Television and the quality of family life. *Commun. Q.* 38, 312–324.
- Matthews, K.A., Kuller, L.H., 2002. The relationship between psychological risk attributes and the metabolic syndrome in healthy women: antecedent or consequence? *Metabolism* 51, 1573–1577.
- Matthews, C.E., Chen, K.Y., Freedson, P.S., et al., 2008. Amount of time spent in sedentary behaviours in the United States, 2003–2004. *Am. J. Epidemiol.* 167, 875–881.
- McVeigh, J.A., Winkler, E.A., Howie, E.K., et al., 2016. Objectively measured patterns of

- sedentary time and physical activity in young adults of the Raine study cohort. *Int. J. Behav. Nutr. Phys. Act.* 13, 41–53.
- Melville, C.A., Cooper, S.A., Morrison, J., et al., 2008. The prevalence and determinants of obesity in adults with intellectual disabilities. *J. Appl. Res. Intellect. Disabil.* 21, 425–437.
- Melville, C.A., Mitchell, F., Stalker, K., et al., 2015. Effectiveness of a walking programme to support adults with intellectual disabilities to increase physical activity: walk well cluster-randomised controlled trial. *Int. J. Behav. Nutr. Phys. Act.* 12, 125–136.
- Melville, C.A., Oppewal, A., Elinder, L.S., et al., 2017. Definitions, measurement and prevalence of sedentary behaviour in adults with intellectual disabilities—a systematic review. *Prev. Med.* 97, 62–71.
- Mikulovic, J., Vanhelst, J., Salleron, J., et al., 2014. Overweight in intellectually-disabled population: physical, behavioural and psychological characteristics. *Res. Dev. Disabil.* 35, 153–161.
- Mitchell, F., Melville, C., Stalker, K., et al., 2013. Walk well: a randomised controlled trial of a walking intervention for adults with intellectual disabilities: study protocol. *BMC Public Health* 13, 620–673.
- Nordstrøm, M., Hansen, B.H., Paus, B., et al., 2013. Accelerometer-determined physical activity and walking capacity in persons with Down syndrome, Williams syndrome and Prader–Willi syndrome. *Res. Dev. Disabil.* 34, 4395–4403.
- O'Donoghue, G., Perchoux, C., Mensah, K., et al., 2016. A systematic review of correlates of sedentary behaviour in adults aged 18–65 years: a socio-ecological approach. *BMC Public Health* 16, 163–188.
- Owen, N., Sugiyama, T., Eakin, E.E., et al., 2011. Adults' sedentary behaviour: determinants and interventions. *Am. J. Prev. Med.* 41, 189–196.
- de Rezende, L.F.M., Lopes, M.R., Rey-López, J.P., et al., 2014. Sedentary behaviour and health outcomes: an overview of systematic reviews. *PLoS One* 9, 1–7.
- Sallis, J.F., Prochaska, J.J., Taylor, W.C., 2000. A review of correlates of physical activity of children and adolescents. *Med. Sci. Sports Exerc.* 32, 963–975.
- Sallis, J.F., Owen, N., Fisher, E.B., 2008. *Ecological Models of Health Behaviour*. Health Behaviour and Health Education: Theory, Research, and Practice, 5th ed. Jossey-Bass, San Francisco, pp. 43–64.
- Schuna, J.M., Johnson, W.D., Tudor-Locke, C., 2013. Adult self-reported and objectively monitored physical activity and sedentary behaviour: NHANES 2005–2006. *Int. J. Behav. Nutr. Phys. Act.* 10, 126–138.
- Siperstein, G.N., Parker, R.C., Drascher, M., 2013. National snapshot of adults with intellectual disabilities in the labor force. *J. Vocat. Rehabil.* 39, 157–165.
- Sparrow, S.S., Balla, D.A., Cicchetti, D.V., 1984. *Vineland Adaptive Behaviour Scales*. American Guidance Service, Circle Pines, MN.
- Teychenne, M., Ball, K., Salmon, J., 2010. Sedentary behaviour and depression among adults: a review. *Int. J. Behav. Med.* 17, 246–254.
- Thorp, A.A., Owen, N., Neuhaus, M., et al., 2011. Sedentary behaviours and subsequent health outcomes in adults: a systematic review of longitudinal studies, 1996–2011. *Am. J. Prev. Med.* 41, 207–215.
- Tremblay, M.S., Aubert, S., Barnes, J.D., et al., 2017. Sedentary behaviour research network (SBRN)—terminology consensus project process and outcome. *Int. J. Behav. Nutr. Phys. Act.* 14, 75–92.
- Troiano, R.P., Berrigan, D., Dodd, K.W., et al., 2008. Physical activity in the United States measured by accelerometer. *Med. Sci. Sports Exerc.* 40, 181–188.
- Verdonschot, M.M., De Witte, L.P., Reichrath, E., et al., 2009. Community participation of people with an intellectual disability: a review of empirical findings. *J. Intellect. Disabil. Res.* 53, 303–318.
- Wilmot, E.G., Edwardson, C.L., Achana, F.A., et al., 2012. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia* 55, 2895–2905.