



OPEN

Aggregated transfer factor of ^{137}Cs in edible wild plants and its time dependence after the Fukushima Dai-ichi nuclear accident

Momo Takada^{1✉}, Tetsuo Yasutaka¹, Seiji Hayashi², Mai Takagi² & Keiko Tagami³

We obtained the aggregated transfer factor (T_{ag}) for 10 common edible wild plant species (four perennial spermatophytes, bamboo shoot, two tree species, and three perennial pteridophytes) in northeastern Japan. Measurement of T_{ag} was carried out in 2012–2019 and we also used publicly available data for 2012–2019: food monitoring data and total deposition data from an airborne survey. The T_{ag} obtained from actual measurements agreed well with T_{ag} values calculated from the publicly available data. The sampling locations were only identified at the municipal level and uncertainty of the deposition for the publicly available data, and thus T_{ag} values showed substantial variation. The T_{ag} of the perennial spermatophytes, including bamboo shoot, and perennial pteridophytes showed single exponential decline with effective half-lives of approximately 2 years, whereas those of tree species did not show distinct temporal change. These results imply that data since 2014 are applicable for T_{ag} estimation for long-term potential ingestion dose in the future to the public because of the slow decline. The calculated T_{ag} values of all species for 2014–2019 ranged from 6.1×10^{-5} to $5.2 \times 10^{-3} \text{ m}^2/\text{kg}$ -fresh mass. The maximum T_{ag} value was observed for the tree koshiabura (*Chengiopanax sciadophylloides*) and the minimum value was observed for the perennial spermatophyte giant butterbur (*Petasites japonica*). Tree species showed higher T_{ag} than spermatophyte and pteridophyte perennials.

Enhanced radiocesium levels in wild foodstuffs in forests persist for longer time periods than in agricultural crops because of factors such as the small amounts of discharge and recycling within the ecosystem^{1,2}. Although 10 years have passed since TEPCO's Fukushima Dai-ichi nuclear power plant accident (hereafter, the Fukushima accident), the radiocesium concentrations in wild foodstuffs (e.g., wild mushrooms and edible wild plants) often exceed the Japanese limit of 100 Bq/kg for ^{134}Cs and ^{137}Cs on a fresh mass (FM) basis, whereas exceedances of this limit have rarely been reported for agricultural products since 2015³. Edible wild plants, comprising spermatophytes and pteridophytes from which new shoots are consumed as leaf vegetables, are the most familiar wild foods in Japan. The collection of edible wild plants in fields or mountains in spring is a tradition in Japanese culture. Estimation of the potential ingestion dose in the future to the public from the consumption of edible wild plants containing ^{134}Cs and ^{137}Cs is required from a radiation protection point of view. This estimation will improve the quality of life for local residents hoping to restore the enjoyment of eating wild foods⁴.

To evaluate the potential ingestion dose in the future to the public from consumption of edible wild plants, estimation of the radionuclide content in edible wild plants is required, for which use of the aggregated transfer factor (T_{ag}) is appropriate. This is defined as the radionuclide activity concentration in edible parts (Bq/kg) divided by radionuclide deposition in the soil (Bq/m²). T_{ag} is commonly applied to estimate radionuclide concentrations in foodstuffs such as berries, mushrooms, and game harvested in natural or semi-natural ecosystems including upland areas or forests⁵. Previous studies have compiled T_{ag} data for trees, berries, mushrooms, and game animals^{6–8}. The T_{ag} assumes equilibrium conditions in the environment by definition, but T_{ag} values usually change over time owing to soil fixation processes and radionuclide migration^{2,9}. Few T_{ag} values of Japanese edible wild plants in an equilibrium condition from before the Fukushima accident are available, although decrease in the radiocesium concentration in several edible wild plants has been observed after the accident^{10,11}

¹National Institute of Advanced Industrial Science and Technology, 1-1-1 Higashi, Tsukuba, Ibaraki 305-8567, Japan. ²National Institute for Environmental Studies, 10-2 Fukasaku, Miharu, Fukushima 963-7700, Japan. ³National Institutes for Quantum Science and Technology, 4-9-1 Anagawa, Inage-ku, Chiba 263-8555, Japan. ✉email: momo.takada@aist.go.jp

and, accordingly, the T_{ag} values decreased. To estimate the long-term potential ingestion dose attributable to consumption of edible wild plants, T_{ag} should be calculated using data that do not show large temporal change because high T_{ag} values soon after contamination cause an overestimation of the long-term ingestion dose. Tagami and Uchida¹⁰ examined the temporal change in radiocesium concentration for three perennial edible wild plants from July 2011 to May 2016 in Chiba Prefecture. The decline of the concentrations could be approximated by exponential functions with two components; effective half-lives representing the rapid and slow loss were 103–230 days and 970–3830 days, respectively. Based on these results, T_{ag} values for assessing long-term potential ingestion dose should be estimated from data since 2014. Tagami et al.¹² observed no decreasing trend in radiocesium concentrations in five species (perennial and woody spermatophytes, and pteridophytes) since 2016 when analyzing monitoring results reported from local governments in Fukushima Prefecture, and thus T_{ag} values were calculated with data since 2016.

Following the Fukushima accident, although the T_{ag} of edible wild plants has been studied previously, the number of species analyzed to date remains limited. A far greater number of edible wild plant species are utilized in Japan and about 20 species of edible wild plants have been consumed, according to interview-based surveys of local residents in rural areas of northeastern Japan^{13,14}. The T_{ag} for additional species is required to accurately estimate the potential ingestion dose to the public from consumption of edible wild plants. However, most edible wild plant species are not cultivated in agricultural fields, and finding them requires searching a wide area in fields and mountains on foot. Therefore, for many species, conducting a continuous field survey to directly determine T_{ag} values is difficult.

Food radioactivity has been monitored since the Fukushima accident by the Ministry of Health, Labour and Welfare (MHLW)¹⁵. The monitoring data are publicly available and include the radioactivity of various edible wild plants. These data may be applicable for calculation of T_{ag} for a variety of edible wild plant species. To calculate T_{ag} , soil radioactivity data at each sampling location are required, but the MHLW food monitoring data are not accompanied by the corresponding soil data. We considered total radiocesium deposition data from an airborne survey by the Ministry of Education, Culture, Sports, Science and Technology (MEXT)¹⁶ to be appropriate to calculate T_{ag} , instead of soil contamination data. The reason for this is that radiocesium was intercepted by the forest canopy soon after the accident, and then the radiocesium mostly transferred to the forest floor with time¹⁷. Kato et al. reported that soil radiocesium activity in June–July 2011 agreed well with values to the results of the third airborne monitoring survey of May–July, 2011¹⁸.

In the present study, we compared T_{ag} values of edible wild plants calculated using publicly available data, and those obtained with actual measured samples, to confirm the applicability of the publicly available monitoring data. To determine an appropriate period for estimating the potential long-term ingestion dose resulting from the intake of edible wild plants, we also calculated the effective half-life (T_{eff}) from the publicly available data, taking into account the physical half-life and ecological half-life. The T_{ag} values for the edible wild plant species were calculated according to the T_{eff} , which can be appropriate to estimate the long-term potential ingestion dose resulting from consumption of these edible wild plants.

Methods

Edible wild plants and soil samples for T_{ag} calculation. *Sample collection.* We collected edible wild plants and soil samples, and calculated T_{ag} for three different parts of two deciduous perennial spermatophytes [giant butterbur (limb and petiole of *Petasites japonicas*), butterbur scape (flower bud of *Petasites japonicas*), and udo (*Aralia cordata*)], two deciduous trees [fatsia sprout (*Aralia elata*) and koshiabura (*Chengiopanax sciadophylloides*)], and one deciduous pteridophyte perennial [western bracken fern (*Pteridium aquilinum*)]. The sampling locations were located on the campus of the National Institutes for Quantum and Radiological Science and Technology in Chiba Prefecture (about 220 km south of the Fukushima Dai-ichi Nuclear Power Plant) for giant butterbur and butterbur scape in 2012–2019, and in fields and mountains in Kawamata town and Iitate village, Fukushima Prefecture (about 40 km northwest of the power plant) for udo, fatsia sprout, koshiabura, and western bracken fern in 2018–2019. The names of the plants in the present study are based on traditional distinctions in the regions, not biological classification, and we identified the samples with local people in the collection areas. For the fields and mountains in Kawamata town and Iitate village, we obtained permission from the landowners to enter the fields for collection, and we worked with the residents to collect wild plants. Litter and soil samples were also collected separately at each sampling location for the edible wild plants, from 20 × 20-cm areas and with a core sampler (5 cm diameter and 5 cm depth). The Forestry Agency reported that 75–88% of soil radiocesium was observed in 0–5 cm layers in six forests in Fukushima prefecture in 2019¹⁹. In 10 undisturbed croplands in eastern Japan, about 70% of ¹³⁴Cs in soil was kept in 0–5 cm layers in 2018²⁰. Accordingly, we used ¹³⁷Cs concentration data in soil 0–5 cm as soil deposition in the present study. Litter and soil samples were collected from 2–3 points at each sampling site.

Measurement. The collected samples of the edible wild plants were immediately transported to the laboratory and the fresh mass was recorded. All samples were oven dried at 80 °C, the dried weight was recorded, and the samples were crushed. Each sample was mixed well and enclosed in a 100 ml plastic container for radioactivity measurement. The litter and soil samples were also oven dried at 80 °C and enclosed in 100 ml plastic containers. The radioactivity of each sample was measured using a HPGc detector (Seiko EG&G, Japan). Radioactivity was measured for as long as 200,000 s, depending on the radioactivity of the samples. Radiocesium activity concentrations (Bq/kg-FM) of the edible wild plants were derived on a FM basis from the ratio between the fresh and dry masses. Deposition densities of radiocesium (Bq/m²) were combined for litter and soil samples because the litter layer still contains negligible amounts of radiocesium¹⁷.

Publicly available data. *Radiocesium activity in edible wild plants from food monitoring data.* Data for edible wild plants were downloaded from the MHLW food monitoring data¹⁵ on 28 June, 2019. The data were recorded from both cultivated and wild material. We excluded data for cultivated plants from the present analysis. Twelve parts of 11 species of edible wild plants were included as more than 300 measurements were recorded in the period 2011–2019: five deciduous perennial spermatophytes [giant butterbur, butterbur scape, udo, uwabamisou (*Elatostema umbellatum*), momijigasa (*Parasenecio delphinifolius*), and ohbagiboushi (*Hosta sieboldiana*)], bamboo shoot (*Phyllostachys* spp.), two deciduous trees (fatsia sprout and koshiabura), and three deciduous perennial pteridophytes [western bracken fern, ostrich fern (*Matteuccia struthiopteris*), and Japanese royal fern (*Osmunda japonica*)]. These 11 species are indicated to be common edible wild plants in the study region based on the large number of records. However, ohbagiboushi was excluded from the calculation because only 28 radioactivity detections were recorded in the total of ~500 measurements. Data for 2012–2019 were used for the present analysis because of the low number of measurements in 2011. As the season for collecting some edible wild plants starts in December and continues until July in the following year, the data for a specific year comprised those from December of the previous year to July of the current year. The monitoring data contained the sampling location at the municipal level, but more detailed location data were not identified. We selected the monitoring data collected at the municipalities where radiocesium deposition data were monitored. We calculated T_{ag} for the edible wild plants on a FM basis (m^2/kg -FM) in the present study. Most T_{ag} estimates for forest products, such as trees, mushrooms, and berries, have been evaluated on a dry-weight basis in previous studies⁶ and in international literature^{5,21}. However, the radioactivity concentrations of the present monitoring data were recorded on a FM basis (Bq/kg-FM), but the water content of the samples was not recorded.

Radiocesium deposition data from airborne survey. The total radiocesium deposition data were obtained from the results of the fifth airborne monitoring survey of 28 July, 2012 conducted by MEXT¹⁶. The measurement value is the average of measurements within a circle of about 600 m diameter under the aircraft, with a flight path width of 1.85 km within 80 km of the nuclear plant and 3 km outside the 80 km. Our obtained data comprised radiocesium (¹³⁴Cs and ¹³⁷Cs) deposition densities at the center points of the sampling squares (approximately 250 × 250 m). The results of only one monitoring event, on 28 July, 2012, were adapted for the entire period of the food monitoring data in the present study, although airborne monitoring surveys by MEXT were conducted several times. The residence half-time of the radiocesium in agricultural soil in Japan is approximately 15 years²², and radiocesium discharge into rivers from catchments was small after the Fukushima accident²³. Thus, only one survey for deposition data could be used for food monitoring data in 2012–2019.

We used a representative deposition value for each municipality because sampling locations of the edible wild plants from the food monitoring data were determined at the municipal level. A geometric mean value of the deposition densities was used as the representative value for each municipality as the deposition densities in each municipality showed log-normal distributions rather normal distributions. For areas where the deposition density was less than 10,000 Bq/m², we used the half value, 5000 Bq/m², because the data indicated “< 10,000 Bq/m²” uniformly. The municipalities in which more than 30% of the area had a deposition density less than 10,000 Bq/m² were excluded from the present analysis because of the large uncertainty in the deposition. The final area for the present analysis comprised 95 municipalities in seven prefectures (Iwate, Miyagi, Fukushima, Ibaraki, Tochigi, Gunma, and Chiba).

Data used for T_{ag} calculation. The data for the 11 parts of 10 species used for T_{ag} calculation are summarized in Table 1. All detected samples for the publicly available data were used in the T_{ag} calculation. The detected sample number accounted for 18–95% of the total number of measurements; for most samples, the reported results were less than the detection limit. Data less than the detection limit were excluded for T_{ag} calculation from publicly available data, which avoids underestimation of ingestion dose although there is the possibility of overestimation of T_{ag} .

Analysis. We calculated the T_{ag} for ¹³⁷Cs because ¹³⁷Cs has a long half-life ($T_{1/2}$ = 30 years) and therefore is suitable for long-term dose assessment. Radioactivity of ¹³⁷Cs for all samples was decay-corrected to 11 March, 2011. The aggregated transfer factor (T_{ag}) was calculated in accordance with IAEA-TRS 472⁵:

$$T_{ag} (m^2/kg - FM) = {}^{137}\text{Cs in edible part (Bq/kg - FM)} / {}^{137}\text{Cs deposition (Bq/m}^2\text{)}. \quad (1)$$

To determine an appropriate period for estimation of long-term potential ingestion dose rate, we also calculated the effective half-life (T_{eff}), which takes into account the physical half-life and ecological half-life. T_{eff} was determined using temporal change in ¹³⁷Cs concentrations normalized by the deposition that is temporal change in T_{ag} value in the present study, which is defined as

$$T_{eff} = \ln(2)/\lambda, \quad (2)$$

where λ is the ¹³⁷Cs loss rate in edible parts of the plants. The value of λ was obtained using the following equation:

$$A_t = A_0 \times \exp(-\lambda \times t), \quad (3)$$

where A_t is T_{ag} of ¹³⁷Cs at time t , and A_0 is estimated initial T_{ag} . Temporal change in T_{ag} under a non-equilibrium condition is adequately described by a two-component exponential decline, comprising rapid and slow loss components⁵. In the present study, we adopted a single exponential equation in accordance with a previous study; Tagami and Uchida¹⁰ reported that the short effective half-life for three edible wild plant species of deciduous perennial spermatophytes was 103–230 days. Given that the present data were for the period 2012–2019, the

Vernacular name	Scientific name	Plant category	Publicly available data ^a				Measured data ^a				
			No. of measurements	Percentage of measurements above the detection limit (%)	No. of prefectures/municipalities	Concentration range (Bq/kg-FM)	Corresponding soil deposition range (kBq/m ²)	No. of measurements	Location and sampling period	Concentration range (Bq/kg-FM)	Corresponding soil deposition range (kBq/m ²)
Giant butterbur	<i>Petasites japonicus</i>	Spermatophyte, deciduous perennial	707	18	7/63	ND-156	12.2–7.34 × 10 ²	44	QST ^b (Chiba), 2012–2018	0.4–8.4	9.2–14.8
Butterbur scape	<i>Petasites japonicus</i>	Spermatophyte, deciduous perennial	536	29	5/59	ND-324	12.2–6.87 × 10 ²	22	QST (Chiba), 2012–2019	0.8–23.2	9.2–14.8
Udo	<i>Aralia cordata</i>	Spermatophyte, deciduous perennial	383	18	4/53	ND-332	14.0–7.34 × 10 ²	2	Kawamata ^c (Fukushima), 2018–2019	6.8–41.6	33.9–54.7
Uwabamisou	<i>Elatostema umbellatum</i>	Spermatophyte, deciduous perennial	149	70	3/29	ND-435	12.6–1.51 × 10 ²				
Momijigasa	<i>Parasenecio delphinijolius</i>	Spermatophyte, deciduous perennial	100	57	4/22	ND-151	12.2–1.21 × 10 ²				
Bamboo shoot	<i>Phyllostachys</i> spp.	Spermatophyte	3596	69	7/74	ND - 737	12.2–6.87 × 10 ²				
Fatsia sprout	<i>Aralia elata</i>	Spermatophyte, deciduous tree	501	49	7/62	ND-898	12.2–6.87 × 10 ²	3	Kawamata (Fukushima), 2019	29.6–2.25 × 10 ²	1.30 × 10 ² –2.87 × 10 ²
Koshiabura	<i>Chengio-panax sciadophylloides</i>	Spermatophyte, deciduous tree	85	95	6/56	ND-7.86 × 10 ³	12.2–6.87 × 10 ²	7	Kawamata, Iitate ^d (Fukushima), 2018–2019	6.07 × 10 ² –1.43 × 10 ⁴	78.8–8.65 × 10 ²
Western bracken fern	<i>Pteridium aquilinum</i>	Pteridophyte, deciduous perennial	1109	38	7/61	ND-452	12.2–6.87 × 10 ²	2	Kawamata (Fukushima), 2018–2019	2.9–22.1	76.5–3.57 × 10 ²
Ostrich fern	<i>Matteuccia struthiopteris</i>	Pteridophyte, deciduous perennial	265	49	5/52	ND-412	12.2–6.87 × 10 ²				
Japanese royal fern	<i>Osmunda japonica</i>	Pteridophyte, deciduous perennial	73	64	6/28	ND-2.99 × 10 ³	14.0–6.87 × 10 ²				

Table 1. Data used for calculation of the aggregated transfer factor of 11 parts of 10 edible wild plant species in 2012–2019. ^aRadioactivity is for ¹³⁷Cs and is decay-corrected to the day of sample collection. ^bNational Institutes for Quantum and Radiological Science and Technology (QST) in Chiba Prefecture, about 220 km south of the Fukushima Dai-ichi Nuclear Power Plant. ^cKawamata town in Fukushima Prefecture, about 40 km northwest of the power plant. ^dIitate village in Fukushima Prefecture, about 40 km northwest of the power plant.

effect of the rapid loss component was considered to be small. Parameter estimation in the single exponential model was conducted using the `nls()` function in R software 4.0.0²⁴.

As the T_{ag} values showed a log-normal distribution for most species, Spearman's rank correlation test was applied to examine the relationship between ¹³⁷Cs deposition and concentration of ¹³⁷Cs in the edible wild plants. One-way ANOVA and Tukey's post hoc test on log-transformed T_{ag} values were used to evaluate the significance of differences among the species. These analyses were conducted using R software 4.0.0²⁴. Significant differences were determined at a probability level of 0.05.

Consent for publication. All authors have read and agreed to the published version of the manuscript.

Results and discussion

Comparison of T_{ag} calculated from publicly available data and actual measurement data. The calculated T_{ag} (m²/kg-FM) in each year is summarized for each species in Supplemental Table 1:

- The geometric means (GMs) of T_{ag} values calculated using the collected samples ranged from 8.1×10^{-6} to 2.5×10^{-2} m²/kg-FM; the minimum was for western bracken fern in 2019 and the maximum was for koshiabura in 2018 at Kawamata, Fukushima.
- The GMs of T_{ag} values calculated using the publicly available data ranged from 1.6×10^{-5} to 1.2×10^{-2} m²/kg-FM and thus were similar to the actual measurement data. The minimum GM was for udo in 2019 and the maximum was for koshiabura in 2019. The geometric standard deviation (GSD) range was 1.5–4.5.

Annual GMs of T_{ag} values calculated from publicly available data and actual measurement data are compared in Fig. 1. The values for individual years are represented by different points. The T_{ag} values were distributed close




Figure 1. Comparison of annual geometric means of the aggregated transfer factor (T_{ag}) calculated from publicly available data and actual measurement data. Circles, diamonds, and triangles indicate deciduous perennial spermatophytes, deciduous tree spermatophytes, and deciduous perennial pteridophytes, respectively. Values for individual years are represented by different points. Error bars indicate the geometric standard deviation in cases where more than three samples were available.




Figure 2. Correlation between deposition and concentration of ^{137}Cs in three edible wild plants. Circles, diamonds, and triangles indicate butterbur scape, fatsia sprout, and western bracken fern, respectively. The three species are representative deciduous perennial and tree spermatophyte, and deciduous perennial pteridophyte, respectively, in the year of the maximum number of detections.

to the 1:1 line, which suggested that T_{ag} values calculated from the publicly available data generally agreed with those calculated from actual measurements. Hence, an obvious overestimation of T_{ag} from the publicly available data described above was not observed in the present data. We confirmed that T_{ag} calculated from the publicly available food monitoring data and the total deposition data from the airborne survey are reliable surrogates for actual measurement samples. We discuss T_{ag} calculated from the publicly available data hereafter.

Relationship between soil deposition and radioactivity in edible wild plants from publicly available data. We confirmed the relationship between deposition and concentration of ^{137}Cs for the publicly available data for butterbur scape, fatsia sprout, and western bracken fern in a year (Fig. 2), as a representative deciduous perennial and tree spermatophyte, and deciduous perennial pteridophyte, respectively, in the year of the maximum number of detections. Butterbur scape, fatsia sprout, and western bracken fern showed positive




Figure 3. Temporal change in the aggregated transfer factor (T_{ag}) in the period 2012–2019. Circles, diamonds, and triangles indicate deciduous perennial spermatophytes, deciduous tree spermatophytes (including bamboo shoot), and deciduous perennial pteridophytes, respectively. Single exponential fitted lines are shown. Solid lines indicate statistically significant parameters (see Table 2).

significant, nonsignificant, and weak negative significant correlations, respectively (Spearman's rank correlation, butterbur scape, $p=0.001$, $r_s=0.45$; fatsia sprout, $p=0.85$, $r_s=-0.03$; western bracken fern, $p=0.03$, $r_s=-0.21$). Among 29 subdata with more than 20 detections for each species in a year, in addition to the data shown in Fig. 2, 13 showed statistically significant positive correlations (Butterbur scape in 2014 and 2016; bamboo shoot in 2012, and 2014–2019; fatsia sprout in 2013 and 2016; koshiabura in 2013; and ostrich fern in 2012), and western bracken fern in 2017 showed a significant negative correlation. These weak correlations may be affected by uncertainty in the deposition data. We used a representative deposition value for each municipality and the original deposition data grid was of low resolution (see the “Methods” section Radiocesium deposition data from airborne survey). Especially for the cases lacking a clear positive correlation, the degree of radiocesium absorption by edible wild plants was largely different even in the same deposition. Radiocesium uptake by plants in an environment is also affected by other factors (e.g., soil characteristics^{25,26}). The edible wild plants targeted

Vernacular name	A_0 (m ² /kg-FM)			λ (/day)			T_{eff} (year) ^a				
Giant butterbur	7.27×10^{-4}	±	2.95×10^{-4}	*	1.31×10^{-3}	±	5.31×10^{-4}	*	1.5	±	0.6
Butterbur scape	9.59×10^{-4}	±	2.19×10^{-4}	***	8.97×10^{-4}	±	3.12×10^{-4}	**	2.1	±	0.7
Udo	4.28×10^{-4}	±	2.07×10^{-4}	*	2.06×10^{-4}	±	4.24×10^{-4}				
Uwabamisou	8.73×10^{-4}	±	1.83×10^{-4}	***	2.99×10^{-4}	±	1.56×10^{-4}	†	(6.4	±	3.3) ^b
Momijigasa	6.43×10^{-4}	±	2.79×10^{-4}	*	1.31×10^{-4}	±	2.68×10^{-4}				
Bamboo shoot	9.80×10^{-4}	±	6.28×10^{-5}	***	2.27×10^{-4}	±	3.84×10^{-5}	***	8.3	±	1.4
Fatsia sprout	3.74×10^{-4}	±	1.81×10^{-4}	†	-1.14×10^{-4}	±	2.40×10^{-4}				
Koshiabura	1.16×10^{-2}	±	2.45×10^{-3}	***	-3.06×10^{-5}	±	1.47×10^{-4}				
Western bracken fern	1.74×10^{-3}	±	3.26×10^{-4}	***	9.75×10^{-4}	±	1.93×10^{-4}	***	1.9	±	0.4
Ostrich fern	1.97×10^{-3}	±	5.23×10^{-4}	***	1.11×10^{-3}	±	4.02×10^{-4}	**	1.7	±	0.6
Japanese royal fern	7.20×10^{-3}	±	3.72×10^{-3}	†	1.48×10^{-3}	±	7.91×10^{-4}	†	(1.3	±	0.7) ^b

Table 2. Estimated parameters and standard errors for correlations of T_{ag} (m²/kg-FM) in the period 2012–2019 with time (day) calculated using Eq. (3) and effective half-lives [T_{eff} (year)] calculated using Eq. (2) for 11 parts of 10 edible wild plant species. A_0 is estimated initial T_{ag} , and λ (/day) is the ¹³⁷Cs loss rate in edible parts of the plants. *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$, † $p < 0.1$. ^a T_{eff} was calculated only when λ was statistically significant. ^bRough estimation values with nonsignificant ($0.05 < p < 0.1$) parameters.

in the present study were not cultivated but were collected in a variety of environments, such as forests with high organic matter content in the soil and paddy field margins with poorly drained soil high in clay content, although we cannot precisely confirm the growth environment of each species included in the present study.

Temporal change in T_{ag} . The time-dependence of T_{ag} for each species in the period 2012–2019 is shown in Fig. 3. The T_{ag} values of deciduous perennial spermatophytes and pteridophytes showed a decreasing trend with time. Given that the bioavailability of ¹³⁷Cs in the soil in the plant root zone decreased with time, as observed in previous studies^{27,28}, we also observed a decrease in T_{ag} . The T_{ag} of deciduous trees did not show a decreasing trend with time.

After the Chernobyl nuclear accident, radiocesium concentrations in deciduous tree leaves decreased with time owing to the effect of direct deposition at an early stage and the following root uptake effect²⁹, and the T_{ag} of tree leaves decreased accordingly. In previous studies conducted in orchards after the Chernobyl and Fukushima accidents, radiocesium concentrations in deciduous tree leaves showed a decreasing trend^{30,31}. The lack of a declining trend for woody edible wild plants T_{ag} in the present study may be due to a smaller effect of direct deposition at the early stage resulting from interception by tall tree canopies in the vicinity. The height of trees with edible wild plants is usually at eye level. The samples collected soon after the accident were possibly affected by direct deposition, whereas in the latter study period, many of the data were from trees grown after the accident. If the effect of direct deposition was large, a declining trend in T_{ag} might have been observed as observed in orchards. Thus, the absence of a declining trend in T_{ag} indicates that the effect of direct deposition was relatively small.

As an additional possibility for the absence of a declining trend in tree T_{ag} , the continuous supply of bioavailable radiocesium from the organic layer on the forest floor may affect the temporal change in T_{ag} . Compared with the managed conditions in orchards of previous studies^{30,31}, an organic layer develops on the soil surface in a forest and, therefore, reabsorption of radiocesium from the organic layer via the roots may be more active. Imamura et al.¹⁷ also observed a similar trend to that in the present study, namely that radiocesium concentrations in leaves of the canopies of the deciduous tree konara oak (*Quercus serrata*) did not show a temporal change from 2011 to 2015 in two Fukushima forests. These authors' results included the effect of direct deposition on the tree bodies at an early stage of the accident, although the emergence of leaves was after the deposition. Nevertheless, a clear decreasing trend in the radiocesium concentration was not observed, which implies that a deciduous tree actively absorbs radiocesium via the roots in Fukushima forests, and a sufficient amount of radiocesium is absorbed to conceal a decline at an early stage owing to the effect of direct deposition.

Single exponential fitted lines for each species are shown in Fig. 3. The estimated parameters and the T_{eff} (year) calculated with Eq. (2) in “Methods” section are presented in Table 2. The T_{eff} for T_{ag} values that showed a decreasing trend was approximately 2 years, except for bamboo shoot. Tagami and Uchida¹⁰ reported that the T_{eff} of the slow loss component for three edible wild plants of deciduous perennial spermatophytes was 970–3830 days. The ¹³⁷Cs decline in pteridophytes, and deciduous shrub and herbaceous species on the floor of European forests was reported to be 1.2–8 years for T_{eff} excluding the rapid loss component after the Chernobyl nuclear accident³². The present results are thus within the range of previous studies.

For bamboo shoot, applying a single exponential function, a relatively long T_{eff} of 8.3 years was estimated. The T_{ag} decreased between 2012 and 2014, and thereafter no notable change was observed. This observation may reflect the effect of rapid and a slow loss components. Indeed, we applied a two-component exponential function for bamboo shoot, and observed T_{eff} of 0.7 years and –7.8 years for the rapid and slow loss components, respectively. For edible wild tree species, statistically significant single exponential fitted lines were not observed, which reflected the absence of change in T_{ag} with time, as discussed above in this section.

Vernacular name	Plant category	T_{ag} (m ² /kg-FM)					Comparison with previous studies	
		N	GM*	GSD	Minimum	Maximum		
Koshiabura	Spermatophyte, deciduous tree	22	5.2×10^{-3}	a	3.5	5.4×10^{-4}	5.3×10^{-2}	$7.3 \times 10^{-3,12}$
								7.2×10^{-2} on a dry-weight basis ³³
Fatsia sprout	Spermatophyte, deciduous tree	115	4.3×10^{-4}	b	4.0	1.1×10^{-5}	1.1×10^{-2}	$1.1 \times 10^{-3,12}$
Japanese royal fern	Pteridophyte, deciduous perennial	24	4.2×10^{-4}	bc	3.9	5.1×10^{-5}	5.0×10^{-3}	1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰
Bamboo shoot	Spermatophyte	2170	3.9×10^{-4}	b	2.5	2.8×10^{-5}	6.9×10^{-3}	$5.1 \times 10^{-4,12}$
Uwabamisou	Spermatophyte, deciduous perennial	69	3.6×10^{-4}	b	2.1	6.6×10^{-5}	1.6×10^{-3}	5.3×10^{-5} – 1.6×10^{-4} for deciduous spermatophyte perennials ¹⁰
Momijigasa	Spermatophyte, deciduous perennial	40	2.5×10^{-4}	bcd	2.5	7.8×10^{-5}	5.8×10^{-3}	1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰
Ostrich fern	Pteridophyte, deciduous perennial	52	2.0×10^{-4}	cde	2.8	2.3×10^{-5}	4.4×10^{-3}	1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰
Western bracken fern	Pteridophyte, deciduous perennial	329	1.9×10^{-4}	de	3.3	1.3×10^{-5}	2.7×10^{-3}	1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰
Butterbur scape	Spermatophyte, deciduous perennial	96	1.7×10^{-4}	de	2.2	3.0×10^{-5}	1.7×10^{-3}	5.3×10^{-5} – 1.6×10^{-4} for deciduous spermatophyte perennials ¹⁰
Udo	Spermatophyte, deciduous perennial	40	1.1×10^{-4}	e	4.0	9.0×10^{-6}	3.7×10^{-3}	1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰
Giant butterbur	Spermatophyte, deciduous perennial	69	6.1×10^{-5}	f	3.1	9.2×10^{-6}	1.9×10^{-3}	$1.4 \times 10^{-4, 12}$
								5.3×10^{-5} – 1.6×10^{-4} for deciduous spermatophyte perennials ¹⁰
								1.4×10^{-4} – 7.0×10^{-4} for spermatophyte and pteridophyte perennials ¹⁰

Table 3. Aggregated transfer factor (m²/kg-FM) calculated from publicly available data for 2014–2019 for 11 parts of 10 edible wild plant species. T_{ag} , Aggregated transfer factor; GM, Geometric mean; GSD, geometric standard deviation. *Different lower-case letters indicate a significant difference (one-way ANOVA with Tukey's post hoc test, $p < 0.05$).

The T_{ag} varied for all species, varying by 1–3 orders of magnitude within a year that included more than two detections (Fig. 3, Supplemental Table 1). As demonstrated in previous studies⁵, the present study also showed substantial variation in T_{ag} values, which may be for several reasons. Recently, Tagami et al.¹² calculated T_{ag} using the radiocesium concentration in edible wild plants measured by local municipalities from higher-resolution publicly available data (accurate to district level) for giant butterbur, bamboo shoot, fatsia sprout, and koshiabura. The municipalities in these authors' study are located within the present study area. These authors' results differed in being one or two orders of magnitude smaller than the present results. The lower resolution of the present deposition data may be one of the causes of the greater T_{ag} variation. The other source of variation is the site dependency of radiocesium absorption by edible wild plants from the soil as described above. Clarification of factors that contribute to the variation in T_{ag} other than ¹³⁷Cs deposition, and its trends consistent with species, is necessary, which will decrease uncertainty and lead to more accurate estimation of T_{ag} of ¹³⁷Cs with wild plants.

Summary of T_{ag} for estimation of long-term ingestion dose to the public. To estimate long-term potential ingestion dose to the public, T_{ag} with small temporal variability excluding high values at the early stage after the accident is required. However, for the edible wild plant species in the present study, no T_{ag} information in an equilibrium condition from before the Fukushima accident is available. Therefore, average values of T_{ag} for the period after the decrease in T_{ag} has weakened and a certain number of samples is available would be appropriate. The T_{eff} for T_{ag} showing a decreasing trend was approximately 2 years except for bamboo shoot, which has not shown any temporal variation since 2014. The T_{ag} for the other species, udo, uwabamisou, momijigasa, fatsia sprout, koshiabura and Japanese royal fern, has not shown temporal variation throughout 2012–2019 (see the “Results and discussion” section Temporal change in T_{ag}). Therefore, T_{ag} values since 2014 are applicable for estimation of long-term potential ingestion dose to the public. The GMs and GSDs of the T_{ag} values for 2014–2019 for each species are shown in Table 3 listed in order of decreasing GM.

Significant differences in T_{ag} were observed among the species (one-way ANOVA with Tukey's post hoc test, $p < 0.05$). The maximum GM was 5.2×10^{-3} m²/kg-FM for koshiabura. The second-highest GM was that observed for fatsia sprout (4.3×10^{-4} m²/kg-FM). The tree species showed higher T_{ag} values than those of the five spermatophyte perennials (6.1×10^{-5} to 3.6×10^{-4} m²/kg-FM). The T_{ag} for bamboo shoot, which shows characteristics of tree species and herbaceous plants, was intermediate (3.9×10^{-4} m²/kg-FM) between those two plant categories. The T_{ag} values for pteridophytes ranged from 1.9×10^{-4} to 4.3×10^{-4} m²/kg-FM, and thus were similar or slightly higher than those of spermatophyte perennials. These results implied that the T_{ag} of edible wild plants showed certain trends consistent with the plant category.

Similar T_{ag} values have been reported in previous studies (Table 3). Tagami et al.¹² reported T_{ag} values for the tree species koshiabura and fatsia sprout of 7.3×10^{-3} and 1.1×10^{-3} m²/kg-FM, respectively, and values of 5.1×10^{-4} and 1.4×10^{-4} m²/kg-FM for the spermatophyte deciduous perennials bamboo shoot and giant butterbur, respectively. Kiyono et al.³³ reported T_{ag} for koshiabura on a dry-weight basis of 7.2×10^{-2} m²/kg, which was similar to the present result adjusted for the water content of the actual measured samples of 86%–88%.

Tagami and Uchida¹⁰ calculated that the T_{ag} of three species of deciduous spermatophyte perennials ranged from 5.3×10^{-5} to 1.6×10^{-4} m²/kg-FM, and that for five species of spermatophyte and pteridophyte perennials ranged from 1.4×10^{-4} to 7.0×10^{-4} m²/kg-FM. The exceptionally high ¹³⁷Cs concentration in koshiabura has been well documented in the Fukushima region^{34,35}. Several factors have been proposed: the shallow root zone of koshiabura corresponds to a surface layer of forest soil containing a high concentration of bioavailable radiocesium, and the root endophytic bacteria of koshiabura make radiocesium in the soil easier to absorb for koshiabura^{36,37}. The T_{ag} of koshiabura is an order of magnitude higher than that of most other species studied (and two orders of magnitude for giant butterbur). Therefore, consumption of koshiabura contributes to a higher potential ingestion dose compared with those of the other species.

No data are available for T_{ag} evaluated before the Fukushima accident for the present study species. Therefore, we cannot verify whether the obtained T_{ag} values are in the range under an equilibrium condition. Somewhat informative data have been recorded by the Nuclear Regulation Authority, and include measurements of ¹³⁷Cs radioactivity in mugwort (*Artemisia indica* Willd.) and soil collected before the accident in Miyagi Prefecture, which is in the same region as Fukushima³⁸. Mugwort is an edible deciduous spermatophyte perennial of the Asteraceae that grows in fields or mountains, but was not included in the present study. The giant butterbur, butterbur scape, and momijigasa are also deciduous spermatophyte perennials of the Asteraceae. The ¹³⁷Cs activities in 2001–2010 for mugwort and soil were 0.029–0.16 Bq/kg-FM and 108–1455 Bq/m², respectively (decay-corrected to 1 January, 2001). The T_{ag} values derived from the measurement data are generally at a level consistent with these three species in Table 3. The T_{ag} values summarized in 2014–2019 are presumed to be close to those observed under an equilibrium condition, and are suggested to be useful to estimate the long-term potential ingestion dose to the public resulting from consumption of these edible wild plants.

Conclusion

We calculated T_{ag} for 11 parts of 10 common edible wild plants using publicly available data for 2012–2019. Each species shows variation in T_{ag} of about two orders of magnitude in 1 year. The reason may partly be because the sampling locations were identified only at the municipal level and uncertainty in the deposition data. Several species showed trends for decline in T_{ag} and the trends have become relatively weaker since 2014. Variation in T_{ag} of each species was substantial, but averaged T_{ag} values for each species since 2014 are applicable for estimation of long-term potential ingestion dose to the public. The averaged T_{ag} ranged over approximately two orders of magnitude among the 10 species, and to a certain extent trends were consistent with plant category. Tree species tend to show a higher T_{ag} than spermatophyte and pteridophyte perennials, and bamboo shoot T_{ag} is intermediate between that of trees and spermatophyte perennials. Clarification of the factors underlying this variation is necessary, which will lead to accurate T_{ag} calculation.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Received: 21 December 2021; Accepted: 9 March 2022

Published online: 25 March 2022

References

- Sombré, L. *et al.* Long-term radiocesium behaviour in spruce and oak forests. *Sci. Total Environ.* **157**, 59–71. [https://doi.org/10.1016/0048-9697\(94\)90565-7](https://doi.org/10.1016/0048-9697(94)90565-7) (1994).
- Tikhomirov, F. A. & Shcheglov, A. I. Main investigation results on the forest radioecology in the Kyshtym and Chernobyl accident zones. *Sci. Total Environ.* **157**, 45–57. [https://doi.org/10.1016/0048-9697\(94\)90564-9](https://doi.org/10.1016/0048-9697(94)90564-9) (1994).
- Ministry of Agriculture, Forestry and Fisheries. *Results of inspections on radioactivity levels in agricultural products*, https://www.maff.go.jp/e/policies/food_safety/agri_gaiyo.html. Accessed 28 Jan 2021.
- Takada, M., Yasutaka, T., Kanai, Y. & Kuroda, Y. Factors affecting resumption of forest or satoyama usage by former evacuees following their return after the Fukushima Dai-ichi nuclear power plant accident. *Radioprotection* **55**, 325–334. <https://doi.org/10.1051/radiopro/2020085> (2021).
- International Atomic Energy Agency. *Handbook of Parameter Values for the Prediction of Radionuclide Transfer in Terrestrial and Freshwater Environments. Technical Reports Series No. 472.* IAEA, Vienna. https://www-pub.iaea.org/mtcd/publications/pdf/trs472_web.pdf. (2010).
- Calmon, P., Thiry, Y., Zibold, G., Rantavaara, A. & Fesenko, S. Transfer parameter values in temperate forest ecosystems: a review. *J. Environ. Radioact.* **100**, 757–766. <https://doi.org/10.1016/j.jenvrad.2008.11.005> (2009).
- Ohashi, S., Okada, N., Tanaka, A., Nakai, W. & Takano, S. Radial and vertical distributions of radiocesium in tree stems of *Pinus densiflora* and *Quercus serrata* 1.5 y after the Fukushima nuclear disaster. *J. Environ. Radioact.* **134**, 54–60. <https://doi.org/10.1016/j.jenvrad.2014.03.001> (2014).
- Tagami, K., Howard, B. J. & Uchida, S. The time-dependent transfer factor of radiocesium from soil to game animals in Japan after the Fukushima Dai-ichi Nuclear accident. *Environ. Sci. Technol.* **50**, 9424–9431. <https://doi.org/10.1021/acs.est.6b03011> (2016).
- Albers, B. P., Steindl, H., Schimmack, W. & Bunzl, K. Soil-to-plant and plant-to-cow's milk transfer of radiocesium in alpine pastures: significance of seasonal variability. *Chemosphere* **41**, 717–723. [https://doi.org/10.1016/s0045-6535\(99\)00432-4](https://doi.org/10.1016/s0045-6535(99)00432-4) (2000).
- Tagami, K. & Uchida, S. Changes of effective half-lives of ¹³⁷Cs in three herbaceous plants and bioavailable ¹³⁷Cs fraction in soil after the Fukushima nuclear accident. *Appl. Geochem.* **85**, 162–168. <https://doi.org/10.1016/j.apgeochem.2017.01.013> (2017).
- Tagami, K. & Uchida, S. Aggregated transfer factor of radiocesium from soil to wild edible herbaceous plants, In: Bessho, K., Matsumura, H., Miura, T., Tagami, K., Takamiya, K. (ed), *Proceedings of the 19th Workshop on Environmental Radioactivity. KEK Proceedings 2018.* High Energy Accelerator Research Organization, Tsukuba, 194–199 (in Japanese with English abstract). <https://lib-extopc.kek.jp/preprints/PDF/2018/1825/1825007.pdf>. (2018).
- Tagami, K., Takata, M., Yasutaka, T. & Uchida, S. Changes of radiocesium concentrations in wild edible plants with time: Do we find radiocesium concentration decrease in recent years? In: Bessho, K., Matsumura, H., Miura, T., Yoshida, G. (ed), *Proceedings of the 21st Workshop on Environmental Radioactivity. KEK Proceedings 2020.* High Energy Accelerator Research Organization, Tsukuba, 63–68 (in Japanese with English abstract). <https://lib-extopc.kek.jp/preprints/PDF/2020/2025/2025004.pdf>. (2020).

13. Saito, A. Change of carrying capacity and gathering pressure on wild edible plants and mushrooms (original article): A case study in Sawauchi Village, Iwate Prefecture. *For. Econ. Res. Inst.* **59**, 2–16. https://doi.org/10.19013/rinrin.59.3_2 (2006) (in Japanese).
14. Matsuura, T., Hayashi, M., Sugimura, K., Tanaka, N. & Miyamoto, A. Ecosystem services evaluation of harvesting edible wild plants/mushrooms: A case study in Tadami Town, Fukushima Prefecture. *Jpn. J. For. Plann.* **47**, 55–81. https://doi.org/10.20659/jjfp.47.2_55 (2013) (in Japanese).
15. Ministry of Health, Labour and Welfare. *Levels of radioactive materials in foods tested in respective prefectures*. https://www.mhlw.go.jp/english/topics/2011eq/index_food_radioactive.html. Accessed 28 Jan 2021.
16. Japan Atomic Energy Agency. *Airborne monitoring in the distribution survey of radioactive substances*. <https://radioactivity.nsr.go.jp/en/list/307/list-1.html>. Accessed 10 Feb 2022.
17. Imamura, N. *et al.* Temporal changes in the radiocesium distribution in forests over the five years after the Fukushima Daiichi Nuclear Power Plant accident. *Sci. Rep.* **7**, 8179. <https://doi.org/10.1038/s41598-017-08261-x> (2017).
18. Kato, H., Onda, Y., Gao, X., Sanada, Y. & Saito, K. Reconstruction of a Fukushima accident-derived radiocesium fallout map for environmental transfer studies. *J. Environ. Radioact.* **210**, 105996. <https://doi.org/10.1016/j.jenvrad.2019.105996> (2019).
19. Forestry Agency. *Result of a Survey of Radioactive Cesium Distribution in Forest*. https://www.rinya.maff.go.jp/j/kaihatu/jyosen/attach/pdf/R1_jittaihaaku-12.pdf. Accessed 20 April 2021.
20. Tagami, K. & Uchida, S. Radiocaesium vertical distributions changes with time in undisturbed grassland soils observed after the Fukushima Accident, In: Bessho, K., Matsumura, H., Miura, T., Yoshida, G. (ed), *Proceedings of the 21st Workshop on Environmental Radioactivity. KEK Proceedings 2020*. High Energy Accelerator Research Organization, Tsukuba, 63–68 (in Japanese with English abstract). <https://lib-extopc.kek.jp/preprints/PDF/2020/2025/2025004.pdf>. (2020).
21. International Atomic Energy Agency. *Environmental Transfer of Radionuclides in Japan Following the Accident at the Fukushima Daiichi Nuclear Power Plant: Report of Working Group 4 Transfer Processes and Data for Radiological Impact Assessment Subgroup 2 on Fukushima Data*. IAEA-TECDOC-1927, <https://www-pub.iaea.org/MTCD/Publications/PDF/TE-1927web.pdf>. (2020).
22. Komamura, M. *et al.* Long-term monitoring and analysis of ⁹⁰Sr and ¹³⁷Cs concentrations in rice, wheat and soils in Japan from 1959 to 2000. *Technical Report No. 24 of the National Institute for Agro-Environmental Sciences, Tsukuba* (in Japanese with English abstract). <http://agriknowledge.affrc.go.jp/RN/2010724549.pdf>. (2006).
23. Taniguchi, K. *et al.* Transport and redistribution of radiocesium in Fukushima fallout through rivers. *Environ. Sci. Technol.* **53**, 12339–12347 (2019).
24. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, <https://www.R-project.org/>. (2020).
25. Sanchez, A. L. *et al.* High plant uptake of radiocesium from organic soils due to Cs mobility and low soil K content. *Environ. Sci. Technol.* **33**, 2752–2757. <https://doi.org/10.1021/es990058h> (1999).
26. Grytsyuk, N., Arapis, G. & Davydchuk, V. Root uptake of ¹³⁷Cs by natural and semi-natural grasses as a function of texture and moisture of soils. *J. Environ. Radioact.* **85**, 48–58 (2006).
27. Schimmack, W. & Bunzl, K. Migration of radiocesium in two forest soils as obtained from field and column investigations. *Sci. Total Environ.* **116**, 93–107. [https://doi.org/10.1016/0048-9697\(92\)90367-2](https://doi.org/10.1016/0048-9697(92)90367-2) (1992).
28. Takeda, A., Tsukada, H., Nakao, A., Takaku, Y. & Hisamatsu, S. I. Time-dependent changes of phytoavailability of Cs added to allomorphic Andosols in laboratory cultivations and extraction tests. *J. Environ. Radioact.* **122**, 29–36 (2013).
29. Antonopoulos-Domis, M., Clouvas, A. & Gagianas, A. Compartment model for long-term contamination prediction in deciduous fruit trees after a nuclear accident. *Health Phys.* **58**, 737 (1990).
30. Antonopoulos-Domis, M., Clouvas, A. & Gagianas, A. Long term radiocesium contamination of fruit trees following the Chernobyl accident. *Health Phys.* **71**, 910–914 (1996).
31. Kusaba, S., Matsuoka, K., Saito, T., Kihou, N. & Hiraoka, K. Changes in radiocesium concentration in a Japanese chestnut (*Castanea crenata* Sieold & Zucc.) orchard following radioactive fallout. *Soil Sci. Plant Nutr.* **61**, 165–168 (2015).
32. Pröhl, G. *et al.* Ecological half-lives of ⁹⁰Sr and ¹³⁷Cs in terrestrial and aquatic ecosystems. *J. Environ. Radioact.* **91**, 41–72. <https://doi.org/10.1016/j.jenvrad.2006.08.004> (2006).
33. Kiyono, Y., Akama, A., Iwaya, M. & Yoshida, Y. Transfer of radiocesium released in the 2011 Fukushima Daiichi Nuclear Power Station accident to *Eleutherococcus sciadophyloides*, a wild tree that produces edible sprouts. *Bull. For. For. Prod. Res. Inst.* **18**, 195–211. https://doi.org/10.20756/ffpri.18.2_195 (2019).
34. Kiyono, Y. & Akama, A. Radioactive cesium contamination of edible wild plants after the accident at the Fukushima daiichi nuclear power plant. *Jpn. J. For. Environ.* **55**, 113–118. https://doi.org/10.18922/jjfe.55.2_113 (2013).
35. Sugiura, Y., Kanasashi, T., Ogata, Y., Ozawa, H. & Takenaka, C. Radiocesium accumulation properties of *Chengiopanax sciadophyloides*. *J. Environ. Radioact.* **151**, 250–257. <https://doi.org/10.1016/j.jenvrad.2015.10.021> (2016).
36. Muramatsu, Y., Sugiyama, M., Ohno, T., Sato, M. & Sato, M. Transfer of radiocaesium into forest products (2 mechanisms and mitigation measures of radiocesium transfer to plants, Special issue Countermeasures for radiocesium contaminated soil). *Jpn. J. Soil Sci. Plant Nutr.* **85**, 117–120. https://doi.org/10.20710/dojo.85.2_117 (2014) (in Japanese).
37. Yamaji, K. *et al.* Root endophytic bacteria of a ¹³⁷Cs and Mn accumulator plant, *Eleutherococcus sciadophyloides*, increase ¹³⁷Cs and Mn desorption in the soil. *J. Environ. Radioact.* **153**, 112–119. <https://doi.org/10.1016/j.jenvrad.2015.12.015> (2016).
38. Nuclear Regulation Authority. *Environmental Radioactivity and Radiation in Japan*, <https://www.kankyo-hoshano.go.jp/data/database/>. Accessed 10 Feb 2022.

Acknowledgements

We thank the authorities of Kawamata and Iitate, and local residents for their assistance during our field survey. We also thank Drs Mirai Watanabe and Masanori Tamaoki, and Mr Seiichi Takechi at the National Institute for Environmental Studies for their support and useful comments regarding field survey and analysis. We thank Robert McKenzie, Ph.D., from Edanz Group (<https://en-author-services.edanz.com/ac>), for editing a partial draft of this manuscript.

Author contributions

M.T. Collection of the field data, and the data analysis, writing the main manuscript. T.Y.: Conception and design of the work, the data analysis. S.H. and M.T.: Collection of the field data. K. T.: Conception and design of the work, collection of the field data. All authors reviewed the manuscript.

Funding

This study was partly supported by JSPS KAKENHI (Grant Numbers 18H04141 and 19J01605).

Competing interests

The authors declare no competing interests.

Additional information

We comply with relevant guidelines and legislation regarding the sample collection in the present study. The plant species in the study are not endangered. The edible wild plant samples, udo, fatsia sprout, koshiabura, and western bracken fern in 2018–2019 collected in Kawamata town and Iitate village, were collected by local residents as part of their wild foods collection activities, and we do not own the measurement samples. The samples of giant butterbur and butterbur scape in 2012–2019 collected on the campus of QST in Chiba Prefecture are stored in QST. Voucher specimens of plant materials in the present study do not exist.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-09072-5>.

Correspondence and requests for materials should be addressed to M.T.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022