



Preventing and Mitigating Post-Traumatic Stress: A Scoping Review of Resilience Interventions for Military Personnel in Pre Deployment

Zhanying Sun ^{1,2,*}, Jie Song ^{2,*}, Jingru Chen¹, Xueyang Gan¹, Yi Li³, Chen Qiu², Weili Zhang¹, Yuan Gao²

¹Graduate School, Chinese PLA Medical School, Beijing, 100853, People's Republic of China; ²Nursing Department, the First Medical Center, Chinese PLA General Hospital, Beijing, 100853, People's Republic of China; ³Department of Orthopedics, the Forth Medical Center, Chinese PLA General Hospital, Beijing, 100853, People's Republic of China

*These authors contributed equally to this work

Correspondence: Yuan Gao, Email gaoyuanzd@163.com

Purpose: Resilience is considered as a protective factor that can assist individuals to reduce post-traumatic stress reactions. In recent years, armies in many countries have widely implemented resilience training programs before deployment to prevent or reduce post-deployment combat stress reactions. Therefore, this study aims to review what is known about resilience interventions for military personnel in pre deployment.

Methods: Based on Arskey and O'Malley's framework, a scoping review was completed. This review was performed through searching databases including PubMed, Embase, Web of Science, Medline and the Cochrane Library, and screening literature to extract data, finally summarizing the findings.

Results: A total of 25 studies focusing on resilience interventions for military personnel in pre deployment were involved and analyzed using intervention approaches, outcome measures, intervention effects, and so on.

Conclusion: Based on the existing evidence in this review, it is cautiously believed that the resilience intervention program for military personnel before deployment is effective. However, there is no single effective best method even the same type intervention can make different effects in different situations and populations. Therefore, the population differences and context should be fully considered in constructing and implementing program to build military personnel resilience.

Keywords: resilience, military personnel, prevention, post-traumatic stress response

Introduction

Modern military operations are characterized by their particularity, complexity, and uncertainty. Therefore, military personnel usually face a variety of stressors, including live ammunition, bloody terror, disability, and death. Excessive stress can lead to serious mental disorders and cognitive impairment, manifesting as anxiety, fear, suicide or homicide, lack of attention, and loss of behavioral control. Finally, non-combat attrition may appear.^{1,2} Two longitudinal studies on veterans of Operational Iraqi Freedom (OIF) and Operational Enduring Freedom (OEF) have indicated that the incidence of posttraumatic stress disorder (PTSD) is 2 to 3 times higher among those exposed to combat relative to those who did not report significant combat exposure.^{3,4}

Large-scale non-combat attrition significantly affects the ability and efficiency of troops to carry out diversified military tasks, lowering overall combat effectiveness and increasing the economic burden of the military's health care system. The US military healthcare system spends \$3.1 billion annually on treatment for PTSD alone, depression alone, and comorbid PTSD and depression.⁵

Prophylactic intensive training targeting protective factors of traumatic stress has been found to lower the incidence and severity of post-traumatic reactions.^{6,7} Resilience, the ability to resist, adapt, recover, and grow from challenges,⁸ is currently recognized as a protective factor in post-traumatic stress response.⁹ Multiple studies have revealed that individuals with higher resilience have a more stable emotional state and mental health when experiencing stressful events,^{10–12} develop lower stress levels, and are better capable of dealing with challenges and complete tasks.

In a longitudinal study performed by the Canadian Forces,¹³ it was found that post-deployment individuals' resilience was higher than that of undeployed individuals at baseline, and deployment experience significantly improved the resilience of some military personnel. Therefore, it is suggested that stress stimulation may positively impact resilience, providing a feasible basis for resilience training strategies.

Realizing the importance of resilience, the US Department of Defense conducted extensive research on measuring and nurturing resilience and launched the Resilience Program in 2007.^{14,15} This program aims to transform disease-focused care models within the military into a mental health-focused prevention model. Then, various branches of the US military proposed a series of programs to improve the resilience of military personnel, including the Comprehensive Soldier Fitness (CSF), the Air Force Resilience Training (ART), the Operational Stress Control (OSC), the Naval Operational Stress Control (OSC), and the Marine Corps Combat Operations Stress Control (COSOC) program.¹⁶ Many countries around the world have widely taken resilience training as a preventive program rather than a medical program, such as the US military's Master Resilience Training (MRT),¹⁷ and the British Army's Traumatic Risk Management, TRiM.¹⁸

Unfortunately, many of these programs have not been empirically evaluated prior to implementation,¹⁹ and some resilience-building programs may exert little or no positive impact on mental health outcomes for military personnel, with interventions diminishing over time.^{20,21} Based on a systematic review,²² Skeffington et al discovered that no high-quality randomized controlled study confirmed the effectiveness of PTSD prevention strategies in 2013. However, a study by the Irish Defence Forces showed that an effective resilience intervention plan was necessary before deployment.²³ Hourani et al²⁴ surveyed resilience interventions undertaken by forces in the United States, the United Kingdom, and the United Nations, concluding that pre-traumatic exposure strategies and stress reduction training methods were the most promising ways to prevent PTSD in 2011. Leppin et al²⁵ reported that resiliency training programs had a small to moderate impact on enhancing resilience and other mental health outcomes within three months of follow-up in 2014, while the research subjects did not include active military personnel. A recent meta-analysis on randomized controlled trials of resilience interventions for military personnel and emergency responders suggests that cognitive-behavioral interventions and mindfulness training can enhance the resilience of military personnel in pre deployment.²⁶

This study aims to review what is known about resilience interventions for military personnel in pre deployment. Using a scoping approach, the intervention of building military resilience can be discussed and elucidated, which is beneficial for understanding how to enhance resilience and formulate effective military personnel resilience intervention programs. It is of great significance for promoting the resilience building of military personnel, preventing and alleviating post-traumatic stress response, lowering the adverse consequences of post-traumatic stress and strengthening the overall combat effectiveness of troops in future studies.

Materials and Methods

This scoping review was based on the scope review framework proposed by Arksey et al²⁷ and the scope review guidelines issued by Peters et al²⁸ The stages are shown below:

1. Identifying questions
2. Identifying studies
3. Selecting studies
4. Extracting data
5. Collating, summarizing, and reporting results

Stage 1: Identifying Questions

This scoping review was performed to deal with the following questions:

1. What resilience interventions are available for pre deployment military personnel?
2. Which concepts do these interventions use and what theories do they depend on?
3. What are the characteristics, purposes and contents of these resilience interventions?
4. What are the (main) outcomes and effects of these resilience interventions?

Stage 2: Identifying Studies

A systematic search was performed between November 2023 and March 2024. Five databases, namely PubMed, Embase, Web of Science, Medline and the Cochrane Library, were chosen. In terms of each database, there was a detailed research string combining the term “military personnel” or “military” or “army” or “soldier” or “national guard” or “marines” or “navy” or “airforce” or “pilots” or “airman” or “armed forces” or “coastguard” or “sailor” or “submariner” and “Resilience” or “psychological, adaptation” or “psychological, stress” or “psychological, adjustment” or “stress” or “adjustment disorders”. The search was performed within titles, abstracts, and keywords. Furthermore, an additional search was made by snowballing the reference lists of retrieved literature.

Two review authors independently assessed the titles and abstracts of studies which were identified by the literature search following the predefined inclusion criteria. We resolved any disagreement through consensus or, if necessary, by consultation with a third review author. If necessary, the original study author was contacted by Email or telephone to acquire uncertain but vital information for the present study. A total of 1645 papers were retrieved (772 PubMed, 206 Embase, 595 Web of Science, 9 Medline, and 63 The Cochrane Library). Among them, 421 repetitive papers were removed, while 105 studies were included after reading the title and abstract. Finally, 25 studies were involved after reading the full text.

Inclusion Criteria

1. Participants: military personnel of all services and arms, including the National Guard Force and the reserve service.
2. Intervention Characteristics: any specifically resilience-based intervention, irrespective of content, duration, setting, or delivery media in pre deployment.
3. Outcome Measures: resilience (measured with specific resilience scales) and resilience related indicators in mental health, physical health, psychological functioning, and work performance.
4. Study Design: randomized controlled trials (RCTs), cluster-RCTs, cross-over trials, Quasi-RCTs, and controlled trials.
5. Published in English and peer-reviewed

Exclusion Criteria

1. Studies evaluating treatment interventions for established psychological illness.
2. Unable to get full text.

Stage 4: Extracting Data

Selected data were tabulated. Extracted information includes the first author, publication year, location, study design, duration of intervention, sample size and pertinent outcomes (see [Table 1](#)).

Stage 5: Collating, Summarizing, and Reporting Results

Types of Interventions

Cognitive-behavioral therapy-based interventions were performed in six studies; seven studies adopted mindfulness and relaxation-based training; two studies were concentrated on stress inoculation training; four studies used biofeedback-based programmes; two studies applied neuropsychological-based programmes; two studies used psychoeducational-based programmes, and in the final two studies, family-centered resilience interventions were performed (See [Table 2](#) for a brief description of each intervention).

Table 1 Characteristics of Included Studies

Reference	Year	Location	Study Design	Duraion	Sample Size (IG/CG)	Pertinent Outcomes
Adler et al ²⁹	2015	USA	Cluster-RCT	10 weeks	2019 (1019/1000)	Anxiety (GAD-7) Depressive symptoms (PHQ-D) Sleep problems Group Cohesion
Bouchard et al ³⁰	2012	Canada	RCT	3 days	41 (21/20)	Objective stress levels: cortisol; heart rate
Cacioppo et al ³¹	2015	USA	RCT	5 days	1138 (688/450)	Empathy (the Empathic concern sub scale) Generalized trust (Rosenberg's Faith in People Scale) Loneliness (revised UCLA scale) Hardiness Alcohol misuse (TICS)
Chongruksa et al ³²	2015	Thailand	RCT	12 weeks	60 (30/30)	Resilience (THMI: adapted from CD-RISC) Mental well-being (THMI-54) anxiety, social dysfunction, depression (GHQ-28 (Thai))
Cohn and Pakenham ³³	2008	Australia	RCT	45 days	174 (101/73)	Attributional Style (Real Events Attributional Style Questionnaire) Coping Strategies (Brief COPE scales) Questionnaire-12 (GHQ-12)
Crane et al ³⁴	2018	Australia	Cluster-RCT	4 weeks	226 (130/96)	Stressor frequency (23 item list created by the researchers) Depression symptoms (PHQ-8) Anxiety (GAD-7)
Fikretoglu et al ²¹	2019	Canada	Cluster-RCT	9 weeks	2831 (1452/1379)	Psychological distress (K-10) Subjective levels of distress (SUDS) Resilience (CD-RISC) Anxiety (GADS-7) Depression (PHQ) Mental health service use (CAF-MHSUQ)
Fornette et al ³⁵	2014	French	Quasi experimental	11 weeks	21 (10/11)	The Profile of Mood States (POMS) questionnaire The State-Trait Anxiety Inventory
Hourani et al ³⁶	2016	USA	Cluster-RCT	7 months	348 (251/97)	Heart period (HP) Respiratory sinus arrhythmia (RSA) PTSD (PCL-C) Perceived Stress (coping; Perceived Stress Scale)
Hourani et al ³⁷	2017	USA	Cluster-RCT	18 months	891 (469/422)	Perceived stress (Personal Stress Situation) PTSD symptoms (PCL-M)
Jha et al ³⁸	2010	USA	Quasi experimental	8 weeks	48 (31/17)	The Ospan score Positive and negative affect (PANAS)
Johnson et al ³⁹	2014	USA	Cluster-RCT	8 weeks	287 (153/134)	Heart rate. Breathing rate. Neuropeptide Y and norepinephrine levels

Jones et al ⁴⁰	2019	UK	RCT	9 weeks	707 (358/349)	Symptoms of common mental disorder General Health Questionnaire (GHQ) Symptoms of PTSD (Caseness; PC-PTSD-5) Alcohol use (AUDIT-C)
Keynan et al ⁴¹	2019	Israel	RCT	4 weeks	180 (90/90)	Emotional regulation: measured using resilience training on the eStroop task Anxiety (STAI, Hebrew version) Alexithymia (TAS-20, Hebrew version)
Lester et al ⁴²	2012	USA and Japan	Quasi experimental	8 sessions	488 unique families	Brief Symptom Inventory (BSI) the McMaster Family Assessment Device (FAD)
Lester et al ⁴³	2016	USA and Japan	Quasi experimental	8 sessions	2165 unique families	The self-report Brief Symptom Inventory-18 (BSI-18) the McMaster Family Assessment Device (FAD) PTSD (PCL-C)
Lewis et al ⁴⁴	2015	USA	Cluster-RCT	15 minutes	891 (469/422)	PTSD (PCL-C) Heart rate and Respiratory sinus arrhythmia (RSA)
Meland et al ⁴⁵	2015	Norwegian	Quasi experimental	12 months	21	Five Facet Mindfulness Questionnaire (FFMQ) Norwegian Mental Skills Questionnaire for Athletes (NMS-42) A Norwegian version of the Sport and Anxiety Scale
Pyne et al ⁴⁶	2018	USA	Cluster-RCT	12 months	342 (200/142)	PTSD (PCL-M) Heart rate variability (HRV)
Regula et al ⁴⁷	2023	Switzerland	RCT	15 weeks	99 (35/46)	The Primary Appraisal and Secondary Appraisal Questionnaire (PASA) Visual analogue scales (VAS) Positive and negative affect (PANAS) Cortisol Heart Rate
Stanley et al ⁴⁸	2011	USA	Case report	8 weeks	47	Five Facet Mindfulness Questionnaire (FFMQ) Perceived Stress Scale (PSS)
Taylor et al ⁴⁹	2011	USA	RCT	12 days	18 (10/8)	Dissociation (CADSS) PTSD symptomatology (IESR)
Wald et al ⁵⁰	2016	Israel	RCT	6 weeks	719 (363/356)	PTSD prevalence Self reported PTSD (PCL-M) Depression symptoms (PHQ-9)
Wessemann et al ⁵¹	2016	Germany	RCT	1.5 days	67 (36/31)	Current mental state (BSI) PTSD symptoms (PDS, German version)
Williams et al ⁵²	2007	USA	Cluster-RCT	9 weeks	1199 (583/616)	Beck Depression Inventory, Second Edition (BDI-II) Perceived Stress Scale (PSS) Interpersonal Relationship Inventory Sense of Belonging Inventory-Psychological (SOBI-P) Coping Styles Questionnaire (CSQ) Perceived Cohesion Scale (PCS)

Abbreviations: IG, intervention group; CG, control group; GAD, Generalised Anxiety Disorder; PHQ, Patient Health Questionnaire; UCLA, University of California, Los Angeles; TICS, two-item conjoint screen; TMHI, Thai Mental Health Indicator; CD-RISC, Connor Davidson Resilience Scale; GHQ, General Health Questionnaire; Brief-COPE, Brief Coping Orientation to Problems Experienced; K-10, Kessler Psychological Distress scale; SUDS, Subjective Units of Distress Scale; CAF-MHSUQ, Canadian Armed Forces Mental Health Service Use Questionnaire; PCL-C, PTSD Checklist- Civilian Version; PCL-M, PTSD checklist-Military version; AUDIT-C, Alcohol Use Disorders Identification Test; STAI, State-Trait Anxiety Inventory; TAS-20, Toronto Alexithymia Scale.

Table 2 A Brief Description of Interventions

Type	Reference	Population	Intervention	Comparator
Cognitive-behavioral therapy-based interventions	Cacioppo et al ³¹	US Army Soldiers	Social Resilience Training (SRT) was delivered in largely a psycho-educational format with some skills based rehearsal. Then, it was designed to address feelings of isolation and maladaptive social cognition with an emphasis on modifying maladaptive social cognition.	Afghanistan cultural awareness training
	Chongruksa et al ³²	Thai Army Rangers	The used integrated Group Counselling (IGC) was an 'Interactive model of existential therapy incorporating art therapy by drawing mandalas, and CBT practices.	Informational control
	Cohn and Pakenham ³³	Australian Army recruits	Brief cognitive-behavioral program was consisted of two 40-minute sessions of psychoeducation on cognitive restructuring and coping strategies.	Discussion of training experiences
	Fornette et al ³⁵	French Air Force flying school cadets	Cognitive adaptation training involved in automatic mental mode and adaptive mental mode was aimed to enhance metacognitive skills pertaining to the two major types of human cognitive processes.	Basic flying training
	Regula et al ⁴⁷	Switzerland officer cadets	Resilience training (RT) is based on a cognitive behavioral approach and on positive psychology and adopted for the needs of the Swiss Armed Forces to improve skills for managing stressful situations and build resilience to prevent negative stress outcomes.	Diversity management training (DMT)
	Williams et al ⁵²	US Navy recruits	BOOT Camp Survival Training included learning and practicing strategies for coping, and managing emotional responses and stress.	No additional training
Mindfulness-and relaxation-based training	Crane et al ³⁴	Australian Army recruits	Self-reflection training was operationalised by means of guided coping and emotion regulatory self-reflection in order to increase resilience.	Coping skills training
	Fikretoglu et al ²¹	Canadian military recruits	Road to Mental Readiness (R2MR) training was consisted of stress management skills training, psycho-educational instruction and small group based vignette exercise to reinforce skills learned.	Delayed intervention control condition
	Jha et al ³⁸	US Marine Corps reservists	Mindfulness-Based Mind Fitness Training (MMFT) blended mindfulness skills training with concrete applications for the operational environment and information and skills concerning stress, trauma, and resilience.	Assessment only
	Johnson et al ³⁹	US Marine Corps	MMFT	No additional training
	Meland et al ⁴⁵	Norwegian combat aircraft squadron	Mindfulness Training (MT) was consisted of theoretical lectures, guided practice on yoga, body scanning and seated meditation.	Not applied
	Stanley et al ⁴⁸	US Marine Corps reserve	MMFT blended mindfulness skills training with concrete applications for the operational environment and information and skills about stress, trauma, and resilience.	No additional intervention
Stress inoculation training	Taylor et al ⁴⁹	US Navy personnel	Psychological skills training (PST) was consisted of arousal control, mental imagery, goal setting, and positive self-talk.	No treatment control condition
	Hourani et al ³⁶	USMC Marines	Pre-deployment Stress Inoculation Training (PRESIT) consisted of education, skills acquisition and practice in a simulated environment.	"Current best practice" brochure + lecture on stress management
	Hourani et al ³⁷	US Army Soldiers	PRESIT	Stress management training

Biofeedback-based programmes	Bouchard et al ³⁰	Canadian Soldiers	Immersion and Practice of Arousal Control training (ImPACT) was a stress-management training programme with biofeedback. Tactical breathing (TB) involved a process of timed inhalations, holds and exhalations.	Training as usual control condition
	Lewis et al ⁴⁴	US Army Paratroopers	Predeployment stress inoculation therapy (PRESTINT) began with “Battle Breathing” techniques, followed by biofeedback training, ending with a Multimedia Stress Environment exercise.	
	Pyne et al ⁴⁶	US National Guard Soldiers	Heart rate variability biofeedback (HRVB) and Cognitive Bias modification for interpretation (CBM-I). CBM-I intervention shifted a soldier’s interpretive style by presenting an increased proportion of neutral or non-negative interpretations of an ambiguous deployment-related scenario.	Training as usual control condition
	Wessemann et al ⁵¹	German army medical unit personnel	Chaos Driven Situations Management Retrieval System (Charly) refers to a technology-based approach consisting of blended learning, Internal differentiation, simulation techniques and biofeedback.	Training as usual control condition
Neuropsychological-based programmes	Keynan et al ⁴¹	Israeli Soldiers	The theoretical underpinning of amygdala electrical fingerprint neurofeedback (Amyg-EFP-NF) is Cognitive Neurofeedback. Downregulation of the amygdala activity is considered a vital mechanism in emotional control and the programme aiming to achieve this through NF.	Control electroencephalography NF or NoNF
	Wald et al ⁵⁰	Israeli Soldiers	Attention Bias Modification Training (ABMT) proposed that mechanism of anxiety reduction is its targeting of disruptions in a threat monitoring system responsible for focusing on potential threats in the environment.	Attentional Control Training (ACT) and no-training
Psychoeducational-based programmes	Adler et al ²⁹	US Army Soldiers	This resilience training (RT) programme was consisted of cognitive-education-positive psychology facets, including normalising symptomatology, and encouraging optimism and model of adversity-belief-consequence.	Military history lectures
	Jones et al ⁴⁰	UK Royal Air Force recruits	“SPEAR” is a programme which emphasizes social networks, capitalising on personal strengths, managing emotions as well as enhancing awareness of psychological symptoms and learning methods to promote resilience.	Training as usual
Family-centered resilience interventions	Lester et al ⁴²	US military families	Families OverComing Under Stress (FOCUS) project	Not applied
	Lester et al ⁴³	US military families	FOCUS project	Not applied

Effects of Interventions

Cognitive–Behavioral Therapy-Based Interventions

Cognitive Behavior Therapy (CBT) is a psychological intervention to change or reconstruct negative cognition through cognitive and behavioral techniques. The core of CBT is to change irrational negative beliefs and thoughts, which can thus alter an individual's psychology and behavior.⁵³

Cacioppo et al³¹ reported significant improvements in military hardiness, military environment tolerance, loneliness, empathy, and compassion after the SRT. Similarly, Chongruksa et al³² considered that the Rangers' resilience scale (CD-RISC) score was dramatically higher than that of the control group after Integrated Group Counselling (IGC). However, this intervention program adopted Buddhist concepts and may not apply to groups of other religious or cultural beliefs. Cohn and Pakenham³³ found that the intervention group reported less self-blame, better perceived coping, greater positive states of mind and lower psychological distress than the control group at the end of the brief cognitive–behavioral program. Fornette et al³⁵ discovered that cognitive adaptation training changed their stress management practices, while no differences existed in mood and anxiety between the intervention group and the control group. Regula et al⁴⁷ indicated that cortisol increased in both resilience training (RT) and diversity management (DMT) groups during stress, while there was a slighter cortisol increase in the RT group thereafter. The RT group perceived the military stressor as more challenging, exhibiting higher values in motivation and exerting positive impacts than the DMT group. Williams et al⁵¹ pointed out that BOOTSTRAP Recipients were more likely to develop significantly higher cohesion, better perceived problem-solving strategies and social support, and had less anger-based coping strategies than controls.

Mindfulness-and Relaxation-Based Training

Mindfulness refers to focusing on the present moment while paying attention to observing, and describing feelings, thoughts, and sensations. It is a protective factor for physical and mental health.⁵⁴ In 1979, Kabat-Zinn proposed mindfulness-based stress reduction therapy and pioneered mindfulness training in the medical community.⁵⁵

Crane et al³⁴ discovered that relative to the control group, participants in the self-reflection group had improvements in mental health outcomes on depression, anxiety symptoms, and perceived stressor frequency at longer-term follow-up. Nevertheless, no difference was found in depression symptoms at initial follow-up. Fikretoglu et al²¹ conducted a mindfulness-based program, named the Road to Mental Readiness (R2MR), with Canadian military recruits. Then, it was found that no difference existed in levels of depression and resilience at a five-week follow-up or at a nine-week follow-up. Jha et al³⁸ pointed out that marines receiving greater Mindfulness-Based Mind Fitness Training (MMFT) experienced lower levels of negative emotion and higher levels of positive emotion. Johnson et al³⁹ discovered that recipients exhibited greater reactivity (heart rate), enhanced recovery (heart rate), breathing rate after stressful training and lower plasma neuropeptide Y concentration after MMFT. Meland et al⁴⁵ considered that reductions in somatic anxiety associated with performance occurred, and improvements in self-perceived skills related to mindfulness, attention regulation and arousal regulation were observed after mindfulness training. Taylor et al⁴⁹ reported that psychological skill training (PST) did not notably lower symptoms of post traumatic stress versus a control condition; there existed a reduction in symptoms of post traumatic stress at 24 hours, 1 month, and 3 months after completing training. Nevertheless, the difference in recovery between intervention and control was not significant. Stanley et al⁴⁸ carried out MMFT with Marine Corps, and confirmed that spending more time in engaging in MMFT corresponded with greater self-reported mindfulness, and that increases in mindfulness were related to reductions in perceived stress.

Stress Inoculation Training

Stress Inoculation Training (SIT) refers to a psychological training model that aims to provide a form of vaccination-like immunity against possible future stressors. The idea is that early exposure to stressful events of a certain intensity can prevent future negative stress effects.⁵⁶ SIT, proposed by Meichenbaum in 1985, is a three-stage model, including the concept stage, skill acquisition and consolidation stage, and application and follow-through stage. It mainly involves cognitive restructuring, relaxation training, self-guidance, and self-monitoring, and is used to improve negative emotions, enhance stress resistance,⁵⁷ and prevent post-traumatic stress disorder.⁵⁷

Hourani et al³⁶ in 2016 reported that Pre-Deployment Stress Inoculation Training (PRESIT) was not successful at lowering PTSD diagnosis and self-reported Perceived Stress Scale (PSS) scores compared to the control condition. However, when baseline mental health issues were controlled, participants in the control condition were at 6.9 times the risk of a diagnosis of PTSD compared with the intervention group. Nevertheless, Hourani et al³⁷ in 2017 mentioned that PRESIT was not efficient in lowering the number of subsequent clinical diagnosis and symptoms of PTSD and the score on the PSS.

Biofeedback-Based Programmes

Biofeedback involves using electrical mechanical equipment for measuring physiologic processes occurring in a person and subsequently feeding this information back to them with the purpose of developing a greater awareness and capability of controlling changes within their bodies.⁵⁸

Bouchard et al³⁰ considered that the participants in the biofeedback-assisted management training group were less stressed than control group members in stressful first-aid simulation. Additionally, the intervention group had greater anticipated self efficacy than the control group participants. Lewis et al⁴⁴ indicated that symptoms of post traumatic stress were not significantly decreased by their predeployment stress inoculation training (PRESTINT) compared to controls, while experimental subjects showed greater heart rate variability (HRV) during a post-training combat simulation. HRV is considered a protective factor for reducing traumatic stress.⁵⁹ Pyne et al⁴⁶ believed that no observable overall effect of the “heart rate variability biofeedback” program was found on post-deployment symptoms of post traumatic stress, and there was no obvious difference between intervention and the control arms at 12-month post-deployment. Wessemann et al⁵¹ conducted Chaos Driven Situations Management Retrieval Systems (CHARLY) programme with German military medical personnel. The CHARLY group exhibited a significantly lower level of traumatic stress on the PDS scale following deployment relative to the control group.

Neuropsychological-Based Programmes

Keynan et al⁴¹ revealed the specificity of neurofeedback learning to the targeted amygdala electrical fingerprint (Amyg-EFP) signal, resulting in reduced alexithymia and faster emotional stroop. This suggested better stress coping skills following Amyg-EFP-NF relative to controls. Wald et al⁵⁰ conducted attention bias modification training (ABMT) which is an intervention method, where a pre-designed processing mode is used to directly manipulate attentional bias through task scenarios,⁶⁰ and they reported four sessions of ABMT reduced risk for PTSD associated with the no-training condition. In addition, no other between-group differences were found.

Psychoeducational-Based Programmes

Adler et al²⁹ considered that there existed no obvious differences between conditions concerning depression symptoms, anxiety symptoms, or sleep problems, and resilience training was rated as less useful than military history classes by participants. Jones et al⁴⁰ reported that levels of PTSD, common mental disorder symptoms, alcohol misuse, help-seeking and homesickness were not of significant difference between groups at post intervention or at three-month follow-up.

Family-Centered Resilience Interventions

Family and social support is a protective factor of resilience.⁶¹ The Mental Health Task Force of the US Department of Defense recognizes that maintaining the overall combat effectiveness of the force requires support for the resilience of military personnel and their families. In 2008, the US Marine Corps launched the Families Overcoming Under Stress (FOCUS) project, enhancing communication and understanding among family members through structured narrative to cultivate family resilience and cohesion.⁶² Lester et al⁴² conducted the FOCUS project, and the results indicated that the parents who participated in the project had significantly lower stress, anxiety, depression index, and Family Assessment Device (FAD) unhealthy family function score. In addition, Lester et al⁴³ also suggested that FOCUS training before deployment vastly alleviated parents’ anxiety and depression symptoms and positive coping skills, and improved the overall resilience of the family.

Conclusion and Expectation

By reviewing previous studies, it is evident that the contents and methods of resilience building for military personnel are diverse, when cognitive-behavioral therapy-based interventions, mindfulness and relaxation-based training, stress inoculation training, biofeedback-based programmes, neuropsychological-based programmes, Psychoeducational-based programmes and the family-centered resilience interventions are included. In addition, there is no single effective method; even the effects of the same resilience intervention method can vary in different situations and populations. Therefore, when constructing and implementing intervention program to build military personnel resilience, the population differences and context should be fully considered. For example, cognitive behavioral intervention methods have a certain “cultural specificity” and lack of generalizability.⁶³ Stress inoculation training should further clarify the critical values for different populations to produce learned helplessness and training effects; Besides, ABMT should select appropriate training times that can activate the reward system and reduce stress based on different populations.

Resilience is a complex, multidimensional structure. Therefore, the effectiveness of resilience interventions cannot be measured solely by the diagnosis of PTSD and other psychological disorders, and there is no “gold standard”.²⁶ Mcclung et al⁶⁴ provided an overview of physiological factors impacting or predicting resilience in military populations, which included physical performance, body composition, nutrition and diet quality, and biomarkers. It is no predictive value considering any single influencing factor alone for individual resilience.⁶⁵ Therefore, linking psychological and physiological indicators of resilience in future studies plays a vital role in forming more comprehensive assessments of resilience in military populations.

In future studies, algorithms could be developed and validated to predict PTSD by integrating various data sources, including self-report, vital signs, demographics, psychological traits, stress response, genomics, proteomics, transcriptomics, metabolomics, and epigenetics. Moreover, a multidimensional and multimodal approach across multiple systems, paths, or mechanisms could be employed to build personalized, systematic, and diversified resilience intervention programs enhancing the overall combat effectiveness of the military. The Maximal Adaptability Model predicts that when stressors progress to the extremes of hyper-stress, limits of maximal psychological and physiological adaptability may be exceeded, causing dynamic instability. Thus, we should choose and adjust the appropriate training level and training content according to psychological status and cognitive behavior monitoring of different individuals when carrying out the SIT.⁶⁶ For instance, a smartphone-delivered version of Skills for Life Adjustment and Resilience (SOLAR) could be implemented to manage stress and prevent PTSD.⁶⁷ Similarly, the Biofeedback Assisted Resilience Training (BART) digital health platform could support remote intervention and physiological and behavioral information collection.⁶⁸ Finally, Mcclung et al⁶⁴ believed that various perspectives of physical performance, nutrition, body composition, and biomarkers contribute to physiological resilience in military personnel, which can provide potential targets for future resilience intervention research.

Acknowledgments

The authors thank Dr. Mi Song for her invaluable advice.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Riley E, Mitko A, Stumps A, et al. Clinically significant cognitive dysfunction in OEF/OIF/OND veterans: prevalence and clinical associations. *Neuropsychology*. 2019;33(4):534–546. doi:10.1037/neu0000529
2. Seal KH, Bertenthal D, Samuelson K, Maguen S, Kumar S, Vasterling JJ. Association between mild traumatic brain injury and mental health problems and self-reported cognitive dysfunction in Iraq and Afghanistan Veterans. *J Rehabil Res Dev*. 2016;53(2):185–198. doi:10.1682/JRRD.2014.12.0301
3. Smith TC, Ryan MA, Wingard DL, Slymen DJ, Sallis JF, Kritz-Silverstein D. New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *BMJ*. 2008;336(7640):366–371. doi:10.1136/bmj.39430.638241.AE
4. LeardMann CA, Smith TC, Smith B, Wells TS, Ryan MA. Baseline self reported functional health and vulnerability to post-traumatic stress disorder after combat deployment: prospective US military cohort study. *BMJ*. 2009;338(2):b1273. doi:10.1136/bmj.b1273

5. Vyas KJ, Fesperman SF, Nebeker BJ, et al. Preventing PTSD and depression and reducing health care costs in the military: a call for building resilience among service members. *Mil Med.* 2016;181(10):1240–1247. doi:10.7205/MILMED-D-15-00585
6. Brewin CR, Andrews B, Valentine JD. Meta-analysis of risk factors for posttraumatic stress disorder in trauma-exposed adults. *J Consult Clin Psychol.* 2000;68(5):748–766. doi:10.1037/0022-006X.68.5.748
7. Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychol Bull.* 2003;129(1):52–73. doi:10.1037/0033-2909.129.1.52
8. Brown L, Cohen B, Costello R, Brazhnik O, Galis Z. Conceptualizing a resilience research framework at The National Institutes of Health. *Stress Health.* 2023;39(S1):4–9. doi:10.1002/smi.3260
9. Horn SR, Charney DS, Feder A. Understanding resilience: new approaches for preventing and treating PTSD. *Exp Neurol.* 2016;284(Pt B):119–132. doi:10.1016/j.expneurol.2016.07.002
10. Sefidan S, Pramstaller M, La marca R, et al. Resilience as a protective factor in basic military training, a longitudinal study of the Swiss armed forces. *Int J Environ Res Public Health.* 2021;19(1):18. doi:10.3390/ijerph19010018
11. Daffey-Moore E. Why is building psychological resilience relevant to UK Service personnel in order to improve military mental health? *BMJ Mil Health.* 2020;166(2):89–94. doi:10.1136/jramc-2018-000962
12. Nindl BC, Billing DC, Drain JR, et al. Perspectives on resilience for military readiness and preparedness: report of an international military physiology roundtable. *J Sci Med Sport.* 2018;21(11):1116–1124. doi:10.1016/j.jsams.2018.05.005
13. Sudom KA, Lee JE, Zamorski MA. A longitudinal pilot study of resilience in Canadian military personnel. *Stress Health.* 2014;30(5):377–385. doi:10.1002/smi.2614
14. Scammell J. Resilience in the workplace: personal and organisational factors. *Br J Nurs.* 2017;26(16):939. doi:10.12968/bjon.2017.26.16.939
15. Meredith LS, Sherbourne CD, Gaillott SJ, et al. Promoting psychological resilience in the US Military. *Rand Health Q.* 2011;1(2):2.
16. Bowles SV, Bates MJ. Military organizations and programs contributing to resilience building. *Mil Med.* 2010;175(6):382–385. doi:10.7205/MILMED-D-10-00099
17. Reivich KJ, Seligman ME, McBride S. Master resilience training in the US Army. *Am Psychol.* 2011;66(1):25–34. doi:10.1037/a0021897
18. Greenberg N, Langston V, Jones N. Trauma risk management (TRiM) in the UK Armed Forces. *J R Army Med Corps.* 2008;154(2):124–127. doi:10.1136/jramc-154-02-11
19. Polusny MA, Erbes CR. A dynamic, multilevel approach to conceptualising and designing resilience research in the context of military stress. *Stress Health.* 2023;39(S1):40–47. doi:10.1002/smi.3277
20. van Rooij SJH, Stevens JS, Ely TD, et al. Childhood trauma and COMT genotype interact to increase hippocampal activation in resilient individuals. *Front Psychiatry.* 2016;7:156. doi:10.3389/fpsy.2016.00156
21. Fikretoglu D, Liu A, Nazarov A, Blackler K. A group randomized control trial to test the efficacy of the Road to Mental Readiness (R2MR) program among Canadian military recruits. *BMC Psychiatry.* 2019;19(1):326. doi:10.1186/s12888-019-2287-0
22. Skeffington PM, Rees CS, Kane R. The primary prevention of PTSD: a systematic review. *J Trauma Dissociation.* 2013;14(4):404–422. doi:10.1080/15299732.2012.753653
23. Doody CB, Egan J, Bogue J, Sarma KM. Military personnels' experience of deployment: an exploration of psychological trauma, protective influences, and resilience. *Psychol Trauma.* 2022;14(4):545–557. doi:10.1037/tra0001114
24. Hourani LL, Council CL, Hubal RC, Strange LB. Approaches to the primary prevention of posttraumatic stress disorder in the military: a review of the stress control literature. *Mil Med.* 2011;176(7):721–730. doi:10.7205/MILMED-D-09-00227
25. Leppin AL, Bora PR, Tilburt JC, et al. The efficacy of resiliency training programs: a systematic review and meta-analysis of randomized trials. *PLoS One.* 2014;9(10):e111420. doi:10.1371/journal.pone.0111420
26. Doody CB, Robertson L, Cox KM, Bogue J, Egan J, Sarma KM. Pre-deployment programmes for building resilience in military and frontline emergency service personnel. *Cochrane Database Syst Rev.* 2021;12(12):Cd013242. doi:10.1002/14651858.CD013242.pub2
27. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol.* 2005;8(1):19–32. doi:10.1080/1364557032000119616
28. Peters MDJ, Marnie C, Tricco AC, et al. Updated methodological guidance for the conduct of scoping reviews. *JBI Evid Implement.* 2021;19(1):3–10. doi:10.1097/XEB.0000000000000277
29. Adler AB, Williams J, McGurk D, Moss A, Bliese PD. Resilience training with soldiers during basic combat training: randomisation by platoon. *Appl Psychol Health Well Being.* 2015;7(1):85–107. doi:10.1111/aphw.12040
30. Bouchard S, Bernier F, Boivin E, Morin B, Robillard G, Lucia A. Using biofeedback while immersed in a stressful videogame increases the effectiveness of stress management skills in soldiers. *PLoS One.* 2012;7(4):e36169. doi:10.1371/journal.pone.0036169
31. Cacioppo JT, Adler AB, Lester PB, et al. Building social resilience in soldiers: a double dissociative randomized controlled study. *J Pers Soc Psychol.* 2015;109(1):90–105. doi:10.1037/pspi0000022
32. Chongruksa D, Prinyapol P, Sawatsri S, Pansomboon C. Integrated group counselling to enhance mental health and resilience of Thai army rangers. *Asia Pac J Couns Psychother.* 2015;6(1–2):41–57. doi:10.1080/21507686.2015.1091018
33. Cohn A, Pakenham K. Efficacy of a cognitive-behavioral program to improve psychological adjustment among soldiers in recruit training. *Mil Med.* 2008;173(12):1151–1157. doi:10.7205/MILMED.173.12.1151
34. Crane MF, Boga D, Karin E, et al. Strengthening resilience in military officer cadets: a group-randomized controlled trial of coping and emotion regulatory self-reflection training. *J Consult Clin Psychol.* 2019;87(2):125–140. doi:10.1037/ccp0000356
35. Fornette M-P, Bardel M-H, Lefrançois C, Fradin J, Massioui FE, Amalberti R. Cognitive-Adaptation Training for Improving Performance and Stress Management of Air Force Pilots. *Int J Aviation Psychol.* 2012;22(3):203–223. doi:10.1080/10508414.2012.689208
36. Hourani LL, Tueller SJ, Kizakevich P, Strange L, Nelson J. Effects of Stress Inoculation Training (SIT) with relaxation breathing on posttraumatic stress disorder and other mental health outcomes in the military: a longitudinal study. *J Fam Reprod Health.* 2016. doi:10.13140/RG.2.2.34448.20489
37. Hourani L, Tueller S, Kizakevich P, et al. Effect of stress inoculation training with relaxation breathing on perceived stress and posttraumatic stress disorder in the military: a longitudinal study. *Int J Stress Manage.* 2017;25(S1):124. doi:10.1037/str0000082
38. Jha AP, Stanley EA, Kiyonaga A, Wong L, Gelfand L. Examining the protective effects of mindfulness training on working memory capacity and affective experience. *Emotion.* 2010;10(1):54–64. doi:10.1037/a0018438

39. Johnson DC, Thom NJ, Stanley EA, et al. Modifying resilience mechanisms in at-risk individuals: a controlled study of mindfulness training in Marines preparing for deployment. *Am J Psychiatry*. 2014;171(8):844–853. doi:10.1176/appi.ajp.2014.13040502
40. Jones N, Whelan C, Harden L, Macfarlane A, Burdett H, Greenberg N. Resilience-based intervention for UK military recruits: a randomised controlled trial. *Occup Environ Med*. 2019;76(2):90–96. doi:10.1136/oemed-2018-105503
41. Keynan JN, Cohen A, Jackont G, et al. Electrical fingerprint of the amygdala guides neurofeedback training for stress resilience. *Nat Hum Behav*. 2019;3(1):63–73. doi:10.1038/s41562-018-0484-3
42. Lester P, Saltzman WR, Woodward K, et al. Evaluation of a family-centered prevention intervention for military children and families facing wartime deployments. *Am J Public Health*. 2012;102(Suppl 1):S48–S54.
43. Lester P, Liang LJ, Milburn N, et al. Evaluation of a family-centered preventive intervention for military families: parent and child longitudinal outcomes. *J Am Acad Child Adolesc Psychiatry*. 2016;55(1):14–24. doi:10.1016/j.jaac.2015.10.009
44. Lewis GF, Hourani L, Tueller S, et al. Relaxation training assisted by heart rate variability biofeedback: implication for a military predeployment stress inoculation protocol. *Psychophysiology*. 2015;52(9):1167–1174. doi:10.1111/psyp.12455
45. Meland A, Fonne V, Wagstaff A, Pensgaard AM. Mindfulness-based mental training in a high-performance combat aviation population: a one-year intervention study and two-year follow-up. *Int J Aviation Psychol*. 2015;25(1):48–61. doi:10.1080/10508414.2015.995572
46. Pyne JM, Constans JI, Nanney JT, et al. Heart rate variability and cognitive bias feedback interventions to prevent post-deployment PTSD: results from a randomized controlled trial. *Mil Med*. 2019;184(1–2):e124–e132. doi:10.1093/milmed/usy171
47. Zueger R, Niederhauser M, Utzinger C, Annen H, Ehler U. Effects of resilience training on mental, emotional, and physical stress outcomes in military officer cadets. *Mil Psychol*. 2023;35(6):566–576. doi:10.1080/08995605.2022.2139948
48. Stanley EA, Schaldach JM, Kiyonaga A, Jha AP. Mindfulness-based mind fitness training: a case study of a high-stress predeployment military cohort. *Cognit Behav Pract*. 2011;18(4):566–576. doi:10.1016/j.cbpra.2010.08.002
49. Taylor MK, Stanfill KE, Padilla GA, et al. Effect of psychological skills training during military survival school: a randomized, controlled field study. *Mil Med*. 2011;176(12):1362–1368. doi:10.7205/MILMED-D-11-00149
50. Wald I, Fruchter E, Ginat K, et al. Selective prevention of combat-related post-traumatic stress disorder using attention bias modification training: a randomized controlled trial. *Psychol Med*. 2016;46(12):2627–2636. doi:10.1017/S0033291716000945
51. Wesemann U, Kowalski JT, Jacobsen T, et al. Evaluation of a technology-based adaptive learning and prevention program for stress response—a randomized controlled trial. *Mil Med*. 2016;181(8):863–871. doi:10.7205/MILMED-D-15-00100
52. Williams A, Hagerty BM, Andrei AC, Yousha SM, Hirth RA, Hoyle KS. STARS: strategies to assist navy recruits’ success. *Mil Med*. 2007;172(9):942–949. doi:10.7205/MILMED.172.9.942
53. Schrader C, Ross A. A Review of PTSD and Current Treatment Strategies. *Mo Med*. 2021;118(6):546–551.
54. Smith BW, Ortiz JA, Steffen LE, et al. Mindfulness is associated with fewer PTSD symptoms, depressive symptoms, physical symptoms, and alcohol problems in urban firefighters. *J Consult Clin Psychol*. 2011;79(5):613–617. doi:10.1037/a0025189
55. Benzo RP, Anderson PM, Bronars C, Clark M. Mindfulness for healthcare providers: the role of non-reactivity in reducing stress. *Explore*. 2018;14(6):453–456. doi:10.1016/j.explore.2018.03.008
56. Southwick SM, Charney DS. The science of resilience: implications for the prevention and treatment of depression. *Science*. 2012;338(6103):79–82. doi:10.1126/science.1222942
57. Admi H. Stress intervention. A model of stress inoculation training. *J Psychosoc Nurs Ment Health Serv*. 1997;35(8):37–41. doi:10.3928/0279-3695-19970801-16
58. Thabrew H, Ruppeldt P, Sollers JJ. Systematic review of biofeedback interventions for addressing anxiety and depression in children and adolescents with long-term physical conditions. *Appl Psychophysiol Biofeedback*. 2018;43(3):179–192. doi:10.1007/s10484-018-9399-z
59. Minassian A, Geyer MA, Baker DG, Nievergelt CM, O’Connor DT, Risbrough VB. Heart rate variability characteristics in a large group of active-duty marines and relationship to posttraumatic stress. *Psychosom Med*. 2014;76(4):292–301. doi:10.1097/PSY.0000000000000056
60. Xia HS, Li YX, Zhang QY, et al. Attention bias modification for depression: a systematic review and meta-analysis. *Front Psychiatry*. 2023;14:1098610. doi:10.3389/fpsy.2023.1098610
61. Saltzman WR, Lester P, Beardslee WR, Layne CM, Woodward K, Nash WP. Mechanisms of risk and resilience in military families: theoretical and empirical basis of a family-focused resilience enhancement program. *Clin Child Fam Psychol Rev*. 2011;14(3):213–230. doi:10.1007/s10567-011-0096-1
62. Lester P, Mogil C, Saltzman W, et al. Families overcoming under stress: implementing family-centered prevention for military families facing wartime deployments and combat operational stress. *Mil Med*. 2011;176(1):19–25. doi:10.7205/MILMED-D-10-00122
63. Thompson B. *Foundations of Behavioral Statistics: An Insight-Based Approach*. Guilford Press; 2006.
64. McClung JP, Beckner ME, Farina EK. Assessing the physiological basis for resilience in military personnel. *Stress Health*. 2023;39(S1):33–39. doi:10.1002/smi.3271
65. Osório C, Probert T, Jones E, Young AH, Robbins I. Adapting to stress: understanding the neurobiology of resilience. *Behav Med*. 2017;43(4):307–322. doi:10.1080/08964289.2016.1170661
66. Flood A, Keegan RJ. Cognitive resilience to psychological stress in military personnel. *Front Psychol*. 2022;13:809003. doi:10.3389/fpsyg.2022.809003
67. Metcalf O, Gibson K, Fredrickson J, Finlayson-Short L, Varker T, O’Donnell M. Design, development and randomised controlled trial protocol of a smartphone-delivered version of ‘SOLAR’ for emergency service workers to manage stress and trauma. *BMJ Open*. 2023;13(2):e062710. doi:10.1136/bmjopen-2022-062710
68. Davila MI, Kizakevich PN, Eckhoff R, et al. Use of mobile technology paired with heart rate monitor to remotely quantify behavioral health markers among military reservists and first responders. *Mil Med*. 2021;186(Supplement_1):17–24. doi:10.1093/milmed/usaa395

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>