



# Complete Genome Sequence of *Bacillus methylotrophicus* Strain NKG-1, Isolated from the Changbai Mountains, China

Binghua Liu,<sup>a</sup> Beibei Ge,<sup>a</sup> Noreen Azhar,<sup>b</sup> Wenjun Zhao,<sup>a</sup> Hailan Cui,<sup>a</sup> Kecheng Zhang<sup>a</sup>

<sup>a</sup>Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China

<sup>b</sup>Institute of Biotechnology and Genetic Engineering, The University of Agriculture, Peshawar, Pakistan

**ABSTRACT** We report here the complete genome sequence of *Bacillus methylotrophicus* NKG-1, isolated from rare dormant volcanic soils on the Changbai Mountains in China. The 4.20-Mb genome contains 4,432 genes and has a G+C content of 47.06%.

*Bacillus methylotrophicus* is a Gram-positive bacterium associated with plant roots and has beneficial effects on the growth of many plants (1). The indole-3-acetic acid content was found to be 2- and 3-fold higher in the shoots and roots of bacterized seedlings, respectively, than in the control group (2). This strain can be used as a biofertilizer to promote plant growth and as a biopesticide to target root pathogens, including Gram-negative bacteria and fungi (3, 4). *B. methylotrophicus* NKG-1 was isolated from rare dormant volcanic soils on the Changbai Mountains in China. This strain is an effective biocontrol agent against 13 different kinds of soilborne phytopathogenic fungi and can promote the growth of tomato plants under both greenhouse and field conditions (5). This strain has been preserved at the China General Microbiological Culture Collection Center (CGMCC12055) and can be cultured at 28°C in Luria-Bertani (LB) medium.

Genome sequencing was performed using a third-generation PacBio RSII instrument. The use of a single-molecule sequencing approach has many advantages, including a long reading sequence, which facilitates easier genome assembly. The SMRT Portal software was employed to assemble reads into preliminary results (6, 7), and reads were compared with the assembled sequence to determine the sequencing depth. Chromosome and plasmid sequences were distinguished by sequence length and whether the sequence is annular. The GeneMarkS software (<http://topaz.gatech.edu/GeneMark/genemarks.cgi>) was used to predict bacterial protein-coding genes. Gene functional annotation was performed based on BLASTp searches against nonredundant (NR), Kyoto Encyclopedia of Genes and Genomes (KEGG) (8), Protein Family (Pfam) (9), and Cluster of Orthologous Groups (COG) (10) databases.

The *Bacillus methylotrophicus* genome is 4,197,217 bp, with a G+C content of 47.06%. A single chromosome containing 4,432 genes covering 3,768,021 bp, with an average gene length of 850 bp, accounts for ~89.77% of the genome. In addition, the genome encodes 114 structural RNAs, including 9 5S rRNAs, 9 16S rRNAs, 9 23S rRNAs, and 87 tRNAs.

A previous study showed that NKG-1 fermentation broth inhibits the growth of *Botrytis cinerea* on tomato seedling leaves *in vitro* by 60%, resulting in significant increases in seedling fresh weight (27.4%), seedling length (12.5%), and root length (57.7%) compared to the controls (5). The genome of *B. methylotrophicus* will be compared with those of other *Bacillus* strains in future research to illuminate the mechanisms promoting plant growth and biocontrol activity against fungal diseases, as well as to identify the compounds involved.

**Received** 21 November 2017 **Accepted** 27 November 2017 **Published** 18 January 2018

**Citation** Liu B, Ge B, Azhar N, Zhao W, Cui H, Zhang K. 2018. Complete genome sequence of *Bacillus methylotrophicus* strain NKG-1, isolated from the Changbai Mountains, China. *Genome Announc* 6:e01454-17. <https://doi.org/10.1128/genomeA.01454-17>.

**Copyright** © 2018 Liu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kecheng Zhang, [zhangkecheng@sina.com](mailto:zhangkecheng@sina.com).

**Accession number(s).** The complete genome sequence of *Bacillus methylotrophicus* NKG-1 has been deposited in GenBank under accession no. [CP024203](https://doi.org/10.1093/jqs.0.015487-0).

## ACKNOWLEDGMENTS

This work was supported by grants from the National Natural Science Foundation of China (grant 31401796), the Special Fund for Basic-Scientific Research in Chinese Academy of Agricultural Sciences in China (grants Y2017JC12 and 1610142016017), and the National key Research & Development (R&D) plan in China (grant YFD020130).

## REFERENCES

1. Madhaiyan M, Poonguzhali S, Kwon SW, Sa TM. 2010. *Bacillus methylotrophicus* sp. nov., a methanol-utilizing, plant-growth-promoting bacterium isolated from rice rhizosphere soil. *Int J Syst Evol Microbiol* 60:2490–2495. <https://doi.org/10.1099/ijs.0.015487-0>.
2. Pérez-Flores P, Valencia-Cantero E, Altamirano-Hernández J, Pelagio-Flores R, López-Bucio J, García-Juárez P, Macías-Rodríguez L. 2017. *Bacillus methylotrophicus* M4-96 isolated from maize (*Zea mays*) rhizoplane increases growth and auxin content in *Arabidopsis thaliana* via emission of volatiles. *Protoplasma* 254:2201–2213. <https://doi.org/10.1007/s00709-017-1109-9>.
3. Gagne-Bourgue F, Aliferis KA, Seguin P, Rani M, Samson R, Jabaji S. 2013. Isolation and characterization of indigenous endophytic bacteria associated with leaves of switchgrass (*Panicum virgatum* L.) cultivars. *J Appl Microbiol* 114:836–853. <https://doi.org/10.1111/jam.12088>.
4. Bacon CW, Palencia ER, Hinton DM. 2015. Abiotic and biotic plant stress-tolerant and beneficial secondary metabolites produced by endophytic *Bacillus* species, p 163–177. In Arora NK (ed), *Plant microbes symbiosis: applied facets*. Springer, New Delhi, India.
5. Ge BB, Liu BH, Nwet TT, Zhao WJ, Shi LM, Zhang KC. 2016. *Bacillus methylotrophicus* strain NKG-1, isolated from Changbai Mountain, China, has potential applications as a biofertilizer or biocontrol agent. *PLoS One* 11:e0166079. <https://doi.org/10.1371/journal.pone.0166079>.
6. Berlin K, Koren S, Chin CS, Drake JP, Landolin JM, Phillippy AM. 2015. Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. *Nat Biotechnol* 33:623–630. <https://doi.org/10.1038/nbt.3238>.
7. Koren S, Phillippy AM. 2015. One chromosome, one contig: complete microbial genomes from long-read sequencing and assembly. *Curr Opin Microbiol* 23:110–120. <https://doi.org/10.1016/j.mib.2014.11.014>.
8. Ogata H, Goto S, Fujibuchi W, Kanehisa M. 1998. Computation with the KEGG pathway database. *Biosystems* 47:119–128. [https://doi.org/10.1016/S0303-2647\(98\)00017-3](https://doi.org/10.1016/S0303-2647(98)00017-3).
9. Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, Boursnell C, Pang N, Forslund K, Ceric G, Clements J, Heger A, Holm L, Sonnhammer EL, Eddy SR, Bateman A, Finn RD. 2012. The Pfam protein families database. *Nucleic Acids Res* 40:D290–DD301. <https://doi.org/10.1093/nar/gkr1065>.
10. Tatusov RL, Galperin MY, Natale DA, Koonin EV. 2000. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 28:33–36. <https://doi.org/10.1093/nar/28.1.33>.