

## Preferred Methods of Suicide and Most Common Poisonings in India

Dear Editor,

Tanuj Kanchan

A recently published paper on herbicide poisoning in a tertiary care hospital in south India<sup>[1]</sup> in 'Toxicology International' is certainly a useful addition to the literature. The observations presented by the authors are intended for the clinicians, public health practitioners, epidemiologists as well as medicolegal experts. The views expressed by the authors<sup>[1]</sup> in their opening statement of the abstract ('Herbicide poisoning is most common method of suicide in India') and the Introduction section ('Among pesticides, herbicides like paraquat and glyphosate are most common poisonings responsible for high morbidity and mortality'); however, needs to be refuted for the true understanding of epidemiology of poisoning and suicides in Indian scenario.

Suicide is a serious global health problem, and easy access to lethal means is one of the well-recognized risk factor for suicide.<sup>[2]</sup> Hanging and poisoning are reported as the most common method of completed suicides in the region and most parts in India.<sup>[3]</sup> Poisoning irrespective of its manner is associated with considerable mortality and morbidity worldwide. The causes and means of poisonings vary from country to country, in different setups, regions, and between sexes and age groups.<sup>[4-6]</sup> Agrochemical pesticides have been reported as the most common cause of acute poisonings in the region, while most of the fatalities are associated with organophosphate compounds.<sup>[7]</sup> Studies suggest that phosphides and organophosphates are commonly implicated in fatal poisonings in northern and southern part of India, respectively.<sup>[4,8]</sup>

The present communication, thus, briefs on the preferred methods of suicide and most common poisonings in the region and the country. Cherukuri *et al.*,<sup>[1]</sup> need to be applauded for elaborating on a 'relatively uncommon poisoning' in India with discussion on its clinical characteristics and treatment pattern.

Department of Forensic Medicine, Kasturba Medical College, Mangalore, A Constituent College of Manipal University, Karnataka, India


### Address for correspondence:

Dr. Tanuj Kanchan,  
Department of Forensic Medicine and Toxicology, Kasturba Medical College, Manipal University, Mangalore, Karnataka - 575 001, India.  
E-mail: tanuj.kanchan@manipal.edu

## REFERENCES

1. Cherukuri H, Pramoda K, Rohini D, Thunga G, Vijaynarayana K, Sreedharan N, *et al.* Demographics, clinical characteristics and management of herbicide poisoning in tertiary care Hospital. *Toxicol Int* 2014;21:209-13.
2. Kanchan T. Suicide predictors and statistics. In: Payne-James J, Byard R, editors. *Encyclopedia of Forensic and Legal Medicine*. 2<sup>nd</sup> ed. (In Press). Waltham: Academic Press, Elsevier; 2015.
3. Kanchan T, Menon A, Menezes RG. Methods of choice in completed suicides: Gender differences and review of literature. *J Forensic Sci* 2009;54:938-42.
4. Kanchan T, Menezes RG, Kumar TS, Bakkannavar SM, Bukelo MJ, Sharma PS, *et al.* Toxicoepidemiology of fatal poisonings in Southern India. *J Forensic Leg Med* 2010;17:344-7.
5. Kanchan T, Menezes RG. Suicidal poisoning in Southern India: Gender differences—authors' response. *J Forensic Leg Med* 2009;16:365.
6. Ram P, Kanchan T, Unnikrishnan B. Pattern of acute poisonings in children below 15 years--a study from Mangalore, South India. *J Forensic Leg Med* 2014;25:26-9.
7. Bumrah GS, Krishan K, Kanchan T, Sharma M, Sodhi GS. Phosphide poisoning: A review of literature. *Forensic Sci Int* 2012;214:1-6.
8. Singh B, Unnikrishnan B. A profile of acute poisoning at Mangalore (South India). *J Clin Forensic Med* 2006;13:112-6.

### Access this article online

<b>Quick Response Code:</b> 	<b>Website:</b> <a href="http://www.toxicologyinternational.com">www.toxicologyinternational.com</a>
	<b>DOI:</b> 10.4103/0971-6580.155400