



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Dramatic rise in seroprevalence rates of SARS-CoV-2 antibodies among healthy blood donors: The evolution of a pandemic



Maher A. Sughayer^{a,*}, Asem Mansour^b, Abeer Al Nuirat^a, Lina Souan^a,
Mohammad Ghanem^a, Mahmoud Siag^a

^a Department of Pathology and Laboratory Medicine, King Hussein Cancer Center, Amman, Jordan

^b Department of Radiology, and CEO, King Hussein Cancer Center, Jordan

ARTICLE INFO

Article history:

Received 26 February 2021

Received in revised form 15 April 2021

Accepted 16 April 2021

Keywords:

Covid-19

Seroprevalence

Blood donors

SARS-CoV-2 antibodies

ABSTRACT

Background: Seroprevalence studies of SARS-CoV-2 antibodies are useful in assessing the epidemiological status in the community, and the degree of spread.

Objective: To study the seroprevalence rates of SARS-CoV-2 antibodies among healthy blood donors in Jordan, at various points of time and as the pandemic evolves in the community.

Methods: In total, 1374 blood donor samples, from three groups, were tested for SARS-CoV-2 total immunoglobulin antibodies. In the first group, samples from 734 individuals (from donations made between January and June 2020) were tested in June. In the second group, 348 individuals were tested in September 2020. The third group of 292 individuals was tested in February 2021. A qualitative assay was used for testing (specificity 99.8%, sensitivity 100%).

Results: The first two groups, from January–June and September 2020, when confirmed Covid-19 cases numbered between several hundred and 3000, showed a seroprevalence rate of 0% (95% CI 0.00–0.51%). The third group (early February 2021), when the number of confirmed cases had reached 100 times that of September 2020, revealed a seroprevalence of 27.4% (95% CI 22.5–32.9%).

Conclusions: A dramatic rise in seroprevalence of SARS-CoV-2 antibodies was seen among healthy blood donors in Jordan, in parallel with widespread intracommunity transmission of the disease. This information is useful for assessing the degree of herd immunity, and provides for better understanding of the pandemic.

© 2021 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The coronavirus disease 2019 (COVID-19) pandemic had resulted in more than 147 million cases of confirmed infection, and more than 3 million deaths worldwide as of week 2021-16, 2021 (European Centre for Disease Prevention and Control, 2021).

Population-based seroprevalence studies are extremely important for understanding the evolution of the pandemic, and to estimate infection rates and prevalence. They are also important for calculating absolute risks of infection as well as death rates, and to predict the spread of the virus in communities based on the level of herd immunity following infections and/or vaccination. In addition, they are used for planning and monitoring the impact of implementation and relaxation of epidemic mitigation policies (Busch and Stone, 2021)

The true prevalence of infection is believed to be several times more than the number of PCR-confirmed cases because of the large number of asymptomatic infections and/or mild infections that have gone untested, especially early in the pandemic (Huang et al., 2020; McLaughlin et al., 2020; Busch and Stone, 2021). The ratio of estimated to reported infections can be as much as 12.5 (Bajema et al., 2020).

Numerous population seroprevalence studies have been conducted (Bajema et al., 2020; Lai et al., 2020; Chughtai et al., 2020; McLaughlin et al., 2020; Naranbhai et al., 2020; Sam et al., 2021; Sutton et al., 2020; Menachemi et al., 2020; Silveira et al., 2020; Kar et al., 2021; Vena et al., 2020; Pollán et al., 2020; Bogogiannidou et al., 2020; Poustchi et al., 2020; Ng et al., 2020; Figueiredo-Campos et al., 2020; Stringhini et al., 2020; Havers et al., 2020; Ho et al., 2020; Xu et al., 2020; Qutob et al., 2021; Capai et al., 2020; Sood et al., 2020; Godbout et al., 2020; Rostami et al., 2020) in efforts to estimate the true prevalence of COVID-19 infection.

The largest seroprevalence study was that by Bajema et al., in the USA, which showed that by September 2020 the estimated

* Corresponding author at: 202 Q. Rania St., Amman, 11941, Jordan.
E-mail address: Msughayer@KHCC.jo (M.A. Sughayer).

population seroprevalence was less than 10% in the majority of tested communities, although it ranged from 0% up to 23.3% in the hardest-hit areas. This wide variation reflected the levels of transmission in the tested communities and the numbers of PCR-confirmed reported cases (Bajema et al., 2020).

These population-based seroprevalence studies varied in the targeted populations tested and in the recruitment strategies used, which may explain some of the variation. Among the populations tested were healthy blood donors (Amorim Filho et al., 2020; Sughayer et al., 2020; Slot et al., 2020; Nesbitt et al., 2021; Gallian et al., 2020; Olariu et al., 2021; Younas et al., 2020; Banjar et al., 2021; Uyoga et al., 2021; Busch and Stone, 2021; Martinez-Acuña et al., 2020; Saeed et al., 2021; Fiore et al., 2021). Again, the seroprevalence among healthy blood donors and other population groups studied varied according to the community tested and the time of testing, in terms of the pandemic evolution (Lai et al., 2020; Rostami et al., 2020).

The aim of this longitudinal study was to estimate the seroprevalence of Covid-19 among healthy blood donors in Jordan at different points in time, in order to assess the degree of community spread and herd immunity, and to further understand the evolution of the pandemic. To our knowledge, this is the first longitudinal seroprevalence study with blood donors, and the first from Jordan.

Methods

Study population

The study population consisted of three groups of healthy blood donors. The distribution of donors in terms of donation period is detailed in Table 1. Individuals in the first group were randomly chosen from donors attending the blood bank between January and June 2020. The aim was to include donors from January and February 2020, even though the first case of COVID-19 in Jordan was recorded in March of that year. The number chosen from each month was arbitrary. The smallest number in the first group was in June because testing was carried out in the first week of that month. The second and third group were randomly chosen in the first half of September 2020 and late January–early February 2021, respectively. The timing of the second group was chosen to assess the situation when society began to open up again in early September. The third group was chosen to assess the impact of the COVID-19 wave that peaked in January of 2021.

Table 1
Demographics and period of donation of all donors.

Demographics of the 1374 healthy donors, No. (%)	
Male	1177 (85.7%)
Female	197 (14.3%)
Residential area: Amman	1077 (78.4%)
West Amman	706 (65.6%)
East Amman	371 (34.4%)
District outside Amman	297 (21.6%)
North	128 (9.3%)
Center	138 (10.1%)
South	31 (2.3%)
Period	
January 2020	104 (7.6%)
February 2020	89 (6.5%)
March 2020	90 (6.6%)
April 2020	147 (10.7%)
May 2020	229 (16.7%)
June 2020	75 (5.4%)
September 2020	348 (25.3%)
Jan 28–Feb 5, 2021	292 (21.3%)

Samples

Leftover sera and/or plasma collected routinely during the process of blood or apheresis platelet donations were used for the study. The donors were healthy, asymptomatic subjects between the ages of 18 and 63, who underwent routine screening to determine their acceptability for donation, as per standard practice. The sera were tested in three batches at three different points in time. The first batch, consisting of 734 samples representing the period from January to June 2020, was tested in June of that year. The second batch of 348 samples was collected and tested in September 2020, while the third batch of 292 samples was collected during the period January 28–February 5, 2021 and tested in the same period. Thus the total number of donors tested in the study was 1374.

Testing methodology

The samples were tested on Cobas 6000 or Cobas Pro Roche analyzers, using the Elecsys-Anti-SARS-CoV-2 kit (Roche Diagnostics GmbH, Mannheim, Germany), according to the manufacturer's recommendations. This commercially available FDA-approved kit for total immunoglobulins against SARS-CoV-2 is a qualitative assay that uses an electrochemiluminescence immunoassay (ECLIA) for the in vitro detection of total antibodies (IgA, IgM, and IgG) to SARS-CoV-2 in human serum and plasma. The assay uses a recombinant protein representing the nucleocapsid (N) antigen for the determination of antibodies against SARS-CoV-2. The analyzer automatically calculates the cutoff, based on the measurements of two calibrators. The result for a sample is given either as reactive or non-reactive, as well as in the form of a cutoff index (COI; signal sample/cutoff). If the COI is <1.0 then the assay is non-reactive (negative for anti-SARS-CoV-2 antibodies); if COI is ≥ 1.0 then the assay is considered reactive (positive for anti-SARS-CoV-2 antibodies).

The test was validated by the manufacturer using 5272 samples, including blood donors, routine diagnostics, other corona viruses, and common cold panels. The specificity was determined to be 99.8% (95% CI, 99.7–99.9%), while the sensitivity was 100% (95% CI, 88.3–100.0%).

In-house validation using serum samples obtained from previously RT-PCR-confirmed COVID-19-infected patients who had recovered at least 1 month prior to sampling was performed on two occasions: in June 2020 using four samples and in September 2020 using six samples.

Statistical methods

Descriptive statistics were used to analyze the results. The chi-square test was used to compare the seropositive donors' characteristics with the seronegative ones. Student's *t*-test was used to compare the mean ages of the seropositive and seronegative donors. The estimated crude prevalence rate was adjusted to the test's sensitivity and specificity using an online tool (<http://www2.univet.hu/users/jreiczig/CI4prevSeSp/calc02/index.php>) in line with methods described by Lang and Reiczigel (Lang and Reiczigel, 2014).

Results

The in-house validation of the 10 samples from known confirmed previously infected patients revealed positive results for the presence of total anti-SARS-CoV-2 nucleocapsid antibodies.

The demographics of the donors and the periods of donation are summarized in Table 1. The donors were mostly males (86%) and from the capital city of Amman, Jordan (78%).

Table 2
Comparison of seronegative and seropositive donors in the third group

Category	Seropositive donors 80 of 292	Seronegative donors 212 of 292	p-value	Crude prevalence rate for seropositive donors
Number of donors	80 (%)	212 (%)		27.4%
Male	70 (87.5%)	184 (86.8%)	0.9	27.6%
Female	10 (12.5%)	28 (11.8%)		26.3%
Age (yrs) 18–30	47 (58.8%)	116 (54.7%)	0.4	28.8%
31–40	21 (26.3%)	52 (24.5%)		28.8%
41–50	11 (13.7%)	24 (11.3%)		31.4%
51–65	1 (1.2%)	11 (5.2%)		8.3%
Unknown	0	9 (4.2%)		
Blood group: O	33 (41.3%)	88 (41.5%)	0.5	27.3%
A	23 (28.8%)	73 (34.4%)		24.0%
B	15 (18.7%)	37 (17.5%)		28.8%
AB	9 (11.2%)	14 (6.6%)		39.1%
Rhesus blood type: +	70 (87.5%)	194 (91.5%)	0.3	70 (26.5%)
–	10 (12.5%)	18 (8.5%)		10 (35.7%)
Residential location North	7 (8.8%)	33 (15.6%)	0.3	17.5%
Center, including Amman	68 (85.0%)	168 (79.2%)		28.8%
South	5 (6.2%)	11 (5.2%)		31.3%
History of previous COVID-19:				
PCR-confirmed past infection	16 (20%)	0	NA	NA
PCR negative/not performed	47 (58.8%)	180 (84.9%)		
No information available	17 (21.2%)	32 (15.1%)		

The first group was previously reported in a preprint (Sughayer et al., 2020) and, along with the second group, revealed a seroprevalence rate of 0% (95% CI, 0.00–0.51%).

The third group, which represented the most recent period of January 28–February 5, 2021, showed a positive serological test for SARS-CoV-2 in 80 of 292 donors, giving a crude seroprevalence rate of 27.4%. The adjusted estimated seroprevalence rate was 27.3% (95% CI, 22.5–32.9%). The ages of individuals in this group ranged between 18 and 65 years. The mean ages (\pm SD) of the seronegative donors and seropositive donors were 31.4 ± 9.75 and 29.7 ± 8.56 years, respectively, with no statistically significant difference between these mean ages ($p = 0.13$). The demographics and characteristics of the seropositive donors in comparison with the seronegative ones are shown in Table 2. Most of those who tested positive (85%) were in the 18–40 years age group. However, there were no statistically significant differences between the seropositive and seronegative donors in terms of gender, age, blood group, or residence. Males and females were almost equally affected (27.6% vs 26.3%).

One fifth of the seropositive donors were retrospectively found to have been confirmed positive for COVID-19 infection by PCR testing. Forty seven (58.8%) were not known to have had the disease, and so either did not undergo PCR testing or, if they did, produced a negative result. There was no information with regard to previous infections for 17 of the seropositive donors.

Discussion

The importance of serological testing for SARS-CoV-2 antibodies has been previously highlighted (Busch and Stone, 2021; Raoult, 2021). Among the advantages of such testing is an understanding of the evolution of the pandemic in terms of generating a rough estimate of the prevalence of infection. This will help health planners and decision makers to properly enforce or relax mitigation measures. And, most importantly in these times when vaccines are being rolled out, it serves in estimating the risk rates for infection and the degree of herd immunity, and helps in prioritizing vaccine recipients.

In this study the seroprevalence rates in healthy blood donors were measured at three points in time. The results are striking in that they show a dramatic change from 0% early and in the middle of the pandemic, up to 27.4% in February 2021. These findings

would seem reasonable if we consider the cumulative number of confirmed cases around these times in Jordan. Figure 1 shows the cumulative daily cases of COVID-19 in Jordan. In the first period up to June 2020 there were only several hundred confirmed cases, which increased gradually to around 3000 cases in early September (WHO, 2020). However, towards the end of September and beyond the cumulative number of cases started a steep rise such that, by February 5, 2021, the number had increased 100-fold since September 2020. It is worth mentioning that a strict lockdown was in effect until early June, which was gradually relaxed over the following 3 months, with full opening of all sectors, including schools and international travel, in September 2020. It is clear that the first wave of the COVID-19 pandemic in Jordan actually started in late September when the community was fully open and the intracommunity spread became evident. Before that, the several hundred cases were limited to transmission within known specific hot foci. This explains the extremely low seroprevalence initially found in June and September of 2020, as the infection transmission was under strict control with quarantine routinely imposed on all contacts of index cases.

The crude seroprevalence rate found in early February 2021 of 27.4%, if generalized to the entire Jordan population, would mean that the number of cases was roughly 2.7 million in a population of 10 million. If true, this would mean that there were eight cases for every confirmed case. This high ratio of estimated to reported cases




Figure 1. Cumulative number of confirmed COVID-19 cases in Jordan.

is similar to some of the highest ratios reported by Bajema et al. in the USA (Bajema et al., 2020). Looking at it from a different perspective, we can also see that in our rather small sample, one fifth of the donors had been previously confirmed positive by the PCR test, so we may assume a ratio of estimated to reported cases of five, as a lowest estimate.

Most of our seropositive donors (85%) were young – below 40 years of age. Other studies have shown inconsistent findings, with either similar age group distributions to our cohort or the opposite situation, with higher rates in older individuals (Bajema et al., 2020; Martinez-Acuña et al., 2020; Vena et al., 2020; Olariu et al., 2021). Nevertheless, this finding regarding age helps in setting vaccination priorities for those older than 40 years.

Our findings were also in line with previous studies, which showed that early in the pandemic the seroprevalence of COVID-19 in blood donors or other nontargeted populations was very low, ranging from 0% to 2.0% (Erikstrup et al., 2021; Godbout et al., 2020; Qutob et al., 2021; Xu et al., 2020; Ho et al., 2020; Banjar et al., 2021; Nesbitt et al., 2021; Slot et al., 2020; Fiore et al., 2021; Saeed et al., 2021). However, other studies, especially those from hard-hit communities early in the pandemic, showed relatively high rates of up to 23% (Rosenberg et al., 2020; Percivalle et al., 2020).

The high seroprevalence rate found in our study for early February 2021 was similar to that for communities that were severely affected by the pandemic, such as those of New York City and Chelsea, Massachusetts (Bajema et al., 2020; Naranbhai et al., 2020).

With regard to blood group, in terms of a decreased risk for COVID-19 infection in those of blood group O (Gallian et al., 2020), our limited data did not reveal such an association.

To our knowledge, our study was the first longitudinal study of COVID-19 seroprevalence in healthy blood donors. The results illustrate the evolution of the pandemic, and the dynamics and hidden magnitude of the problem.

There were some limitations in our study, including the small number of blood donors in the third group. In addition, although the age groups represented in our study constituted around 60% of the Jordanian population, with the rest being mostly children under the age of 18, the blood donors were mostly males and of younger age groups. Thus, it may be difficult to draw generalizations, as there may have been selection bias and nonrepresentation of the entire population. Further studies are recommended to include children and the elderly.

In conclusion, a dramatic rise in seroprevalence of SARS-CoV-2 from almost 0% to 27.4% in healthy blood donors was seen as the Jordanian community entered into the peak of the first wave of COVID-19. An estimate of the true prevalence was achieved through this longitudinal serological study, leading to much better insight and understanding of the evolution of the pandemic in Jordan.

Several gaps are present in this study. These include the limitations previously mentioned, especially the small number of the donors, the bias with regard to age and gender, and the consequent lack of full representation of the entire community. Nevertheless, this study represents an important addition to our means of controlling and fighting COVID-19.

Conflicts of interest

The authors declare no conflicts of interest.

Funding

No external funding was obtained.

Ethical approval

The institutional review board (IRB) at King Hussein Cancer Center approved the study.

References

- Amorim Filho L, Szwarcwald CL, de Oliveira Garcia Mateos S, de Leon ACMP, de Andrade Medronho R, Veloso VG, et al. Seroprevalence of anti-SARS-CoV-2 among blood donors in Rio de Janeiro, Brazil. *Rev Saude Publica* 2020;54:69.
- Bajema KL, Wiegand RE, Cuffe K, Patel SV, Iachan R, Lim T, et al. Estimated SARS-CoV-2 seroprevalence in the US as of September 2020. *JAMA Intern Med* 2021;181(4):450–460. doi:<http://dx.doi.org/10.1001/jamainternmed.2020.7976>.
- Banjar A, Al-Tawfiq JA, Alruwaily A, Alserehi H, Al-Qunaibet A, Alaswad R, et al. Seroprevalence of antibodies to SARS-CoV-2 among blood donors in the early months of the pandemic in Saudi Arabia. *Int J Infect Dis* 2021;104:452–7. doi:<http://dx.doi.org/10.1016/j.ijid.2021.01.028>.
- Bogiannidou Z, Vontas A, Dadouli K, Kyritsi MA, Soteriades S, Nikoulis DJ, et al. Repeated leftover serosurvey of SARS-CoV-2 IgG antibodies, Greece, March and April 2020. *Eurosurveillance* 2020;25:2001369.
- Busch MP, Stone M. Serosurveillance for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) incidence using global blood donor populations. *Clin Infect Dis* 2021;72:254–6. doi:<http://dx.doi.org/10.1093/cid/ciaa1116>.
- Capai L, Ayhan N, Masse S, Canarelli J, Priet S, Simeoni M-H, et al. Seroprevalence of SARS-CoV-2 IgG antibodies in Corsica (France), April and June 2020. *J Clin Med* 2020;9;. doi:<http://dx.doi.org/10.3390/jcm9113569>.
- Chughtai OR, Batool H, Khan MD, Chughtai AS. Frequency of COVID-19 IgG antibodies among Special Police Squad Lahore, Pakistan. *J Coll Physicians Surg Pak* 2020;30:735–9. doi:<http://dx.doi.org/10.29271/jcpsp.2020.07.735>.
- Nesbitt Daniel J, Jin Daniel P, Hogan Joseph W, Chan Philip A, Simon Melissa J, Vargas Matthew, et al. Low seroprevalence of SARS-CoV-2 in Rhode Island blood donors determined using multiple serological assay formats. *BMC Infect Dis* (preprint available at research square) 2021;. doi:<http://dx.doi.org/10.21203/rs.3.rs-76664/v1> Submitted for publication.
- Erikstrup C, Hother CE, Pedersen OBV, Mølbak K, Skov RL, Holm DK, et al. Estimation of SARS-CoV-2 infection fatality rate by real-time antibody screening of blood donors. *Clin Infect Dis* 2021;72:249–53.
- European Centre for Disease Prevention and Control, Covid-19. <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases>. [Accessed 11 February 2021].
- Figueiredo-Campos P, Blankenhaus B, Mota C, Gomes A, Serrano M, Ariotti S, et al. Seroprevalence of anti-SARS-CoV-2 antibodies in COVID-19 patients and healthy volunteers up to 6 months post disease onset. *Eur J Immunol* 2020;50:2025–40. doi:<http://dx.doi.org/10.1002/eji.202048970>.
- Fiore JR, Centra M, De Carlo A, Granato T, Rosa A, Sarno M, et al. Results from a survey in healthy blood donors in South Eastern Italy indicate that we are far away from herd immunity to SARS-CoV-2. *J Med Virol* 2021;93:1739–42.
- Gallian P, Pastorino B, Morel P, Chiaroni J, Ninove L, de Lamballerie X. Lower prevalence of antibodies neutralizing SARS-CoV-2 in group O French blood donors. *Antivir Res* 2020;181:104880.
- Godbout EJ, Pryor R, Harmon M, Montpetit A, Greer J, Bachmann LM, et al. Severe acute respiratory coronavirus virus 2 (SARS-CoV-2) seroprevalence among healthcare workers in a low prevalence region. *Infect Control Hosp Epidemiol* 2020;1–3;. doi:<http://dx.doi.org/10.1017/ice.2020.1374>.
- Havers FP, Reed C, Lim T, Montgomery JM, Klenna JD, Hall AJ, et al. Seroprevalence of antibodies to SARS-CoV-2 in 10 sites in the United States, March 23–May 12, 2020. *JAMA Intern Med* 2020;180:1576–86.
- Ho H-L, Wang F-Y, Lee H-R, Huang Y-L, Lai C-L, Jen W-C, et al. Seroprevalence of COVID-19 in Taiwan revealed by testing anti-SARS-CoV-2 serological antibodies on 14,765 hospital patients. *Lancet Reg Health West Pac* 2020;3:100041. doi:<http://dx.doi.org/10.1016/j.lanwpc.2020.100041>.
- Huang L, Zhang Xiuwen, Zhang Xinyue, Wei Z, Zhang L, Xu J, et al. Rapid asymptomatic transmission of COVID-19 during the incubation period demonstrating strong infectivity in a cluster of youngsters aged 16–23 years outside Wuhan and characteristics of young patients with COVID-19: a prospective contact-tracing study. *J Infect* 2020;80:e1–e13. doi:<http://dx.doi.org/10.1016/j.jinf.2020.03.006>.
- Kar SS, Sarkar S, Murali S, Dhodapkar R, Joseph NM, Aggarwal R. Prevalence and time trend of SARS-CoV-2 infection in Puducherry, India, August–October 2020. *Emerg Infect Dis J* 2021;27:666. doi:<http://dx.doi.org/10.3201/eid2702.204480>.
- Lai C-C, Wang J-H, Hsueh P-R. Population-based seroprevalence surveys of anti-SARS-CoV-2 antibody: an up-to-date review. *Int J Infect Dis* 2020;101:314–22. doi:<http://dx.doi.org/10.1016/j.ijid.2020.10.011>.
- Lang Z, Reizigal J. Confidence limits for prevalence of disease adjusted for estimated sensitivity and specificity. *Prev Vet Med* 2014;113:13–22. doi:<http://dx.doi.org/10.1016/j.prevetmed.2013.09.015>.
- Martinez-Acuña N, Avalos-Nolazco D, Rodriguez-Rodriguez D, Martinez-Liu C, Taméz RC, Flores-Arechiga A. Seroprevalence of anti-SARS-CoV-2 antibodies in blood donors from Nuevo Leon state, Mexico, during the beginning of the COVID-19 pandemic. *medRxiv* 2020;20240325. doi:<http://dx.doi.org/10.1101/2020.11.28.20240325> 2020.11.28.20240325 In preparation.

- McLaughlin CC, Doll MK, Morrison KT, McLaughlin WL, O'Connor T, Sholukh AM, et al. High community SARS-CoV-2 antibody seroprevalence in a ski resort community, Blaine County, Idaho, US. Preliminary results. *Medrxiv* [Preprint] 2020;32743610, doi:<http://dx.doi.org/10.1101/2020.07.19.20157198>.
- Menachemi N, Yiannoutsos CT, Dixon BE, Duszynski TJ, Fadel WF, Wools-Kaloustian KK, et al. Population point prevalence of SARS-CoV-2 infection based on a statewide random sample – Indiana, April 25–29, 2020. *Morb Mortal Wkly Rep* 2020;69:960.
- Naranbhai V, Chang CC, Beltran WFG, Miller TE, Astudillo MG, Villalba JA, et al. High seroprevalence of anti-SARS-CoV-2 antibodies in Chelsea, Massachusetts. *J Infect Dis* 2020;222:1955–9.
- Ng DL, Goldgof GM, Shy BR, Levine AG, Balcerak J, Bapat SP, et al. SARS-CoV-2 seroprevalence and neutralizing activity in donor and patient blood. *Nat Commun* 2020;11:1–7.
- Olariu TR, Lighezan R, Ursoniu S, Craciun AC, Paduraru AA, Lupu MA, n.d. Seroprevalence of SARS-CoV-2 antibodies in 2115 blood donors from Romania. *Clin Microbiol Infect*. <https://doi.org/10.1016/j.cmi.2020.12.027>.
- Percivalle E, Cambiè G, Cassaniti I, Nepita EV, Maserati R, Ferrari A, et al. Prevalence of SARS-CoV-2 specific neutralising antibodies in blood donors from the Lodi Red Zone in Lombardy, Italy, as at 06 April 2020. *Eurosurveillance* 2020;25:2001031.
- Pollán M, Pérez-Gómez B, Pastor-Barriuso R, Oteo J, Hernán MA, Pérez-Olmeda M. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet* 2020;396:535–44, doi:[http://dx.doi.org/10.1016/S0140-6736\(20\)31483-5](http://dx.doi.org/10.1016/S0140-6736(20)31483-5).
- Poustchi H, Darvishian M, Mohammadi Z, Shayanrad A, Delavari A, Bahador-imonfared A. SARS-CoV-2 antibody seroprevalence in the general population and high-risk occupational groups across 18 cities in Iran: a population-based cross-sectional study. *Lancet Infect Dis* 2020;21(4):473–81, doi:[http://dx.doi.org/10.1016/S1473-3099\(20\)30858-6](http://dx.doi.org/10.1016/S1473-3099(20)30858-6).
- Qutob N, Awartani F, Salah Z, Asia M, Khader IA, Herzallah K, et al. Seroprevalence of SARS-CoV-2 in the West Bank region of Palestine: a cross-sectional seroepidemiological study. *BMJ Open* 2021;11:.
- Raoult D. How useful is serology for COVID-19?. *Int J Infect Dis* 2021;102:170–1, doi:<http://dx.doi.org/10.1016/j.ijid.2020.10.058>.
- Rosenberg ES, Tesoriero JM, Rosenthal EM, Chung R, Barranco MA, Styer LM, et al. Cumulative incidence and diagnosis of SARS-CoV-2 infection in New York. *Ann Epidemiol* 2020;48:23–9.
- Rostami A, Sepidarkish M, Leeftang MMG, Riahi SM, Nourollahpour Shiadeh M, Esfandyari S, et al. SARS-CoV-2 seroprevalence worldwide: a systematic review and meta-analysis. *Clin Microbiol Infect* 2020;., doi:<http://dx.doi.org/10.1016/j.cmi.2020.10.020> S1198-743X(20)30651-0.
- Saeed S, Drews SJ, Pambrun C, Yi Q-L, Osmond L, O'Brien SF. SARS-CoV-2 seroprevalence among blood donors after the first COVID-19 wave in Canada. *Transfusion* 2021;61(3):862–72, doi:<http://dx.doi.org/10.1111/trf.16296> n/a.
- Sam I-C, Chong YM, Tan CW, Chan YF. Low postpandemic wave SARS-CoV-2 seroprevalence in Kuala Lumpur and Selangor, Malaysia. *J Med Virol* 2021;93:647–8, doi:<http://dx.doi.org/10.1002/jmv.26426>.
- Silveira MF, Barros AJ, Horta BL, Pellanda LC, Victora GD, Dellagostin OA, et al. Population-based surveys of antibodies against SARS-CoV-2 in Southern Brazil. *Nat Med* 2020;26:1196–9.
- Slot E, Hogema BM, Reusken CBEM, Reimerink JH, Molier M, Karregat JHM, et al. Low SARS-CoV-2 seroprevalence in blood donors in the early COVID-19 epidemic in the Netherlands. *Nat Commun* 2020;11:5744, doi:<http://dx.doi.org/10.1038/s41467-020-19481-7>.
- Sood N, Simon P, Ebner P, Eichner D, Reynolds J, Bendavid E, et al. Seroprevalence of SARS-CoV-2-specific antibodies among adults in Los Angeles County, California, on April 10–11, 2020. *JAMA* 2020;323:2425–7.
- Stringhini S, Wisniak A, Piumatti G, Azman AS, Lauer SA, Baysson H, et al. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Geneva, Switzerland (SEROCoV-POP): a population-based study. *Lancet* 2020;396:313–9.
- Sughayer MA, Mansour A, Nuirat AA, Souan L, Ghanem M, Siag M. The effect of strict lock down measures on Covid-19 seroprevalence rate and herd immunity. *medRxiv* 2020;., doi:<http://dx.doi.org/10.1101/2020.06.06.20123919> [Preprint] 2020.06.06.20123919.
- Sutton M, Cieslak P, Linder M. Notes from the field: seroprevalence estimates of SARS-CoV-2 infection in convenience sample – Oregon, May 11–June 15, 2020. *Morb Mortal Wkly Rep* 2020;69:1100.
- Uyoga S, Adetifa IMO, Karanja HK, Nyagwange J, Tuju J, Wanjiku P. Seroprevalence of anti-SARS-CoV-2 IgG antibodies in Kenyan blood donors. *Science* 2021;371:79, doi:<http://dx.doi.org/10.1126/science.abe1916>.
- Vena A, Berruti M, Adessi A, Blumetti P, Brignole M, Colognato R, et al. Prevalence of antibodies to SARS-CoV-2 in Italian adults and associated risk factors. *J Clin Med* 2020;9:2780.
- WHO. WHO coronavirus disease (COVID-19) dashboard. Geneva: World Health Organization; 2020 Available online: <https://covid19.who.int/>. (last cited: 14 February 2021).
- Xu X, Sun J, Nie S, Li H, Kong Y, Liang M, et al. Seroprevalence of immunoglobulin M and G antibodies against SARS-CoV-2 in China. *Nat Med* 2020;26:1193–5, doi:<http://dx.doi.org/10.1038/s41591-020-0949-6>.
- Younas A, Waheed S, Khawaja S, Imam M, Borhany M, Shamsi T. Seroprevalence of SARS-CoV-2 antibodies among healthy blood donors in Karachi, Pakistan. *Transfus Apher Sci* 2020;59;., doi:<http://dx.doi.org/10.1016/j.transci.2020.102923>.