

Research Article

T_m -Shift Detection of Dog-Derived *Ancylostoma ceylanicum* and *A. caninum*

Yeqi Fu,¹ Mingwei Wang,¹ Xinxin Yan,¹ Auwalu Yusuf Abdullahi ^{1,2} Jianxiong Hang,¹ Pan Zhang,¹ Yue Huang,¹ Yunqiu Liu,¹ Yongxiang Sun,¹ Rongkun Ran,¹ and Guoqing Li ¹

¹Guangdong Provincial Zoonosis Prevention and Control Key Laboratory, College of Veterinary Medicine, South China Agricultural University, Guangzhou 510542, China

²Animal Science Department, Kano University of Science and Technology, Wudil, PMB 3244, Kano 20027, Nigeria

Correspondence should be addressed to Guoqing Li; gqli@scau.edu.cn

Received 26 February 2018; Accepted 4 April 2018; Published 13 May 2018

Academic Editor: Francesca Mancianti

Copyright © 2018 Yeqi Fu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

To develop a T_m -shift method for detection of dog-derived *Ancylostoma ceylanicum* and *A. caninum*, three sets of primers were designed based on three SNPs (ITS71, ITS197, and ITS296) of their internal transcribed spacer 1 (ITS1) sequences. The detection effect of the T_m -shift was assessed through the stability, sensitivity, accuracy test, and clinical detection. The results showed that these three sets of primers could distinguish accurately between *A. ceylanicum* and *A. caninum*. The coefficient of variation in their T_m values on the three SNPs was 0.09% and 0.15% (ITS71), 0.18% and 0.14% (ITS197), and 0.13% and 0.07% (ITS296), respectively. The lowest detectable concentration of standard plasmids for *A. ceylanicum* and *A. caninum* was 5.33×10^{-6} ng/ μ L and 5.03×10^{-6} ng/ μ L. The T_m -shift results of ten DNA samples from the dog-derived hookworms were consistent with their known species. In the clinical detection of 50 fecal samples from stray dogs, the positive rate of hookworm detected by T_m -shift (42%) was significantly higher than that by microscopic examination (34%), and the former can identify the *Ancylostoma* species. It is concluded that the T_m -shift method is rapid, specific, sensitive, and suitable for the clinical detection and zoonotic risk assessment of the dog-derived hookworm.

1. Introduction

Hookworms are common intestinal parasites distributed globally, which can cause serious harm to the health of dogs, cats, and human beings. There are four species of hookworms, that is, *Ancylostoma caninum*, *A. ceylanicum*, *A. braziliense*, and *Uncinaria stenocephala*, that can infect dogs [1]. According to a recent survey, stray dogs are infected mainly by *A. ceylanicum* and *A. caninum* in China [2]. It was reported that *A. ceylanicum* has become the second most common species of hookworm that can infect humans in Asia [3, 4], especially in Southeast Asian countries such as China [5, 6], Japan [7], Malaysia [8], Laos [9], Thailand [10], and India [11]. Occasionally, *A. caninum* can reach adulthood in people [1]. These two hookworms are zoonotic parasites that can parasitize the host's intestinal tract and cause severe symptoms such as intestinal bleeding, anemia, malnutrition,

and dermatitis [12, 13]. Therefore, it is very important to establish a rapid and accurate identification method for the two dog-derived hookworms for the prevention and control of hookworm disease.

Fecal examination is a traditional method for detecting hookworms, but it is easily missed under mild infection. Because of morphological similarities of different hookworm eggs, it is difficult to identify them only by their appearance. Presently, many molecular detection technologies have been applied to the identification of hookworms, such as specific polymerase chain reaction (PCR), multiplex polymerase chain reaction (multi-PCR), restriction fragment length polymorphism (RFLP), and high resolution melting (HRM) [14–16]. However, there are still some shortcomings, such as being unable to detect a large quantity of samples at the same time, complicated operation, and high cost.

Single nucleotide polymorphism (SNP) has become a molecular genetic marker of the third generation after RFLP and microsatellite sequence (MS) [17, 18]. The T_m -shift based on the SNP is a new molecular detection method, where two specific primers and a reverse primer are designed as follows: the 3'-end of specific primer binds to SNP loci, and two GC-rich sequences with unequal length and no pairing with the template are added at the 5'-end, respectively. Then, PCR products of samples are detected according to different peaks of melting curve in different T_m values [19]. The T_m -shift method can satisfy the high-throughput research and has efficient, accurate, and inexpensive advantages [20]. So far, this method has been applied in many fields including medicine, microbiology, and aquatic biology [21–23]. Recently, we established T_m -shift genotyping method for detection of cat-derived *Giardia lamblia* assemblages A and F [24], which has developed a new genotyping method for veterinary parasitic protozoa.

This study developed a T_m -shift method for detecting *A. ceylanicum* and *A. caninum* from dogs based on three SNP loci of rDNA ITS1 sequences of two hookworms, in order to establish a rapid and accurate method for identifying the species of dog-derived hookworms which could provide a new technical means for clinical detection of hookworms and zoonotic risk assessment.

2. Materials and Methods

2.1. Source of Samples. Adult *A. ceylanicum* and *A. caninum* samples were isolated and identified by Liu et al. [2, 6], preserved in 75% alcohol, and stored in our laboratory. Fecal samples were collected from stray dogs of pet shelters in Shantou, Foshan, and Shaoguan districts of Guangdong province in China, preserved in 2.5% potassium dichromate, and stored at -20°C for use.

2.2. Genomic DNA Extraction. The adult hookworms preserved were repeatedly washed with double-distilled water (ddH_2O), and total genomic DNA was extracted using the Wizard[®] SV Genomic DNA Purification System (Promega, Guangzhou, China) according to the manufacturer's protocols. All clinical fecal samples were dissolved in ddH_2O and then pretreated with 5 cycles of heating at 100°C for 5 min, followed by immediate freezing at -80°C for 5 min. Genomic DNA of clinical fecal samples was extracted using Stool DNA extraction kit (Omega, Guangzhou, China) according to the manufacturer's protocols. Extracted DNAs were then stored at -20°C for use.

2.3. PCR Amplification of ITS1 Sequence. The ITS1 sequences of *A. ceylanicum* and *A. caninum* were amplified using a forward primer AF (5'-CTTTGTCGGGAAGGTTGG-3') and a reverse primer AR (5'-TTCACCACTCTAAGCGTCT-3') designed by Liu et al. [15]. The predictive amplification fragment was 404 bp. Polymerase chain reactions were performed in $25\ \mu\text{L}$, including $12.5\ \mu\text{L}$ of Premix Ex-Taq polymerase (TaKaRa, Dalian, China), $9.5\ \mu\text{L}$ of ddH_2O , $0.5\ \mu\text{L}$ of each primer AF/AR ($50\ \mu\text{mol/L}$), and $2\ \mu\text{L}$ of DNA sample. PCR cycling parameters were as follows: 1 cycle at 94°C for

5 min and then 35 cycles at 94°C for 30 sec, at 61.5°C for 30 sec, and at 72°C for 45 sec, followed by 1 cycle at 72°C for 7 min. The PCR products were analysed by gel electrophoresis in 1.5% agarose gels, stained with 0.2 mg/ml ethidium bromide, and visualized on a UV transilluminator.

2.4. Preparation of Standard Plasmids. The purified PCR products were cloned in *Escherichia coli* and connected with pMD18-T (TaKaRa, Dalian, China) and then transferred into DH5 α Competent Cells (TaKaRa, Dalian, China). Positive clones were screened by bacterial PCR and sent to Shenggong Corporation (Shanghai, China) for sequencing. The plasmid DNAs were extracted using the Plasmid Kit (Omega, Guangzhou, China). These plasmids containing ITS1 sequence of *A. ceylanicum* and *A. caninum* were named AceP and AcaP, respectively. The A260/A280 value and concentration of plasmids ($1\ \mu\text{L}$) were measured using ultra-micro-spectrophotometer, where A260/A280 value and optimal concentration should be 1.8~2.0 and $50\ \text{ng}/\mu\text{L}$. These plasmids were stored at -20°C for use.

2.5. Establishment of T_m -Shift Method Based on SNP. Three sets of T_m -shift primers (including two forward specific primers and one common reverse primer) based on three SNPs (ITS71, ITS197, and ITS296) were designed by software Primer Premier 5.0, referencing to two sequences (KM066110.1, LC177192.1) downloaded from GenBank. Their sequence composition and predictive amplification fragment are shown in Table 1. The above-mentioned primers were synthesized by Shenggong Corporation (Shanghai, China), diluted with sterile double-distilled water at a final concentration of $10\ \text{pmol}/\mu\text{L}$, and stored at -20°C for use. PCR amplification and T_m -shift reaction were performed once in Rotor-Gene Q. The reaction system was performed in $20\ \mu\text{L}$, including $10\ \mu\text{L}$ of SYBR[®]Premix Ex-Taq[™] II (2x), $0.4\ \mu\text{L}$ of long-tail primer, $0.4\ \mu\text{L}$ of short-tail primer, $0.8\ \mu\text{L}$ of common reverse primer, $7.4\ \mu\text{L}$ of ddH_2O , and $1.0\ \mu\text{L}$ of plasmid. PCR cycling parameters were as follows: initial denaturation at 95°C for 5 min and then 40 cycles at 95°C for 10 sec and 63°C for 30 sec. Melting process was from 70 to 95°C at the rate of $0.5^{\circ}\text{C}/\text{sec}$.

2.6. Stability, Sensitivity, and Accuracy Test. The detection effect of the T_m -shift was assessed through the stability, sensitivity, and accuracy test. We tested the reproducibility of two known standard plasmids (AceP and AcaP). Intra-assays were tested seven times, and interassays were tested three times for each plasmid. For detection of the sensitivity of the T_m -shift method, two standard plasmids (AceP and AcaP) were diluted ten times and detected according to the concentration of $1:10\sim 1:10^8$. To evaluate the T_m -shift method's accuracy, we detected ten samples with known hookworm species. All reaction system and cycling parameters were the same as mentioned above.

2.7. Clinical Detection. Fifty fecal samples from the stray dogs were examined by the flotation technique with saturated zinc sulfate and the T_m -shift method. T_m -shift classification results were further confirmed by DNA sequencing.

TABLE 1: Primers for T_m -shift method based on five SNPs.

Primer	Nucleotide sequence (5'-3')	Product length (bp)
ITS71TF(Aca)	<u>GCGGCG</u> CTATGTGCAGCAAGAGT	
ITS71CF(Ace)	<u>GCGGGCAGGGCGGG</u> CTATGTGCAGCAAGAGC	113
ITS71R	ACAAGCAGTAAGGCGGCATTCA	
ITS197GF(Aca)	<u>GCGGCG</u> TGAGCATTAGGCTAACGCCTG	
ITS197AF(Ace)	<u>GCGGGCAGGGCGGG</u> TGAGCATTAGGCTAACGCCTA	114
ITS197R	ACGATTCTGCAAACATTAACGTAAAAAGT	
ITS296TF(Ace)	<u>GCGGCG</u> TTTGCAGAATCGTGACTTT	
ITS296GF(Aca)	<u>GCGGGCAGGGCGGG</u> TTTGCAGAATCGTGACTTG	127
ITS296R	TTCACCACTCTAAGCGTCT	
ITS26AF(Aca)	<u>GCGGCG</u> GGAAGTTGGGAGTATCA	
ITS26GF(Ace)	<u>GCGGGCAGGGCGGG</u> GGAAGTTGGGAGTATCG	121
ITS26R	GTCTAAAGCTCAGCGAAAC	
ITS48AF(Aca)	<u>GCGGCG</u> CCGTTACAGCCCTACGTA	
ITS48GF(Ace)	<u>GCGGGCAGGGCGGG</u> CCGTTACAGCCCTACGTG	116
ITS48R	ATGCAATGCTCATCAAGTC	

The GC tails are bold and underlined. Primers at ITS26 and ITS48 failed to distinguish between *A. ceylanicum* and *A. caninum*.

TABLE 2: The stability of T_m -shift method.

Repeat	ITS71 (T_m)		ITS197 (T_m)		ITS296 (T_m)	
	AcaP	AceP	AcaP	AceP	AcaP	AceP
First ($n = 7$)	86.1 ± 0.1°C	88.0 ± 0.1°C	85.1 ± 0.1°C	85.9 ± 0.1°C	85.0 ± 0.1°C	84.0 ± 0.1°C
Second ($n = 7$)	86.1 ± 0.2°C	88.1 ± 0.1°C	85.1 ± 0.1°C	85.9 ± 0.1°C	85.0 ± 0.1°C	84.0 ± 0.2°C
Third ($n = 7$)	86.1 ± 0.1°C	87.9 ± 0.1°C	85.1 ± 0.2°C	85.9 ± 0.1°C	85.1 ± 0.1°C	84.1 ± 0.1°C
Average ($n = 21$)	86.10°C	88.00°C	85.10°C	85.90°C	85.03°C	84.03°C
CV	0.15%	0.09%	0.14%	0.18%	0.07%	0.13%



FIGURE 1: PCR amplification of the ITS1 sequence of two hookworms. MDL-500 DNA marker; 1: positive control; 2: *A. ceylanicum*; 3: *A. caninum*; 4: negative control.

3. Results

3.1. Amplification of ITS1 Fragment and Preparation of Plasmids. The amplified ITS1 fragment from two genomic DNAs of dog-derived *A. ceylanicum* and *A. caninum* was 404 bp long (Figure 1) and the generated sequence data were submitted to GenBank (accession numbers: MG733994, MG733993). BLAST analysis indicated the highest similarity (100%) with *A. ceylanicum* from Japan (LC036567) and the

highest similarity (99%) with *A. caninum* from Australia (KP844730). Thus, two hookworms were identified as *A. ceylanicum* and *A. caninum*. The A260/A280 values of positive plasmids AceP and AcaP were between 1.8 and 2.0, respectively. Their concentrations were 53.3 ng/μL and 50.3 ng/μL, respectively.

3.2. Detection of qPCR- T_m -Shift. The standard curves of T_m -shift based on ITS71, ITS197, and ITS296 for standard plasmid AceP and AcaP are shown in Figure 2. The result showed that T_m -shift method based on three SNPs could distinguish between *A. ceylanicum* and *A. caninum*. Software analysis showed that the T_m values of AceP (*A. ceylanicum*) and AcaP (*A. caninum*) in three sets of primers were 88.0°C and 86.0°C (ITS71), 86.0°C and 85.0°C (ITS197), and 84.0°C and 85.0°C (ITS296), respectively.

3.3. Stability, Sensitivity, and Accuracy. The stability test results are shown in Table 2. In primer ITS71, the coefficient of variation (CV) of AceP and AcaP melting temperature (T_m) was 0.15 and 0.09%. In primer ITS197, the CV of AceP and AcaP melting temperature was 0.14 and 0.18%. And in primer ITS296, the CV of AceP and AcaP melting temperature was

TABLE 3: The sensitivity of T_m -shift method.

Dilution	ITS71 (T_m)		ITS197 (T_m)		ITS296 (T_m)	
	AcaP	AceP	AcaP	AceP	AcaP	AceP
1:10 ¹	86.1°C	88.0°C	85.0°C	86.0°C	85.0°C	83.9°C
1:10 ²	85.9°C	88.1°C	85.0°C	86.0°C	85.0°C	84.0°C
1:10 ³	86.0°C	88.1°C	85.1°C	86.0°C	85.0°C	84.0°C
1:10 ⁴	86.0°C	88.0°C	85.0°C	86.0°C	85.0°C	84.0°C
1:10 ⁵	86.1°C	88.1°C	84.9°C	86.1°C	85.0°C	83.9°C
1:10 ⁶	86.1°C	88.0°C	85.1°C	86.1°C	84.9°C	83.9°C
1:10 ⁷	86.0°C	88.0°C	85.1°C	86.1°C	85.0°C	83.9°C
1:10 ⁸	—	—	—	—	—	—

—: not detected.

TABLE 4: The accuracy of T_m -shift method.

Sample	Species	ITS71 (T_m)	ITS197 (T_m)	ITS296 (T_m)	GenBank
D22	<i>A. caninum</i>	86.0°C	85.0°C	85.0°C	KC755016
D55	<i>A. ceylanicum</i>	88.0°C	86.0°C	84.0°C	KC755015
D60	<i>A. ceylanicum</i>	88.0°C	85.9°C	83.9°C	KC755020
D74	<i>A. ceylanicum</i>	88.0°C	86.0°C	84.0°C	KC755021
D23	<i>A. caninum</i>	86.1°C	85.0°C	85.1°C	KC755017
D28	<i>A. caninum</i>	86.0°C	85.1°C	85.0°C	KC755018
D34	<i>A. caninum</i>	86.1°C	85.1°C	85.0°C	KC755019
D79	<i>A. ceylanicum</i>	88.0°C	86.0°C	84.0°C	KC755022
D23	<i>A. ceylanicum</i>	87.9°C	86.0°C	84.0°C	KF279132
D32	<i>A. ceylanicum</i>	88.0°C	86.0°C	84.0°C	KF279133

Samples D22, D23, D28, D34, D55, D60, D74, and D79 were isolated and identified by Liu et al. [2]. Samples G23 and G32 were isolated and identified by Liu et al. [6].

0.07 and 0.13%. When AceP and AcaP samples were diluted to 1:10⁷ (5.33×10^{-6} ng/ μ L and 5.03×10^{-6} ng/ μ L), the T_m -shift method was still able to identify both of them. However, when the samples were diluted to 1:10⁸, all of them could not be determined (Table 3). A total of ten known hookworm samples from dogs were randomly selected and detected. The T_m -shift detection results based on ITS71, ITS197, and ITS296 were identical to their known hookworm species (Table 4).

3.4. Clinical Detection. Seventeen out of fifty fecal samples from dogs were microscopically positive for hookworm but twenty-one of them were positive for hookworm using T_m -shift method, where fourteen were identified as *A. ceylanicum* and seven as *A. caninum*. The melting curves and detection results are shown in Figure 3. All detection results were identical to their sequencing results.

4. Discussion

Dogs are important reservoir host to zoonotic hookworms *A. ceylanicum* and *A. caninum*. Genetic evolution analysis indicated that dog-derived *A. ceylanicum* and human-derived *A. ceylanicum* from China [25], Malaysia [26], Kampuchea [27], and India [13] have the closest genetic relationship, and the infection rate of *A. ceylanicum* between humans and dogs had a strong correlation in the same region, as well as confirming the possibility of transmission of *A. ceylanicum*

between humans and animals [28]. *A. caninum* mainly infects dogs and cats and sometimes infects people. Dogs are both companion animals for humans and susceptible animals to zoonotic hookworms. With the development of economy and improvement of life quality, the number of people raising dogs has been increasing. Therefore, the establishment of T_m -shift method for detection of *A. ceylanicum* and *A. caninum* has important significance in the prevention and control of zoonotic ancylostomiasis.

Currently, the species identification of hookworm is mainly dependent on molecular means such as specific PCR, multi-PCR, RFLP, and HRM [14–16]. However, there are still some limitations. The conventional PCR can only detect a single pathogen. Although multi-PCR can simultaneously detect multiple pathogens, it is necessary to design more primers and the PCR conditions are demanding. The RFLP technology can only select specific endonuclease, and the operation is tedious and time-consuming, with low sensitivity for false negatives. HRM is similar to the T_m -shift method through test of the melting curve of PCR products. But it is based on the different GC content and base complementarity in the same DNA sequence to detect samples. When difference bases are counteracted in the sequence, two peaks of melting curve are difficult to distinguish and easily cause detection failure [29]. Compared with specific PCR, multi-PCR, and RFLP, the T_m -shift method can detect 36 or 72 samples simultaneously. Its closed tube operation can greatly






FIGURE 2: Standard curves of T_m -shift for two hookworm standard plasmids based on ITS71 (a), ITS197 (b), and ITS296 (c). AceP: *A. ceylanicum* standard plasmid; AcaP: *A. caninum* standard plasmid; Neg: negative control.

reduce the risk of contaminated samples. Also, there is no need to use the poisonous reagent ethidium bromide (EB), which thus can reduce pollution to the environment and ensure safety of the researchers. The T_m -shift method has clear price advantage compared with HRM, because of the use of instruments for ordinary fluorescence PCR and fluorescent dyes for ordinary SYBR Green I rather than saturation dyes Eva Green and LC Green. In addition, the detection efficiency is improved through amplification of the target fragment by two-step method.

Although the T_m -shift detection method has its unique advantages, it also has some limitations. It has higher demand of two specific primers, whose T_m value should be 59~62°C, and their length should be controlled in 15~22 bp. In order to give better amplification efficiency of Taq polymerase, the gap of the T_m values between the reverse primer and the specific primers (before joining at the 5'-end sequence) must be maintained between 4 and 5°C. It is due to the high screening conditions of specific primers that some SNP loci failed to

design suitable primers. Among five SNP loci (Table 1) originally designed in this experiment, two loci (ITS26 and ITS48) failed to distinguish between *A. ceylanicum* and *A. caninum*, thus leading to the failure of detection. Another three SNP loci (ITS71, ITS197, and ITS296) could distinguish between two species of hookworms. The T_m value of the melting curve between *A. ceylanicum* and *A. caninum* differed at 2°C on ITS71 but 1°C on ITS197 and ITS296. The ITS71 locus identification had the best effect. Among 50 dog-derived fecal samples detected, 21 samples were positive for hookworms by T_m -shift method (14 *A. ceylanicum* and 7 *A. caninum*), which had 4 more positive samples than microscopic examination (17 positive samples), and can accurately identify the species of hookworms from stray dogs.



In conclusion, T_m -shift method based on three SNP loci of ITS1 sequences of *A. ceylanicum* and *A. caninum* was used for the first time for their classification. The successful identification of two hookworms indicates that the method is specific, sensitive, economical, practical, and safe, which



Sample No.	Colour	Tm (°C)	Species
AcaP	Green	86.0°C	Ancylostoma caninum
Acep	Red	87.9°C	Ancylostoma ceylanicum
Dog1	Green	86.1°C	Ancylostoma caninum
Dog2	Red	87.9°C	Ancylostoma ceylanicum
Dog3	Red	87.8°C	Ancylostoma ceylanicum
Dog4	Red	87.8°C	Ancylostoma ceylanicum
Dog5	Green	85.9°C	Ancylostoma caninum
Dog6	Red	87.9°C	Ancylostoma ceylanicum
Dog7	Red	87.8°C	Ancylostoma ceylanicum
Dog8	Red	87.9°C	Ancylostoma ceylanicum
Dog9	Red	87.8°C	Ancylostoma ceylanicum
Dog10	Red	87.9°C	Ancylostoma ceylanicum

Sample No.	Colour	Tm (°C)	Species
AcaP	Green	86.0°C	Ancylostoma caninum
Acep	Red	88.0°C	Ancylostoma ceylanicum
Dog11	Green	86.1°C	Ancylostoma caninum
Dog12	Red	88.0°C	Ancylostoma ceylanicum
Dog13	Red	88.0°C	Ancylostoma ceylanicum
Dog14	Green	86.0°C	Ancylostoma caninum
Dog15	Green	86.1°C	Ancylostoma caninum
Dog16	Red	87.9°C	Ancylostoma ceylanicum
Dog17	Red	88.0°C	Ancylostoma ceylanicum
Dog18	Green	86.1°C	Ancylostoma caninum
Dog19	Red	88.0°C	Ancylostoma ceylanicum
Dog20	Red	88.0°C	Ancylostoma ceylanicum
Dog21	Green	86.0°C	Ancylostoma caninum

(a)



Sample No.	Colour	Tm (°C)	Species
AcaP	Blue	85.0°C	Ancylostoma caninum
Acep	Red	86.0°C	Ancylostoma ceylanicum
Dog1	Blue	85.1°C	Ancylostoma caninum
Dog2	Red	85.9°C	Ancylostoma ceylanicum
Dog3	Red	85.9°C	Ancylostoma ceylanicum
Dog4	Red	86.0°C	Ancylostoma ceylanicum
Dog5	Blue	85.0°C	Ancylostoma caninum
Dog6	Red	85.9°C	Ancylostoma ceylanicum
Dog7	Red	85.9°C	Ancylostoma ceylanicum
Dog8	Red	85.9°C	Ancylostoma ceylanicum
Dog9	Red	86.0°C	Ancylostoma ceylanicum
Dog10	Red	85.9°C	Ancylostoma ceylanicum

Sample No.	Colour	Tm (°C)	Species
AcaP	Blue	85.0°C	Ancylostoma caninum
Acep	Red	86.0°C	Ancylostoma ceylanicum
Dog11	Blue	85.1°C	Ancylostoma caninum
Dog12	Red	86.0°C	Ancylostoma ceylanicum
Dog13	Red	86.0°C	Ancylostoma ceylanicum
Dog14	Blue	85.0°C	Ancylostoma caninum
Dog15	Blue	85.1°C	Ancylostoma caninum
Dog16	Red	85.9°C	Ancylostoma ceylanicum
Dog17	Red	86.0°C	Ancylostoma ceylanicum
Dog18	Blue	85.0°C	Ancylostoma caninum
Dog19	Red	86.0°C	Ancylostoma ceylanicum
Dog20	Red	86.0°C	Ancylostoma ceylanicum
Dog21	Blue	85.0°C	Ancylostoma caninum

(b)

FIGURE 3: Continued.




FIGURE 3: Melting curves of T_m -shift based on ITS71 (a), ITS197 (b), and ITS296 (c) and detection results for 21 hookworm-positive samples.

can provide a new technical means for clinical detection of hookworm and zoonotic risk assessment.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Acknowledgments

This work was supported by a grant from the National Natural Science Foundation of China (Grant no. 31672541) and the Science and Technology Planning Project of Guangdong Province, China (Grant no. 2014A020214005).

References

- [1] D. D. Bowman, S. P. Montgomery, A. M. Zajac, M. L. Eberhard, and K. R. Kazacos, "Hookworms of dogs and cats as agents of cutaneous larva migrans," *Trends in Parasitology*, vol. 26, no. 4, pp. 162–167, 2010.
- [2] Y. J. Liu, G. C. Zheng, P. Zhang et al., "Molecular identification of hookworms in stray and shelter dogs from Guangzhou city, China using ITS sequences," *Journal of Helminthology*, vol. 89, no. 2, pp. 196–202, 2015.
- [3] R. J. Traub, "Ancylostoma ceylanicum, a re-emerging but neglected parasitic zoonosis," *International Journal for Parasitology*, vol. 43, no. 12-13, pp. 1009–1015, 2013.
- [4] R. C. A. Thompson, "Neglected zoonotic helminths: Hymenolepis nana, Echinococcus canadensis and Ancylostoma ceylanicum," *Clinical Microbiology and Infection*, vol. 21, no. 5, pp. 426–432, 2015.
- [5] J. Chen, M. J. Xu, D. H. Zhou, H. Q. Song, C. R. Wang, and X. Q. Zhu, "Canine and feline parasitic zoonoses in China," *Parasites Vectors*, vol. 5, p. 152, 2012.
- [6] Y. Liu, G. Zheng, M. Alsarakibi et al., "The zoonotic risk of ancylostoma ceylanicum isolated from stray dogs and cats in guangzhou, south China," *BioMed Research International*, vol. 2014, Article ID e208759, 2014.
- [7] D. Kaya, M. Yoshikawa, T. Nakatani et al., "Ancylostoma ceylanicum hookworm infection in Japanese traveler who presented chronic diarrhea after return from Lao People's Democratic Republic," *Parasitology International*, vol. 65, no. 6, pp. 737–740, 2016.
- [8] R. Ngui, L. S. Ching, T. T. Kai, M. A. Roslan, and Y. A. L. Lim, "Molecular identification of human hookworm infections in economically disadvantaged communities in Peninsular Malaysia," *The American Journal of Tropical Medicine and Hygiene*, vol. 86, no. 5, pp. 837–842, 2012.
- [9] J. V. Conlan, B. Khamlome, K. Vongxay et al., "Soil-transmitted helminthiasis in Laos: a community-wide cross-sectional study of humans and dogs in a mass drug administration environment," *The American Journal of Tropical Medicine and Hygiene*, vol. 86, no. 4, pp. 624–634, 2012.
- [10] R. J. Traub, T. Inpankaew, C. Sutthikornchai, Y. Sukthana, and R. C. A. Thompson, "PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by

- Ancylostoma ceylanicum* in temple communities in Bangkok,” *Veterinary Parasitology*, vol. 155, no. 1-2, pp. 67–73, 2008.
- [11] R. J. Traub, R. P. Hobbs, P. J. Adams, J. M. Behnke, P. D. Harris, and R. C. A. Thompson, “A case of mistaken identity—reappraisal of the species of canid and felid hookworms (*Ancylostoma*) present in Australia and India,” *Parasitology*, vol. 134, no. 1, pp. 113–119, 2007.
- [12] P. J. Hotez, M. Alvado, and M.-G. Basáñez, “The global burden of disease study 2010: interpretation and implications for the neglected tropical diseases,” *PLOS Neglected Tropical Diseases*, vol. 8, no. 7, p. e2865113-119, 2014.
- [13] S. George, B. Levecke, D. Kattula et al., “Molecular Identification of Hookworm Isolates in Humans, Dogs and Soil in a Tribal Area in Tamil Nadu, India,” *PLOS Neglected Tropical Diseases*, vol. 10, no. 8, Article ID e0004891, 2016.
- [14] R. Ngui, Y. A. L. Lim, and K. H. Chua, “Rapid detection and identification of human hookworm infections through high resolution melting (HRM) analysis,” *PLoS ONE*, vol. 7, no. 7, Article ID e41996, 2012.
- [15] Y. Liu, G. Zheng, M. Alsarakibi et al., “Molecular identification of *Ancylostoma caninum* isolated from cats in southern china based on complete ITS sequence,” *BioMed Research International*, vol. 2013, Article ID 868050, 6 pages, 2013.
- [16] W. Hu, S. Wu, X. Yu et al., “A Multiplex PCR for Simultaneous Detection of Three Zoonotic Parasites *Ancylostoma ceylanicum*, *A. caninum*, and *Giardia lamblia* Assemblage A,” *BioMed Research International*, vol. 2015, Article ID 406168, 6 pages, 2015.
- [17] T. Grodzicker, J. Williams, P. Sharp, and J. Sambrook, “Physical mapping of temperature-sensitive mutations of adenoviruses,” *Cold Spring Harbor Symposia on Quantitative Biology*, vol. 39, no. 1, pp. 439–446, 1975.
- [18] D. Tautz and M. Renz, “Simple sequences are ubiquitous repetitive components of eukaryotic genomes,” *Nucleic Acids Research*, vol. 12, no. 10, pp. 4127–4138, 1984.
- [19] J. Wang, K. Chuang, M. Ahluwalia et al., “High-throughput SNP genotyping by single-tube PCR with Tm-shift primers,” *Bio-Techniques*, vol. 39, no. 6, pp. 885–893, 2005.
- [20] S. H. Zhou, M. Liu, W. X. An et al., “Genotyping of human platelet antigen-15 by single closed-tube T m-shift method,” *International Journal of Laboratory Hematology*, vol. 34, no. 1, pp. 41–46, 2012.
- [21] S. Derzelle, C. Mendy, S. Laroche, and N. Madani, “Use of High-Resolution Melting and Melting Temperature-shift assays for specific detection and identification of *Bacillus anthracis* based on single nucleotide discrimination,” *Journal of Microbiological Methods*, vol. 87, no. 2, pp. 195–201, 2011.
- [22] Y. Huang, S. Nie, S. Zhou et al., “PPARD rs2016520 polymorphism and circulating lipid levels connect with brain diseases in Han Chinese and suggest sex-dependent effects,” *Biomedicine & Pharmacotherapy*, vol. 70, no. C, pp. 7–11, 2015.
- [23] J. J. Ahn, Y. Kim, J. Y. Hong, G. W. Kim, S. Y. Kim, and S. Y. Hwang, “Probe-Based Fluorescence Melting Curve Analysis for Differentiating *Larimichthys polyactis* and *Larimichthys crocea*,” *Food Analytical Methods*, vol. 9, no. 7, pp. 2036–2041, 2016.
- [24] W. Pan, Y. Fu, A. Y. Abdullahi et al., “Development of Tm-shift genotyping method for detection of cat-derived *Giardia lamblia*,” *Parasitology Research*, vol. 116, no. 4, pp. 1151–1157, 2017.
- [25] W. Hu, X. G. Yu, S. Wu et al., “Levels of *Ancylostoma* infections and phylogenetic analysis of cox 1 gene of *A. ceylanicum* in stray cat faecal samples from Guangzhou, China,” *Journal of Helminthology*, vol. 90, no. 4, pp. 392–397, 2016.
- [26] R. Ngui, M. A. K. Mahdy, K. H. Chua, R. Traub, and Y. A. L. Lim, “Genetic characterization of the partial mitochondrial cytochrome oxidase c subunit I (cox 1) gene of the zoonotic parasitic nematode, *Ancylostoma ceylanicum* from humans, dogs and cats,” *Acta Tropica*, vol. 128, no. 1, pp. 154–157, 2013.
- [27] T. Inpankaew, F. Schär, V. Khieu et al., “Simple Fecal Flotation Is a Superior Alternative to Quadruple Kato Katz Smear Examination for the Detection of Hookworm Eggs in Human Stool,” *PLOS Neglected Tropical Diseases*, vol. 8, no. 12, Article ID e3313, 2014.
- [28] R. Ngui, Y. A. L. Lim, R. Traub, R. Mahmud, and M. S. Mistam, “Epidemiological and genetic data supporting the transmission of *Ancylostoma ceylanicum* among human and domestic animals,” *PLOS Neglected Tropical Diseases*, vol. 6, no. 2, Article ID e1522, 2012.
- [29] G. H. Reed, J. O. Kent, and C. T. Wittwer, “High-resolution DNA melting analysis for simple and efficient molecular diagnostics,” *Pharmacogenomics*, vol. 8, no. 6, pp. 597–608, 2007.