

# Developmental Control of the Cell Cycle: Insights from *Caenorhabditis elegans*

Edward T. Kipreos<sup>\*.1</sup> and Sander van den Heuvel<sup>†.1</sup>

<sup>\*</sup>Department of Cellular Biology, University of Georgia, Athens, Georgia 30602-2607, <sup>†</sup>Department of Biology, Science 4 Life Program, Faculty of Science, University of Utrecht, 3584CH, The Netherlands  
ORCID IDs: 0000-0001-9650-3558 (E.T.K.); 0000-0001-9015-7463 (S.v.d.H.)

**ABSTRACT** During animal development, a single fertilized egg forms a complete organism with tens to trillions of cells that encompass a large variety of cell types. Cell cycle regulation is therefore at the center of development and needs to be carried out in close coordination with cell differentiation, migration, and death, as well as tissue formation, morphogenesis, and homeostasis. The timing and frequency of cell divisions are controlled by complex combinations of external and cell-intrinsic signals that vary throughout development. Insight into how such controls determine *in vivo* cell division patterns has come from studies in various genetic model systems. The nematode *Caenorhabditis elegans* has only about 1000 somatic cells and approximately twice as many germ cells in the adult hermaphrodite. Despite the relatively small number of cells, *C. elegans* has diverse tissues, including intestine, nerves, striated and smooth muscle, and skin. *C. elegans* is unique as a model organism for studies of the cell cycle because the somatic cell lineage is invariant. Somatic cells divide at set times during development to produce daughter cells that adopt reproducible developmental fates. Studies in *C. elegans* have allowed the identification of conserved cell cycle regulators and provided insights into how cell cycle regulation varies between tissues. In this review, we focus on the regulation of the cell cycle in the context of *C. elegans* development, with reference to other systems, with the goal of better understanding how cell cycle regulation is linked to animal development in general.

**KEYWORDS** *Caenorhabditis elegans*; cell cycle; cell lineage; DNA replication licensing; proliferation; terminal differentiation; WormBook

## TABLE OF CONTENTS

Abstract	797
Core Regulators of the Cell Cycle	799
Developmental and Tissue Cell Cycle Variants	801
<i>Embryonic cell cycles</i>	802
<i>Somatic larval cell cycles</i>	802
<i>Endocycles</i>	802
<i>Germ cell cycles</i>	803
The Regulation of Cell Cycle Progression During Development	804
<i>Oocyte meiotic arrest and maturation</i>	804
<i>Completion of meiosis</i>	804

*Continued*

Copyright © 2019 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.118.301643>

Manuscript received February 11, 2018; accepted for publication October 10, 2018

Available freely online through the author-supported open access option.

<sup>1</sup>Corresponding authors: Kruytgebouw, Padualaan 8, Utrecht University, 3584CH Utrecht, The Netherlands. E-mail: s.j.l.vandenheuvel@uu.nl; and Department of Cellular Biology, University of Georgia, Athens, GA 30602-2607. E-mail: ekipreos@uga.edu

CONTENTS, *continued*

<i>From meiosis to mitosis during the “oocyte-to-embryo” switch</i>	806
<i>Embryonic cell cycles</i>	806
<i>Timing asynchrony</i>	807
<i>Checkpoint-induced cell cycle arrest in the early embryo</i>	808
<i>Lineage-dependent introduction of GAP phases and quiescence</i>	808
<i>Systemic regulation of cell division during larval development</i>	809
<i>Cell cycle arrest of newly hatched larvae</i>	809
<i>Cell cycle entry during larval development</i>	809
<i>Progression through larval cell divisions</i>	810
<i>The timing of larval development and cell division</i>	810
<i>Dauer development</i>	811
The Control of DNA Replication	811
<i>Replication licensing in larvae</i>	811
<i>Tissue-specific differences in regulating replication: germ cells and embryos</i>	812
<i>Bypassing the DNA replication checkpoint</i>	813
Regulation of the Cell Cycle in Germ Cells	813
<i>Cell cycle regulators in the mitosis vs. meiosis transition</i>	815
<i>Cyclin E expression to maintain germ cells in a mitotic state</i>	815
<i>FBF represses CKI-2 expression to prevent germ cell arrest</i>	816
<i>CRL2–LRR-1 promotes mitotic proliferation and prevents checkpoint activation</i>	816
<i>Regulation of the rate of germ cell proliferation</i>	816
<i>TGF-β promotes the mitotic germ cell fate</i>	816
<i>Insulin/IGF-like signaling promotes germ cell proliferation</i>	817
<i>DAF-18<sup>PTEN</sup> signaling decreases germ cell proliferation in response to unfertilized oocytes</i>	817
<i>Bacterial folates are an exogenous signal to promote germ cell proliferation</i>	817
<i>Volatile bacterial odors increase reproductive rate via neuronal signals</i>	817
<i>The regulation of germ cell numbers upon starvation</i>	818
<i>Dafachronic acid inhibits germ cell proliferation in adults and mediates the adult starvation response</i>	818
The Interplay Between Cell Proliferation, Arrest, and Differentiation	818
<i>Rb/E2F-mediated transcriptional repression</i>	818
<i>Lineage-dependent regulation of the cell cycle</i>	819
<i>Coordinating cell cycle arrest and terminal differentiation</i>	820
<i>Neuronal differentiation</i>	820
<i>Muscle formation</i>	820
<i>Coordinating cell cycle exit with differentiation: SWIISNF chromatin remodelers</i>	820
<i>Combining cell proliferation and differentiation through asymmetric cell division</i>	821
Perspectives	822

IN the late 1980s, studies in yeasts, frogs, and cultured mammalian cells came together in the discovery of cyclin-dependent kinases (CDKs) as the central regulators of the eukaryotic cell division cycle (Dorée and Hunt 2002; Hartwell 2002; Nurse 2002). Although we have since learned an enormous amount, understanding the regulatory networks that link developmental and environmental signals to the cell cycle remains a major challenge. Genetic model systems offer opportunities to discover such connections. *Caenorhabditis elegans* has several features that make this tiny animal attractive for the analysis of cell cycle regulation in a developmental

context. In particular, the ease of genetic analysis, the transparency of its body, and the reproducible pattern of its development facilitate the identification and quantitative characterization of cell cycle regulators. As a consequence, specific cell division phenotypes were described at an early stage, following screens for mutants with abnormal cell lineages (*lin* mutants) (Horvitz and Sulston 1980; Sulston and Horvitz 1981). For example, cells in *lin-5* mutants do not complete M phase, but nevertheless continue subsequent rounds of DNA replication. Conversely, postembryonic precursor cells (“blast cells”) skip DNA replication in *mcm-4* (*lin-6*)

mutants, while initiating mitosis at the normal times. Two other mutants, *cul-1* (*lin-19*) and *lin-23*, undergo supernumerary cell divisions during larval development (Kipreos *et al.* 1996, 2000). Subsequent molecular characterizations revealed how these genes fulfill general cell cycle functions (see below).

Homozygous cell cycle mutants are usually sterile and therefore are obtained from heterozygous mothers. In this situation, cell cycle phenotypes are generally observed during postembryonic development, as the presence of wild-type maternal product allows development through embryogenesis and masks early requirements. Since the discovery of RNA-mediated interference (RNAi) (Guo and Kemphues 1995; Fire *et al.* 1998), knockdown of maternal product has frequently been used to detect the requirements for cell cycle genes in the germline and during early embryogenesis. Many additional developments have facilitated progress, including the use of green fluorescent protein fusions (Chalfie *et al.* 1994) and recent success with clustered regularly interspaced short palindromic repeats (CRISPR)/Cas9-assisted recombineering [reviewed in: Waaijers and Boxem (2014), Dickinson and Goldstein (2016)]. An advanced molecular genetic toolkit is now available, which makes it possible to combine sophisticated genetics, cell biology, biochemistry, and genomics approaches to study cell cycle regulation at single-cell resolution in living animals.

Following pioneering studies in other systems, studies utilizing *C. elegans* confirmed the basic understanding of the core cell cycle machinery [reviewed in Kipreos (2005) and van den Heuvel (2005)]. The *C. elegans* studies also uncovered several novel cell cycle regulators. For instance, the molecular characterization of *cul-1* (*lin-19*), which regulates cell cycle exit, revealed the evolutionarily conserved cullin family (Kipreos *et al.* 1996). Cullin scaffolding proteins form part of CRL (cullin-ring-ligase) E3 ubiquitin ligases, which include SCF (Skp1–cullin–F-box protein), and regulate critical cell cycle functions, among many other cellular functions. The molecular characterization of *lin-5* resulted in the discovery of an evolutionarily conserved LIN-5<sup>NuMA</sup>-based protein complex (Lorson *et al.* 2000; Srinivasan *et al.* 2003). This complex is critical for the generation of microtubule pulling forces that contribute to chromosome segregation and determine the cell cleavage plane by positioning the mitotic spindle. These early examples illustrated the potential of *C. elegans* studies in the discovery of cell cycle control mechanisms that operate in animal development.

*C. elegans* is particularly attractive for discovering universal aspects of cell cycle control, and studying the integration of cell division and development. An important topic is the regulation of cell cycle entry and exit, which is regulated in substantial part during the G1 phase of the cell cycle. In this respect, it is of great importance that the critical regulators of G1 progression (described below) are evolutionarily conserved between *C. elegans* and more complex eukaryotes. This review will broadly cover how the cell cycle is regulated


in *C. elegans*, including progression through variant cell cycles such as meiosis and early embryonic cell cleavages, and the integration of developmental as well as environmental signals with cell cycle entry-and-arrest decisions.

## Core Regulators of the Cell Cycle

Before examining developmental control and tissue variations, this section will summarize some of the basic concepts of cell cycle regulation. Because this information is textbook level and encompasses a large number of studies, we will limit the use of references to quotations of parallels between broadly known regulators and genes with similar functions in *C. elegans*.

The central goals of the cell cycle include the exact duplication of genomic DNA during the S (DNA synthesis) phase and the segregation of chromosomes, as well as cleavage of the cytoplasm during the M (mitotic) phase. The S and M phases are usually separated by GAP phases in which the cells grow, repair DNA, and, when needed, arrest cell cycle progress prior to the next phase. As such, the G1 phase separates the completed M phase from S phase, and the G2 phase separates S and M phases (Figure 1A). Cells in G1 can alternatively commit to go through another cell cycle, enter a temporal cell cycle quiescent state known as G0, or permanently withdraw from the cell cycle.

The transient activation of CDKs at specific cell cycle transitions drives progression through the cell cycle. Active CDKs consist of a catalytic CDK subunit and an associated cyclin protein (Figure 1A). In both budding and fission yeasts, a single CDK is required for progression through “START”, the transition in G1 at which the cell commits to go through a new division cycle, as well as for the transition from G2 into mitosis. At these different transitions, distinct cyclins (G1 or mitotic) activate the yeast cell cycle CDK (CDK1; also known as CDC2 in fission yeast and CDC28 in budding yeast). Multicellular organisms use various cyclins as well as multiple catalytic subunits to regulate cell cycle transitions [reviewed in Sherr (1996)]. Specifically, the closely related CDK4 and CDK6 kinases act in association with D-type cyclins to stimulate cell cycle entry in G1 (Figure 1A). The subsequent activation of CDK2-cyclin E kinases further promotes cell cycle commitment and progression through the G1/S transition. Next, CDK2-cyclin A becomes active during S phase and G2, and CDK1 in association with B-type cyclins controls entry into mitosis. Orthologs of CDK4/6, CDK2, and CDK1 are present as single genes in *C. elegans* (Boxem *et al.* 1999; Liu and Kipreos 2000). Moreover, cyclins of each class are expressed in *C. elegans*, including single D- and E-type cyclins, a single A-type cyclin, several presumed cyclin A-related pseudogenes, and multiple B-type cyclins. While the analysis currently remains incomplete, the temporal expression and cell cycle stage-specific functions of distinct CDK-cyclin combinations appear conserved between *C. elegans* and mammals (Figure 1A) (Kipreos 2005; van den Heuvel 2005; van der Voet *et al.* 2009).



**Figure 1** Key regulators of the cell division cycle. (A) Illustration of the cell division cycle and the multiple cyclin-dependent kinase (CDK) complexes that participate in the regulation of cell cycle control. The CDK catalytic subunits (blue) interact preferentially with one or more subfamilies of cyclin proteins. The circle represents a standard somatic cell cycle with sequential G1, S, G2, and M phases. CDK-cyclin complexes (mammalian names, larger font) are positioned according to their approximate time of activity in mammalian cells. The closest *C. elegans* homologs (names listed, smaller font) appear to share conserved functions. (B) Generic regulation of CDK activity. CDKs are positively regulated by cyclin association, activating phosphorylation (by CAK/Cdk7), and the removal of inhibitory phosphorylation (by phosphatases of the CDC25 family). CDKs are negatively regulated by inhibitory phosphorylation by Wee1/Myt1 kinases, and cyclin degradation through CRU/SCF and/or APC/C E3 ubiquitin ligases. In addition (not indicated), association with CDK-inhibitory proteins (CKIs) prevents CDK activity. This includes CKIs of the Cip/Kip family, known as CKI-1 and CKI-2 in *C. elegans*. See text for further details.

Several positive and negative regulators control the activation and inactivation of CDK-cyclin complexes. This includes activating and inhibitory phosphorylations of the catalytic subunit, transcriptional regulation and protein

degradation of cyclins, dephosphorylation of CDKs and their substrates, and expression as well as destruction of small CDK-inhibitory proteins (CKIs). In addition to binding cyclin, CDK activation requires the phosphorylation of a threonine residue in the kinase activation loop (T loop). This phosphorylation is carried out by a distant CDK-cyclin pair, CDK7 in association with cyclin H (Fisher and Morgan 1994) (Figure 1B). CDK7 has dual functions as a CDK-activating kinase and polymerase II C-terminal domain kinase, which appear conserved between *C. elegans* and mammals (Wallenfang and Seydoux 2002). In general, the regulatory phosphorylations of CDK subunits and ubiquitin-dependent degradation of cell cycle proteins involve evolutionarily conserved mechanisms, while the specific transcriptional regulators and CKIs differ substantially between single-cell eukaryotes and metazoans.

The regulation of CDKs by inhibitory phosphorylation has been best described for CDK1. Kinases of the Wee1/Myt1 family are responsible for the inhibitory CDK phosphorylation. These kinases phosphorylate a threonine-tyrosine residue pair in the ATP-binding loop, which likely interferes with CDK substrate binding and phosphate alignment (Jeffrey *et al.* 1995) (Figure 1B). Phosphatases of the CDC25 family remove the inhibitory phosphorylations, which is sometimes used to control the timing of CDK1 activation. At the G2/M transition, the activation of CDK1 by CDC25 initiates a positive feedback loop, in which CDK1-mediated phosphorylation promotes activation of its positive regulator CDC25 and inactivation of its negative regulator, the Wee1 kinase, to rapidly achieve complete CDK1 activation. *C. elegans* expresses two Wee1/Myt1-related kinases: WEE-1.1 and WEE-1.3 (*wee-1.2* is a pseudogene). These kinases appear to inhibit CDK-1 in a lineage-specific manner (Wilson *et al.* 1999; Burrows *et al.* 2006). Likewise, the *C. elegans* genome encodes four different CDC25-family phosphatases, with some tissue specificity at least for CDC-25.1 and CDC-25.2 (Ashcroft *et al.* 1999; Ashcroft and Golden 2002; Kim *et al.* 2010; Lee *et al.* 2016). Developmental roles for CDC25 family members have been well documented in other animal systems, in particular *Drosophila* embryos [reviewed in: Yuan *et al.* (2016)]. Thus, regulation of CDK1 by inhibitory phosphorylation is conserved among eukaryotes, while distinct Wee1/Myt1 kinases and CDC25-related phosphatases serve developmental functions in metazoans.

The sudden activation of CDK1-mitotic cyclin complexes induces a dramatic reorganization of the cell at the onset of mitosis, leading to the formation of a bipolar spindle and chromosome alignment at the metaphase plate. Mitotic CDK phosphorylation also activates the anaphase-promoting complex/cyclosome (APC/C), a multi-subunit E3 ubiquitin ligase, in association with its coactivator CDC20 (known as Fizzy in *Drosophila* and FZY-1 in *C. elegans*; Kitagawa *et al.* 2002) (Figure 1B). The APC/C-CDC20 fulfills two critical functions by inducing the degradation of securin, a protein that inhibits the proteolytic enzyme separase, and mitotic cyclins (Nasmyth 2005). Securin degradation leads to the activation of separase, which then cleaves the cohesion ring

complexes that hold sister chromosomes together. Through this pathway, the APC/C–CDC20 triggers sister chromosome segregation in anaphase. The degradation of mitotic cyclins induced by the APC/C–CDC20 leads to the inactivation of CDK1 and exit from mitosis. Phosphatases that dephosphorylate CDK1 substrates assist this process. Following CDK inactivation, a CDC20/Fizzy-related APC/C coactivator known as Cdh1 or Fizzy-related (FZR-1 in *C. elegans*) replaces CDC20/Fzy, and maintains APC/C activity and mitotic cyclin degradation during late mitosis, and into the next G1 phase (Fay *et al.* 2002; The *et al.* 2015).

Entry into the next cell cycle requires new cyclin synthesis and activation of G1 CDKs. The transcription of G1 cyclin genes is often controlled by cell external factors, while subsequent expression of G1/S, S phase, and M phase cyclins is generally under cell-intrinsic control. Heterodimeric transcription factors of the E2F/DP family (E2 promoter-binding Factor/Dimerization Partner protein) are critical in the regulation of cell cycle genes in metazoans [reviewed in van den Heuvel and Dyson (2008)]. Depending on the specific E2F subunit, the E2F/DP dimer (also named E2F) primarily acts as a transcriptional activator or a repressor. Proteins of the retinoblastoma (Rb) tumor suppressor family bind and block activating E2Fs, and act in association with repressor E2Fs to inhibit cell cycle gene expression. Upon induction of cyclin D expression by external factors, G1 CDKs (CDK4/6 associated with a D-type cyclin) become active and initiate the phosphorylation of Rb-family proteins. This is thought to reduce Rb-mediated transcriptional repression and to allow the expression of cell cycle genes that include cyclin E, cyclin A, and CDK2. CDK2-cyclin E and CDK2-cyclin A further phosphorylate and inhibit Rb. In this way, Rb/E2F and cyclin E form part of a double-negative feedback loop that controls an all-or-none decision to enter S phase. The components and mechanisms of this pathway are conserved in *C. elegans*, with critical roles for cyclin D (CYD-1) and CDK-4 (CDK4/6) in cell cycle entry (Park and Krause 1999; Boxem and van den Heuvel 2001).

The temporal accumulation and inactivation of CKIs provides important additional regulation of cell cycle entry. Vertebrates express two distinct classes of CKIs, the INK family members that specifically block CDK4/CDK6 kinases and the CIP/KIP class that primarily inhibits CDK2/cyclin E [reviewed in Vidal and Koff (2000)]. Induced expression of CIP/KIP inhibitors in response to external or cell intrinsic signals causes cell cycle arrest in a variety of conditions. Removal of CIP/KIP inhibitors at the G1/S transition can be triggered by phosphorylation mediated, among others, by CDK2 (Lu and Hunter 2010). Once phosphorylated, p27<sup>KIP1</sup> is a substrate for SCF-induced ubiquitin-dependent protein degradation. SKP2, the substrate specificity factor of the SCF E3 ligase, is itself a target for the APC/C–FZR1 E3 ligase in mammals (Bashir *et al.* 2004; Wei *et al.* 2004). Promoting SKP2 degradation, and thereby p27<sup>KIP1</sup> accumulation, is one of the mechanisms by which APC/C–FZR1 inhibits cell cycle entry.

All eukaryotes use multiple mechanisms to prevent inappropriate cell cycle entry, but the exact molecules and their relative contributions vary. Worms lack INK-type CDK inhibitors, like flies, but express two CDK inhibitors of the CIP/KIP family (Hong *et al.* 1998; Feng *et al.* 1999). CKI-1 and CKI-2 are equally similar to mammalian p21<sup>CIP1</sup> and p27<sup>KIP1</sup>, and *Drosophila* dacapo. However, only CKI-1 acts as a general inhibitor of cell cycle entry, whose inactivation leads to supernumerary cell divisions (Hong *et al.* 1998; Fukuyama *et al.* 2003). By contrast, CKI-2 has only a limited contribution (Buck *et al.* 2009). Genetic redundancies indicate that CKIs and other negative regulators of the cell cycle (LIN-35<sup>Rb</sup>, SCF-LIN-23<sup>B-TrCP</sup>, and FZR-1<sup>FZR1/Cdh1</sup>) act in parallel (Boxem and van den Heuvel 2001; Fay *et al.* 2002). For instance, the G1 inhibitory role of APC/C–FZR-1 becomes apparent only when other G1/S regulators are disrupted, and probably does not result from CKI-1<sup>p27</sup> stabilization (Fay *et al.* 2002; The *et al.* 2015).

The *C. elegans* CDC-14 phosphatase also promotes cell cycle quiescence, likely by antagonizing degradation of CKI-1 induced by CDK phosphorylation (Saito *et al.* 2004). Budding yeast Cdc14 antagonizes CDK1 phosphorylation to promote exit from mitosis by supporting the accumulation of a CKI (Sic1) and the degradation of mitotic cyclins (Stegmeier and Amon 2004). Such a role has not been reported for fission yeast or mammalian Cdc14, possibly because in these organisms the PP1 and PP2A phosphatases are more important for counteracting CDK1 phosphorylation (Grallert *et al.* 2015). The fact that specific regulators may be critical in some organisms while not in others may reflect different levels of redundancy, and illustrates how studying the cell cycle in multiple systems helps acquire a deeper understanding.

Several additional regulators collaborate with or antagonize CDK-cyclins to govern cell-cycle events. This includes checkpoint control pathways that, for example, arrest cell cycle progression in response to DNA damage or incomplete alignment of chromosomes at the metaphase plate. Moreover, additional kinases act in concert with CDKs to regulate DNA replication (DDK kinase) and mitosis (Plk1, Aurora A, and Aurora B). These regulators are not described in this overview section, but we have tried to include any developmental functions reported in *C. elegans* in the relevant sections below.

## Developmental and Tissue Cell Cycle Variants

*C. elegans* embryonic and larval somatic cell cycles occur in the context of an almost invariant lineage. Somatic cells divide at set developmental times to produce daughter cells with reproducible cell fates (Sulston and Horvitz 1977; Kimble and Hirsh 1979; Sulston *et al.* 1983). The division of somatic cells is restricted to the embryonic and larval stages of development; and adult-stage somatic cells are postmitotic. In contrast, germ cells divide throughout larval and adult stages. Below, we will briefly describe how cell cycles vary in different tissues or developmental stages. More

in-depth descriptions of these differences can be found in previous reviews (Kipreos 2005; van den Heuvel 2005).

### **Embryonic cell cycles**

“Cleavage-type” cell divisions immediately follow fertilization in many animal embryos, including *C. elegans*. These divisions are rapid and occur without an overall increase in size, so that daughter cells become progressively smaller with each subsequent division. In *C. elegans*, the first two mitotic cell cycles last only 15–20 min, with the timing lengthening progressively as cell divisions continue (Deppe *et al.* 1978; Sulston *et al.* 1983). This short cell cycle timing is likely facilitated by abundant maternal product (mRNA and protein) that is supplied to the oocyte from the hermaphrodite parent. The entire process of embryogenesis is completed in 800 min at 20° to produce 671 cells, of which 113 undergo apoptosis (Sulston *et al.* 1983).

The early embryonic cell cycles are streamlined to include only S phase and mitosis, as occurs in the early embryos of many other animals (Edgar and McGhee 1988). S phase is very short in the early cell cycles and only lasts ~8.5 min in the first cell cycle (Sonneville *et al.* 2012). In other animals, the speed of embryonic S phase is accomplished by initiating DNA replication simultaneously at many more replication origins than in later cell cycles (Méchali 2010). Presumably, a similar strategy allows the very short S phase timing in early *C. elegans* embryos.

G2 phase is the first gap phase to appear during embryogenesis in *C. elegans* embryos. G2 phase is initially observed at the 26-cell stage in the two cells that give rise to the intestine shortly before the cells migrate into the interior of the embryo during the process of gastrulation (Edgar and McGhee 1988).

Cyclin D (*CYD-1*) is required for progression through G1 phase in somatic larval cells (Park and Krause 1999; Boxem and van den Heuvel 2001). Yet in the embryo, only a few, very late cell divisions require cyclin D (Boxem and van den Heuvel 2001; Yanowitz and Fire 2005). This suggests that only these late embryonic cell divisions include a G1 phase. However, it is also possible that embryonic cells have G1 phases that do not require the normal G1 regulatory program; or that cyclin D and *CDK-4* are required specifically for reentering G1 phase from a quiescent (G0) state that is present in larval cell lineages but is absent from most embryonic lineages. Notably, the late-dividing embryonic cells that require *CYD-1* activity have substantially longer time periods separating their previous and final divisions than the majority of embryonic cells, which may therefore include a G1 or G0 phase.

### **Somatic larval cell cycles**

Newly hatched L1-stage larvae have 558 nuclei; some of these nuclei are in syncytial cells, and hence there are slightly less than 558 cells. Fifty-five of the cells in the newly hatched larvae will undergo cell or nuclear divisions during the larval stages. These cells are called postembryonic blast cells and include two germline precursor cells. The 53 somatic blast

cells will generate an additional 403 nuclei in the hermaphrodite (Sulston and Horvitz 1977; Kimble and Hirsh 1979). The majority of the somatic postembryonic blast cells contribute to increasing the number of cells in tissues already present in the embryo, including muscle, nerves, intestine, hypodermis, and scavenger cells known as coelomocytes (Sulston and Horvitz 1977). However, two additional structures are created during the larval stages: the vulva [created by the vulval precursor cells (VPCs)] and the somatic gonad, created by the Z1 and Z4 blast cells (Kimble and Hirsh 1979).


The larval-stage somatic cell divisions have full cell cycles that often include lengthy G0/G1 phases prior to initiating DNA replication, for example in the VPCs, seam cells [which produce cells for the general epidermis (hypodermis/skin)], and intestine cells (Figure 2). The length of the G2 phase varies substantially between lineages. A very short G2 phase separates S phase from mitosis in the seam cell and vulval precursor cell lineages, and a much longer G2 phase of several hours is observed in the intestine (Hedgecock and White 1985; Euling and Ambros 1996; Boxem and van den Heuvel 2001; Zhong *et al.* 2003).

### **Endocycles**

Endoreplication occurs when cells undergo rounds of DNA replication without an intervening mitosis. Two tissues in *C. elegans* undergo endoreplication: the hypodermis and intestine. Unlike in *Drosophila*, in which endocycles often replicate only the euchromatin parts of the genome (Lilly and Duronio 2005), endoreplication in the *C. elegans* hypodermis and intestine appears to involve complete (or largely complete) doubling of the genome in each endoreplicative cycle (Hedgecock and White 1985).

The majority of intestine cells undergo a nuclear division at the L1-to-L2 transition, producing cells with two nuclei. All intestinal nuclei then undergo endoreplication to attain a 4C DNA content. At the next three larval stage transitions, the intestinal nuclei endoreplicate to successively generate DNA contents of 8C, 16C, and 32C (Hedgecock and White 1985).

Hyp7 is a large syncytial cell that contains multiple nuclei and acts as the skin below the cuticle for the majority of the body (with the exclusion of the head and tail). Hyp7 is formed as a syncytium in embryos and the embryonically derived hyp7 nuclei remain 2C throughout development (Hedgecock and White 1985). During the larval stages, nuclei are added to the syncytium from the seam cell lineages, which produce daughter cells that fuse with hyp7 (Figure 2, A and B). The cells destined to fuse with hyp7 undergo one round of DNA replication without mitosis shortly after they are born, and then fuse with the hyp7 syncytium as 4C nuclei (Hedgecock and White 1985). In the adult stage, the hyp7 nuclei undergo additional endoreplication to attain an average DNA content of ~10.7C (Flemming *et al.* 2000). The adult hyp7 nuclei do not display discrete peaks of 4C, 8C, and 16C. This suggests either that the endoreplication occurs slowly and continuously without substantial gaps between endocycles, or that the endoreplication only covers part of the genome. An



**Figure 2** Examples of invariant postembryonic cell lineages in *C. elegans*. (A) Lineage of the stem-cell like epithelial seam cells. The *y*-axis indicates the time (hours) of development from hatching; vertical lines represent seam cells, horizontal lines are cell divisions, and an H denotes the *hyp7* fusion fate; anterior is to the left. S indicates DNA synthesis and G1 temporal quiescence. Note the repeated rounds of asymmetric cell divisions, separated by a single proliferative division at the L1/L2 boundary. (B) Immunofluorescence microscopy image of a transgenic animal expressing cell membrane (GFP::PHPLC<sup>18</sup>) and DNA (GFP::H2B) markers in the seam cell lineage (controlled by the *wrt-2* promoter). The animal just completed asymmetric seam cell divisions in the second larval stage (L2). Indicated is a seam cell that just divided, of which the anterior daughter (for simplicity named Vn.a) will fuse with the general epidermis (*hyp7*), while the posterior cell Vn.p will remain present as a seam cell. Also marked is the nucleus of the *hyp7* syncytium. (figure courtesy of S. van der Horst). (C) Cell lineages for the vulval precursor cells that give rise to neurons and glial cells in the L1 larval stage, and the vulva in the L3 stage; and the postembryonic mesoderm lineage that gives rise to body wall muscles, coelomocytes, and sex myoblasts in the L1 stage, and vulval and uterine muscles in the L3 stage.

analysis of copy numbers between genomic regions could distinguish these possibilities, but has not been reported. The adult endoreplication of *hyp7* nuclei is positively regulated by TGF- $\beta$  signaling (Nystrom *et al.* 2002). The extent of endoreplication of the adult *hyp7* nuclei depends on the level of food intake, with diet-restricted animals exhibiting lower levels of endoreplication (Tain *et al.* 2008).

For both the hypodermis and intestine, a possible explanation for why the cells evolved to undergo endoreplication is to preserve the structural integrity of those tissues. Endoreplication provides increased genomic DNA content to allow cell growth without the disorganization of tissue integrity that could occur if the cells underwent mitosis (Hall and Altun 2008).

### Germ cell cycles

The germline contains ~2000 germ cells in the adult hermaphrodite, approximately twice as many cells as are present in the adult soma [reviewed in Hansen and Schedl (2013) and Kimble and Seidel (2013)]. The germline is generated from two primordial germ cells in the newly hatched L1 larvae, Z2 and Z3 (Kimble and Hirsh 1979). The terminology “germ cells” is typically used in the *C. elegans* field to refer to all cells derived from the Z2 and Z3 lineage, encompassing all

four larval stages and adults. During the larval stages, germ cells divide continuously to increase from 2 to ~2000 cells in early-stage adults (Kimble and Crittenden 2005). The rate of proliferation of germ stem cells (GSCs) is higher in larvae than in the adult (Roy *et al.* 2016). During the L3 and L4 larval stages, a subset of GSCs enter meiosis to produce sperm, and then in the adult stage, GSCs enter meiosis to produce oocytes. Germ cells in adults and later larval stages are syncytial with an opening in their plasma membrane that allows a connection to a common cytoplasm.

Adult GSCs are the only cells in *C. elegans* that closely match the description of an adult stem cell population that divides within a stem cell niche. The stem cell niche is located at the distal end of each gonad arm in hermaphrodites. The niche is formed by the somatic gonadal distal tip cell (DTC) that encases the distal end of the gonad, with cellular projections that extend over the region in which mitotic germ cells are present (~20 germ cell diameters); germ cells past this point are nonmitotic and have entered meiosis. Under well-fed conditions, adult-stage GSCs divide continuously within the stem cell niche, and upon leaving the niche, they enter meiosis to produce oocytes. As discussed below, the germ cell cycle differs from somatic larval cell cycles in that the G1 phase is truncated or nonexistent in the majority of germ

cells. Additionally, germ cells differ from somatic larval-stage cells in their regulation of DNA replication. These observations highlight that basic aspects of cell cycle control have evolved differently in germ cells and somatic tissues.

## The Regulation of Cell Cycle Progression During Development

As described above, the pattern and timing of cell divisions in *C. elegans* depend on the developmental context and lineage of the cells. Cells form different lineages starting as early as the first division of the zygote (P0), when cellular components become unequally distributed among an anterior somatic blastomere (AB) and posterior germline precursor (P1). This division has been studied extensively as a model for asymmetric cell division, which create cell diversity, and often segregate the potential to proliferate and differentiate unevenly among the daughter cells. In addition to asymmetric cell division, cell–cell signaling contributes to the lineage-specific division patterns, starting as early as the four-cell stage. Moreover, environmental conditions such as the levels of nutrients and oxygen induce systemic responses that determine developmental progress. The interplay between developmental cues, environmental signals, and cell cycle regulators ultimately determines whether cells initiate or arrest cell division. In this section, we discuss how developmental and environmental processes may be integrated with the cell cycle machinery.

### Oocyte meiotic arrest and maturation

Development starts with a fertilized egg. For fertilization to be successful, oocytes first need to grow, mature, ovulate, and progress through meiosis I. As in most other animals, developing *C. elegans* oocytes arrest cell cycle progression in meiotic prophase I while growth continues (Figure 3). This arrest depends on low CDK-1/cyclin B activity, which is assumed to result from inhibitory phosphorylation of CDK-1 by the WEE-1.3 dual-specificity kinase (Detwiler *et al.* 2001; Burrows *et al.* 2006). The onset of maturation releases the meiotic arrest and allows oocytes to progress from diakinesis to meiotic metaphase, accompanied by nuclear envelope breakdown, meiotic spindle assembly, and cortical rearrangements [reviewed in Von Stetina and Orr-Weaver (2011) and Kim *et al.* (2013)]. These changes suggest that CDK-1 becomes activated during maturation, and indeed, oocyte maturation in *C. elegans* requires CDK-1 and mitotic cyclins (Boxem *et al.* 1999; Burrows *et al.* 2006; van der Voet *et al.* 2009), as well as the polo-like kinase PLK-1 (Chase *et al.* 2000). In this way, *C. elegans* oocyte maturation resembles that in other animals. In fact, studies of this process in frogs provided a breakthrough in understanding cell cycle regulation, when purified maturation-promoting factor (MPF) was found to consist of CDK1 in association with a mitotic cyclin (Masui and Markert 1971; Dunphy *et al.* 1988; Gautier *et al.* 1988).

What came as a surprise was the finding that *C. elegans* uses the sperm-specific cytoskeletal protein MSP (major sperm


protein) as a hormonal trigger for maturation (Miller *et al.* 2001) (Figure 3A). MSPs are released from sperm in the adjacent spermathecae, and bind the Ephrin-related receptor VAB-1 on oocytes and somatic sheet cells of the gonad. This interaction overcomes VAB-1 Eph-mediated inhibition of a mitogen-activated protein kinase (MAPK) pathway in oocytes (Miller *et al.* 2003). In addition to MPF, meiotic progression and maturation depend on MAPK activation in *C. elegans* as well as vertebrate oocytes (Kim *et al.* 2013). Thus, in common with other animals, a hormonal signal overcomes the prophase I meiotic arrest of *C. elegans* oocytes by activating MAPK and MPF. However, the nature of the hormonal signal varies among animals between progesterone in frogs, luteinizing hormone (LH) in humans, and MSP in *C. elegans*.

Another critical step in meiotic maturation is the translational activation of maternally inherited mRNAs. In frogs, this includes the synthesis of cyclin B from mRNAs that are kept dormant in prophase-arrested oocytes. Regulators of mRNA translation are downstream targets of MSP signaling in *C. elegans* meiotic maturation. One of these regulators is the conserved RNA-binding protein LIN-41 TRIM-NHM (Spike *et al.* 2014a,b), originally discovered as a “heterochronic” gene that provides temporal control over larval development (Slack *et al.* 2000). LIN-41 acts as a translational repressor in the germline, both in cooperation with and antagonized by the “oocyte maturation-defective” proteins OMA-1 and OMA-2 (Tsukamoto *et al.* 2017) (Figure 3B). The redundant OMA-1/2 zinc finger proteins regulate multiple processes, either through binding the 3'-UTRs of specific mRNAs or protein association. Genetic and protein expression data indicate complex interactions between LIN-41 and OMA-1/2. LIN-41 is expressed throughout meiotic prophase, promotes meiotic arrest, and prevents CDK-1 activation. In contrast, OMA-1/2 is not expressed until after the pachytene stage, antagonizes LIN-41, and promotes meiotic maturation and CDK1 activation (Detwiler *et al.* 2001; Spike *et al.* 2014a,b; Tsukamoto *et al.* 2017). Despite their opposite functions, both LIN-41 and OMA-1/2 ribonucleoprotein (RNP) complexes associate with and repress *cdc-25.3* mRNA. This probably explains the contribution of LIN-41 in meiotic arrest; however, as *cdc-25.3* remains translationally repressed until early embryogenesis, derepression of *cdc-25.3* mRNA translation does not trigger meiotic maturation in the wild type. Interestingly, some other mRNAs (*spn-4*, *meg-1*) are repressed by LIN-41 while being activated by OMA-1/2 (Tsukamoto *et al.* 2017). The translational activation of these mRNAs and the release of translation regulatory proteins from LIN-41 RNPs probably supports growth and meiotic maturation.

### Completion of meiosis

As in other animals, oocyte maturation in *C. elegans* triggers further progression through meiosis I. RNAi experiments demonstrated that not only CDK-1 but also individual B-type cyclins exert critical roles in meiotic progression (Boxem *et al.* 1999; Chase *et al.* 2000; van der Voet *et al.*





**Figure 3** Meiotic arrest, oocyte maturation, fertilization, and the oocyte-to-embryo transition in the hermaphrodite germline. (A) Schematic illustration of the successive stages from meiosis I-arrested oocytes in the gonad arm (left) to mitotic embryos in the uterus (right). Nuclei are green, sperm are orange. (B) Diagram of various regulators that mediate the cell cycle transitions in time, from left to right. Initially, WEE-1.3 and LIN-41 are thought to antagonize CDK-1 activation. See text for further information.

2009; Deyter *et al.* 2010). *C. elegans* expresses three typical B-type cyclins (CYB-1, CYB-2.1, and CYB-2.2) and an evolutionarily conserved B3-type cyclin (CYB-3) (Kreutzer *et al.* 1995; van der Voet *et al.* 2009). B3 cyclins form a distinct subfamily of cyclins that share sequence motifs with both A- and B-type cyclins. Only simultaneous inhibition of all four B-type cyclins fully resembles *cdk-1* loss and causes arrest in diakinesis of meiosis I (van der Voet *et al.* 2009). Nevertheless, knockdown of individual cyclins leads to distinct defects, which suggests that the B-type cyclins have overlapping as well as specific functions in substrate phosphorylation. The RNAi data indicate that CYB-1 is required specifically for the full condensation and alignment of chromosomes at the metaphase plate in meiosis, as well as mitosis. In contrast, CYB-3 is required for sister chromatid separation at the metaphase-to-anaphase transition in meiosis and mitosis (Cowan and Hyman 2006; van der Voet *et al.* 2009).

The data appear to fit with recent observations in *Drosophila* embryos and mouse oocytes, which support the idea that cyclin B3 stimulates APC/C activation and securin degradation (Yuan and O'Farrell 2015; Zhang *et al.* 2015). However, the mechanism by which cyclin B3 promotes APC/C activation may differ between these species. In contrast to flies and mice, *C. elegans* *cyb-3(RNAi)* embryos show increased kinetochore localization of the spindle assembly checkpoint (SAC) component MAD2 (Deyter *et al.* 2010). Consistently, the metaphase arrest of *cyb-3(RNAi)* embryos could be suppressed by co-inhibition of SAC components (Deyter *et al.* 2010). Thus, the absence of CYB-3 appears to trigger a strong SAC-mediated block of APC/C activation and arrest in metaphase. By contrast, cyclin B3 activation of the APC/C in fly embryos and mouse oocytes is SAC-independent (Yuan and O'Farrell 2015; Zhang *et al.* 2015).

While CDK-1/B-type cyclin complexes promote progression from prophase to metaphase of the meiotic divisions (van der Voet *et al.* 2009), the transition from metaphase to anaphase is triggered by the APC/C-FZY-1<sup>CDC20</sup> (Golden *et al.* 2000; Davis *et al.* 2002). In agreement, forward genetics and genome-wide RNAi screens found that not only *cdk-1* and *cdc-25.1* are required for passage through meiosis, but also multiple *mat* (metaphase-to-anaphase transition defective) genes that encode APC/C components, *fzy-1<sup>CDC20</sup>*, and other ubiquitin-dependent proteolysis factors that include proteasome subunits (Golden *et al.* 2000; Davis *et al.* 2002; Sönnichsen *et al.* 2005).

An interesting question is how CDK and APC/C activities are coordinated with meiotic progression and fertilization. Interestingly, the CRL2-ZYG-11 ubiquitin ligase, containing the ZYG-11 substrate-specificity factor, is required for progression through meiosis II and cyclin B1/B3 degradation in meiosis (Liu *et al.* 2004; Sonnevile and Gönczy 2004; Vasudevan *et al.* 2007). CRL2-ZYG-11 acts redundantly with the APC/C in degrading cyclin B1 and B3 (Figure 3B). CRL2-ZYG-11 targets at the least CYB-1 directly for degradation in *C. elegans*, a function recently found to be conserved in human cells (Balachandran *et al.* 2016). The human and *C. elegans* CRL2-ZYG-11A/B complexes are not required for normal progression through mitosis. However, when the APC/C is kept inactive in response to SAC activation in human cells, CRL2-ZYG-11A/B-mediated degradation of cyclin B1 allows exit from mitosis, which explains a phenomenon known as “mitotic slippage” (Balachandran *et al.* 2016).

Similar to *zyg-11* or *cul-2* mutants, unfertilized oocytes fail to go through anaphase II and retain higher levels of CYB-1 than fertilized eggs (McNally and McNally 2005). It is tempting to speculate that fertilization normally triggers

the activation of the CRL-2-ZYG-11 E3 ligase in meiosis II, which complements APC/C-FZY-1<sup>CD<sub>20</sub></sup> in mitotic cyclin destruction. Only the combined E3 ligases would achieve the complete downregulation of CDK/cyclin activity that is needed for “licensing” of DNA replication origins during the transition from meiosis to mitotic cell division. The extensive cyclin protein degradation in meiosis and, in particular, early mitoses makes it likely that the resynthesis of cyclin B is needed for subsequent embryonic divisions.

In normal development, fertilization occurs immediately after the oocyte enters the spermatheca. Notably, when fertilization does not take place, DNA synthesis eventually reinitiates. This has been observed in multiple conditions. For instance, hermaphrodites that are exhausted of sperm at the end of the brood continue to lay mature oocytes that resume rounds of DNA replication and nuclear envelope breakdown, which is described as an endomitotic oocyte (*Emo*) phenotype (Ward and Carrel 1979). A similar *Emo* phenotype results from interfering with ovulation by ablating a proximal gonadal sheet cell (McCarter *et al.* 1997), and various sperm-defective (*spe*) or fertilization-defective (*fer*) mutants produce unfertilized oocytes with a single polyploid nucleus (L'Hernault *et al.* 1988). Oocytes in such mutants undergo maturation and ovulation, and initiate anaphase I (McNally and McNally 2005). However, the chromosomes do not become segregated into polar bodies, and meiosis II chromosome congression, spindle formation, and anaphase do not occur. Nevertheless, a female pronucleus forms without apparent delay. These data demonstrate that *C. elegans* oogenesis does not include a second meiotic arrest, in contrast to most other animals, in which mature oocytes remain arrested in metaphase I (*Drosophila*) or meiosis II (vertebrates) until fertilization occurs (Von Stetina and Orr-Weaver 2011).

Fertilization coincides with egg activation and induces the completion of both meiotic divisions, the switch to mitotic divisions, and the formation of a protective egg shell. The mechanisms by which fertilization triggers egg activation are poorly understood, but certain *spe/fer* mutants are likely to miss critical steps. Specifically, the sperm-derived factor *SPE-11* is essential for the completion of meiosis and polar body formation (McNally and McNally 2005). However, *SPE-11* does not appear to be evolutionarily conserved and its molecular function currently remains unknown. In vertebrates, release of metaphase II arrest results from activation of the APC/C in response to a transient intracellular Ca<sup>2+</sup> increase that follows sperm entry (Von Stetina and Orr-Weaver 2011). Ca<sup>2+</sup> oscillations have also been described during meiotic resumption of fertilized *C. elegans* eggs (Singaravelu and Singson 2013). Several *spe/fer* mutants affect Ca<sup>2+</sup> channels in the sperm, and the sperm-provided TRP-3 (*SPE-41*) Ca<sup>2+</sup> channel is not only required for fertilization but also contributes to a local Ca<sup>2+</sup> wave upon fertilization (Singaravelu *et al.* 2012). It remains to be determined whether this is critical for the fertilization signal and is linked to the cell cycle.

### **From meiosis to mitosis during the “oocyte-to-embryo” switch**

Following meiotic progression, CDK-1 and APC/C also exert functions specific for the oocyte-to-embryo switch. This transition requires that a number of proteins present in oocytes are degraded or modified. Critical in this process is *MBK-2*, a member of the family of dual-specificity tyrosine phosphorylation-regulated kinases (DYRKs) (Pellettieri *et al.* 2003; Quintin *et al.* 2003; Stitzel *et al.* 2006). DYRKs become tyrosine autophosphorylated during their synthesis, which is critical for kinase activity. However, *C. elegans* *MBK-2* becomes active specifically during egg activation, in part through direct phosphorylation by CDK-1 (Cheng *et al.* 2009) (Figure 3B). Three pseudophosphatases that are required for egg activation, *EGG-3* and the closely related *EGG-4/EGG-5*, keep *MBK-2* tethered at the cortex of oocytes in an inactive form (Maruyama *et al.* 2007; Stitzel *et al.* 2007; Cheng *et al.* 2009; Parry *et al.* 2009). Activation of the APC/C in meiotic metaphase induces the release of active *MBK-2* through degradation of *EGG-3* and probably *EGG-4/5*. Active *MBK-2* phosphorylates the katanin-related protein *MEI-1*, which targets *MEI-1* for ubiquitin-dependent degradation (Quintin *et al.* 2003). *MEI-1* promotes the formation of compact meiotic spindles and its degradation is needed to allow the formation of extensive mitotic spindles. Further, *MBK-2* phosphorylates the *OMA-1/2* proteins, which induces their degradation in the somatic precursors (Shirayama *et al.* 2006; Stitzel *et al.* 2006) and redirects the function of *OMA-1/2* to inhibit transcription in early germline blastomeres (Güven-Ozkan *et al.* 2008). Additionally, *MBK-2* phosphorylates *MEX-5* and *MEX-6* to create a contact site for the polo-like kinase *PLK-1*, which is critical in the establishment of anterior–posterior polarity of cytoplasmic components (Nishi *et al.* 2008). Thus, parallel regulation of *MBK-2* by CDK-1 and APC/C couples the oocyte-to-embryo transition to meiotic progression (Figure 3B).

### **Embryonic cell cycles**

Cell cycle progression in the early embryo is likely driven by a single CDK (*CDK-1*) in association with B-type cyclins. In support of this, the alternating S and M phases of the early cleavage-type embryonic cell divisions do not require *CDK-4*/cyclin D or *CDK-2*/cyclin E kinase activity (Boxem *et al.* 1999; Boxem and van den Heuvel 2001). Following *cye-1* knockdown, cell division continues until approximately the 100-cell stage (Fay and Han 2000). Remarkably, however, *CDK-2* and *CYE-1* are essential for anterior–posterior polarity in the one-cell egg (Cowan and Hyman 2006). Studies of this phenotype revealed that the *CDK-2/CYE-1* kinase promotes assembly of centrosomal proteins, while the mature centrosome provides a symmetry-breaking cue during polarity establishment.

As *cdk-1* knockdown results in meiotic arrest, insight into its mitotic contributions has been obtained mostly by following the inactivation of mitotic cyclins (Boxem *et al.* 1999;


Chase *et al.* 2000; van der Voet *et al.* 2009; Deyter *et al.* 2010). The depletion of individual cyclins results in very similar defects in meiosis and mitosis. As described above (“completion of meiosis”), the observed phenotypes support the notion that the four B-type cyclins act partly redundantly, while *CYB-1* is also specifically required for chromosome condensation and congression, and *CYB-3* is essential for anaphase onset. Moreover, immunostaining of zygotes depleted of *cyb-3* showed severely delayed and reduced localization of PCN-1<sup>PCNA</sup> in paternal pronuclei, probably reflecting delayed S phase progression (Michael 2016). This indicates that *CYB-3* also contributes to S phase control. As such, *CYB-3* could potentially complement or functionally replace cyclin E and cyclin A, which would explain why these latter cyclins are not required for DNA synthesis in early embryos (van der Voet *et al.* 2009).

### Timing asynchrony

The two daughter cells formed during the first mitotic division display asynchrony in the timing of mitotic entry, in stark contrast to the synchronous nuclear divisions in *Drosophila*, or cell cleavages during *Xenopus* and marine invertebrate early embryogenesis. The larger anterior blastomere AB enters mitosis and divides ~2 min before the posterior blastomere P1. One mechanism underlying this asynchrony involves a checkpoint pathway that responds to DNA replication stress (Brauchle *et al.* 2003) (Figure 4). This pathway uses the ATR/ATM-related kinase *ATL-1* and its downstream effector, the *CHK-1* kinase. Inactivating this checkpoint in wild-type embryos reduced the difference in division timing between AB and P1 by ~40% (Brauchle *et al.* 2003). These data suggest that a DNA replication checkpoint is normally engaged in P1 to achieve the proper timing of cell division, and possibly, through lengthening S phase, promoting the fidelity of DNA replication in the germline precursor cells.

Also contributing to timing asynchrony is the asymmetric distribution of the polo-like kinases *PLK-1* and *PLK-2* (Budirahardja and Gönczy 2008; Nishi *et al.* 2008; Rivers *et al.* 2008), and cyclin B3 (Michael 2016) (Figure 4). These proteins become enriched in the AB blastomere in response to anterior–posterior polarity, established through asymmetric cortical localization of PAR (partitioning-defective) proteins. The polo kinases accumulate in the anterior of the one-cell embryo by binding to the *MEX-5/MEX-6* (“muscle excess”) proteins (Nishi *et al.* 2008). *MEX-5/6* are regulators of cytoplasmic asymmetry that accumulate themselves in the anterior in response to phosphorylation by the *PAR-1* kinase (Tenlen *et al.* 2008; Griffin *et al.* 2011). Additional phosphorylation by *MBK-2* primes *MEX-5* for interaction with and phosphorylation by *PLK-1* (see above; Nishi *et al.* 2008). In this way, the *MEX-5/6*–*PLK-1* interaction not only contributes to *PLK-1* localization, but also promotes *MEX-5/6* function in regulating the asymmetric distribution of cytoplasmic determinants in the one-cell embryo.

*CDK-1* appears to act both as an upstream activator and downstream target in the *PLK-1* cell cycle pathway, through



**Figure 4** Cell division timing asynchrony in the two-cell embryo. (Top) Combined fluorescence and DIC microscopy image of a two-cell-stage embryo in which the DNA is visualized by GFP::H2B expression. The anterior blastomere (AB) initiated mitosis almost 2-min earlier than the precursor of the germline P1. (Bottom) Several cell cycle-regulatory mechanisms have been found to underlie the timing asynchrony. One is based on the anterior enrichment of the *MEX-5/6* cytoplasmic determinants, which interact with the *PLK-1,2* Polo kinases. The other involves preferential activation of the *ATL-1*<sup>ATR</sup>–*CHK-1* DNA replication checkpoint pathway in P1. See text for further information.

mechanisms that are conserved in human cells [reviewed in Zitouni *et al.* (2014)]. As an activator, *CDK-1* phosphorylates the *PLK-1* regulator *SPAT-1*<sup>Bora</sup> (Tavernier *et al.* 2015). This promotes *SPAT-1* interaction with *PLK-1* and exposes the *PLK-1* T loop to activating phosphorylation by Aurora A (*AIR-1*) (Figure 4). Active *PLK-1* activates *CDK-1* as a downstream target by promoting nuclear accumulation of the *CDK-1*-activating phosphatase *CDC-25.1* (Rivers *et al.* 2008). Thus, the asymmetric localization of *PLK-1* leads to asymmetry in nuclear *CDC-25.1*, which is expected to contribute to advanced mitotic entry of the anterior blastomere via more rapid activation of *CDK-1/cyclin B* (Figure 4). Indeed, knockdown of the *wee-1.3* Myt1 kinase, which mediates inhibitory phosphorylation of *CDK-1*, advances the cell division timing, and reduces the asynchrony between AB and

P1 (Michael 2016). This confirms that the negative phosphorylation of CDK-1 is a critical timing event in the early embryo.

### Checkpoint-induced cell cycle arrest in the early embryo

In yeast, checkpoint control pathways are essential only after damage or errors occur. Yet, as described above, the *C. elegans* ATL-1/CHK-1 checkpoint pathway increases cell cycle length during normal embryonic development. This is reminiscent of late syncytial divisions in *Drosophila* embryos, in which the cell cycle lengthens during the switch to zygotic gene expression. This midblastula transition is induced by a developmentally regulated DNA replication/damage checkpoint pathway that involves the *Drosophila* mei-41 (ATM) and grapes (Chk1) kinases [reviewed in Yuan *et al.* (2016)].

Despite a constitutive role for checkpoint activation in P1, DNA damage and replication stress still induce a substantial delay in cell cycle progression of early *C. elegans* blastomeres. Interrupting DNA synthesis or UV irradiation generates a cell cycle delay of up to 15 min in the *C. elegans* zygote, with subsequent divisions showing shorter delays (Encalada *et al.* 2000; Brauchle *et al.* 2003; Stevens *et al.* 2016). Notably, the DNA replication checkpoint delays are substantially longer in the germline precursor cells (P0, P1, and P2) than the somatic sister cells. This increased response to DNA replication stress in germline precursors probably evolved to protect the genome of future generations.

Checkpoint genes are also used to respond to environmental conditions. For instance, a pathway related to the SAC induces an extreme form of developmental quiescence in response to severe oxygen deprivation (anoxia) (Nystul *et al.* 2003). This so-called “suspended animation” coincides with a cell cycle arrest of early blastomeres. A substantial percentage of the cells arrest in mitotic metaphase, dependent on the *san-1*<sup>MAD3/BubR1</sup> and *MDF-2*<sup>MAD2</sup> SAC pathway genes. However, anoxia also induces arrest in other phases of the cell cycle, including a prophase arrest that depends on the nucleoporin *NPP-16*<sup>NUP50</sup>. These arrests are critical for embryo survival and development to adulthood following reoxygenation.

Exposing adult animals to low oxygen (hypoxia) induces an “embryonic diapause” that may be a natural form of suspended animation to protect embryos *in utero* (Miller and Roth 2009). This response not only requires *san-1*, probably in the embryo, but also the hypoxia-inducible factor *HIF-1* in neurons of the mother. Mitochondrial functions and reduced ATP levels are likely involved in these responses. One-cell embryos arrest prior to nuclear envelope breakdown in response to RNAi of genes encoding tricarboxylic acid cycle/Krebs cycle components (Rahman *et al.* 2014). These embryos arrest with CDK-1 predominantly in the inhibitory phosphorylated form. The observations point to a possible connection between the levels of ATP, or other metabolites, and cell cycle regulation via the CDC-25 phosphatase.

The SAC also shows an interesting cell size dependence. The SAC delays APC/C activation when the kinetochores of

sister chromatids are not properly attached to microtubules of opposite spindle poles. Whereas early embryos of flies and frogs appear to lack this checkpoint, early *C. elegans* embryos display a moderate delay in anaphase onset in response to spindle defects (Encalada *et al.* 2005). This delay depends on conserved checkpoint proteins, including *MDF-1*<sup>MAD1</sup>, *MDF-2*<sup>MAD2</sup>, and *SAN-1*<sup>MAD3</sup>. Notably, *C. elegans* misses an MPS1 SAC kinase homolog. However, a recent study demonstrated that *PLK-1*<sup>Polo</sup> functionally replaces MPS1 in *C. elegans* and that polo kinase also participates in this checkpoint in mammals (Espeut *et al.* 2015; von Schubert *et al.* 2015). Germline precursors show a stronger SAC response than their somatic sisters, and more strikingly, the SAC-induced metaphase delay increases with each subsequent embryonic cell division (Galli and Morgan 2016). This phenomenon correlates strongly with the kinetochore-to-cytoplasm ratio, and has been explained by the larger fraction of APC/C that can be inhibited by checkpoint proteins in smaller cells. Free kinetochores generate the checkpoint signal through interaction with MAD2–CDC20. Hence, the fact that *C. elegans* has holocentric chromosomes and its egg has a relatively small volume may explain why *C. elegans* embryos display a SAC response, in contrast to the much larger embryos of frogs and flies (Galli and Morgan 2016).

### Lineage-dependent introduction of GAP phases and quiescence

As described in the “Developmental and Tissue Cell Cycle Variants” section, introduction of GAP phases occurs in a lineage-dependent fashion. The first G2 phase appears at the onset of gastrulation at the 26-cell stage. The intestinal precursor cells Ea and Ep complete DNA synthesis before they migrate inward to initiate gastrulation. Mitosis is delayed until after the migration, corresponding to a G2 phase of ~1 hr (Edgar and McGhee 1988). The precursor cell for the germline follows this inward movement of Ea/Ep, and subsequently divides one more time to form the Z2 and Z3 primordial germ cells. These germline precursor cells are the first cells to undergo a prolonged cell cycle arrest, which lasts from 140 min of embryonic development until midway through the first larval stage, a period of ~18 hr (Sulston *et al.* 1983). In most embryonic lineages, the cell cycle gradually lengthens, and nearly all cells arrest division between 4 and 7 hr of embryogenesis, to either become quiescent or initiate terminal differentiation.

Several observations indicate that the extended embryonic cell cycles include a G1 or G0 state. First, a few embryonic cells depend on G1 regulators to reinitiate cell division after an extended interphase. This includes the final embryonic divisions of precursor cells in the intestinal, Q neuroblast, and coelomocyte lineages that depend on zygotic *cyd-1* cyclin D expression (Boxem and van den Heuvel 2001; Yanowitz and Fire 2005; S. van den Heuvel, unpublished data). Thus, at least these specific embryonic cell cycles include a G1 phase. The blast cells that form the postembryonic lineages also appear to enter a G0 or extended G1 state, as their division

during larval development requires cyclin D and CDK-4 (Park and Krause 1999; Boxem and van den Heuvel 2001). Moreover, in the absence of maternal supplies of the G1/S regulators *cki-1*<sup>CIP/KIP</sup>, *cul-1*<sup>cullin</sup>, or *lin-23*<sup>B-TrCP</sup>, embryos die with many additional cells (over 850 instead of 558) (Kipreos *et al.* 1996, 2000; Fukuyama *et al.* 2003). Similar to *Drosophila* dacapo, *cki-1* loss results in a single extra division at the time of normal arrest, as observed in the intestinal and mesodermal lineages (Fukuyama *et al.* 2003). In contrast, the primordial germ cells normally arrest in G2/M with condensed chromosomes, and this arrest is not affected by the absence of *cki-1*<sup>CIP/KIP</sup> (Fukuyama *et al.* 2003, 2006). In short, in a few hours of embryonic development, cells change in a lineage-specific pattern from undergoing rapidly alternating S and M phases to an irreversibly arrested postmitotic state (differentiated cells), or a temporarily arrested state of quiescence (G0 or extended G1) or G2/M arrest (primordial germ cells).

### **Systemic regulation of cell division during larval development**

The timing of the larval cell divisions depends not only on lineage-specific information, but also on environmental and systemic developmental signals. Unfavorable conditions soon after hatching trigger a developmental arrest, while such conditions experienced later in life can induce autophagy in the germline and cell cycle arrest of GSCs. These conditions require synchronization between cell cycle progression and development.

### **Cell cycle arrest of newly hatched larvae**

The first stage larva hatches from its egg in a state without growth or active cell division. In the absence of food or nutrients, the animal remains in this arrested state, also known as L1 diapause, and can survive up to several weeks [Castro *et al.* 2012; reviewed in Baugh (2013)]. Uptake of food immediately triggers the larval developmental program, which includes growth and cell division. Insulin/insulin-like growth factor (IGF) signaling and several other factors have been found to control the switch between L1 arrest and development. IGF signaling has been extensively studied in *C. elegans* because inactivation of specific components of this pathway increases life span and allows entry into the alternative “dauer” larval state. These studies identified a single IGF receptor, DAF-2<sup>ILR</sup>, which activates the phosphoinositide 3-kinase AGE-1<sup>PI3K</sup> and downstream kinases AKT-1/2<sup>Akt/PKB</sup>, while the lipid phosphatase DAF-18<sup>PTEN</sup> acts as an antagonist [reviewed in Murphy and Hu (2013)]. A large variety of insulin-related peptides can either activate or antagonize DAF-2<sup>ILR</sup> receptor signaling. Activation of the pathway leads, through activation of AKT-1/2, to phosphorylation of the DAF-16<sup>FoxO</sup> transcription factor. This phosphorylation interferes with nuclear import of DAF-16, and thereby leads to DAF-16 inactivation. IGF signaling through DAF-16 and other downstream effectors is not only critical for the regulation of life span and dauer arrest, but also for cell cycle control during L1 development.

The initiation of L1 development in the presence of food depends on the DAF-2<sup>ILR</sup> receptor and the expression of a subset of insulin-related peptides (Baugh and Sternberg 2006; Baugh 2013). Conversely, both DAF-16<sup>FoxO</sup> and DAF-18<sup>PTEN</sup> are needed for L1 arrest in the absence of food. Arrested L1 larvae express *cki-1*<sup>CIP/KIP</sup> in somatic blast cells, and this expression is lost in *daf-16* mutants (Hong *et al.* 1998). *daf-16* mutants fail to properly arrest cell division and show poor survival in the absence of food. Interestingly, the Z2 and Z3 primordial germ cells undergo multiple rounds of division in *daf-18* mutants (which have increased insulin signaling), independently of *daf-16* and *cki-1* (Fukuyama *et al.* 2006). This divergence from somatic cells likely reflects the unique arrest of Z2/Z3 at the G2/M transition (see above). Loss of both of the *Drosophila* nanos-related genes *nos-1* and *nos-2* also leads to Z2/Z3 divisions under starvation conditions (Subramaniam and Seydoux 1999). It is currently not known whether *nos-1/nos-2* act as downstream targets of IGF-signaling to regulate the cell cycle arrest of germline precursor cells.

Several additional regulators have been found to affect L1 diapause arrest, at least in part as contributors to IGF signaling. This includes a conserved ATPase ASNA-1 and micro RNA (miRNA) *miR-73*, which are probably involved in the secretion of specific insulin-related peptides from sensory neurons and the intestine (Baugh 2013). The expression of another miRNA, *miR-235*, also contributes to L1 diapause arrest. Feeding leads to *miR-235* downregulation through IGF signaling. This induces expression of the *nhr-91* nuclear hormone receptor (germ cell nuclear factor), a *miR-235* target that promotes L1 development (Kasuga *et al.* 2013). While IGF signaling and *miR-235* affect L1 development in general, the DAF-16-mediated regulation of *cki-1*<sup>CIP/KIP</sup> is currently the only established link to the cell cycle. This connection is particularly relevant because mammalian FoxO transcription factors also have been reported to induce cell cycle arrest through p27<sup>KIP1</sup> regulation, under the control of PI(3)K–Akt/PKB signaling (Medema *et al.* 2000).

### **Cell cycle entry during larval development**

While *cki-1*<sup>CIP/KIP</sup> is critical for cell cycle arrest, activation of the CDK-4/cyclin D kinase likely drives the resumption of cell division during larval development. In support, precursor cells of the somatic cell lineages show transcriptional activation of the *cyd-1*<sup>Cyclin D</sup> and *cdk-4*<sup>CDK4/6</sup> genes coincident with cell cycle entry (Park and Krause 1999; Brodigan *et al.* 2003). Moreover, these blast cells remain arrested in G0/G1 through all larval stages in *cyd-1* and *cdk-4* mutants (Park and Krause 1999; Boxem and van den Heuvel 2001). Further, ectopic expression of G1 cyclins or CDK/cyclin combinations in arrested larvae induces DNA replication and cell division (Park and Krause 1999; Korzelius *et al.* 2011b). Thus, although the underlying regulation currently remains unknown, the timing of *cyd-1* and *cdk-4* transcription and kinase activation appears to control cell cycle entry during larval development.

To induce the G1–S phase transition, the CDK-4/Cyclin D kinase needs to overcome inhibition of cell cycle entry by *lin-35<sup>Rb</sup>*, *cki-1<sup>CIP/KIP</sup>*, and *fzr-1<sup>Cdh1</sup>* (Boxem and van den Heuvel 2001; The *et al.* 2015). These negative regulators show substantial functional redundancy among each other and also with the SCF–LIN-23 <sup>$\beta$ -TrCP</sup> E3 ubiquitin ligase. Indeed, single mutation of the *C. elegans* Rb-related gene *lin-35* or the APC/C coactivator *fzr-1<sup>FZR1/Cdh1</sup>* barely affects cell division. However, combining these mutations results in substantial overproliferation, and either single mutation increases the hyperplasia associated with *cki-1* or *cul-1/lin-23* loss (Boxem and van den Heuvel 2001; Fay *et al.* 2002; Ruijtenberg and van den Heuvel 2015; The *et al.* 2015). While these genes all have general functions in G1/S inhibition, their relative contribution is substantially lineage-dependent. Although possibly more pronounced in *C. elegans*, remarkable redundancies among different regulators of the G1/S transition have also been observed in flies (Buttitta *et al.* 2007) and mice (Wirt *et al.* 2010).

The extra cell divisions in *cul-1<sup>Cullin</sup>* and *lin-23 <sup>$\beta$ -TrCP</sup>* mutants start during the second larval stage, at least in part because maternal product suffices at earlier developmental stages (Fukuyama *et al.* 2003). The overproliferation phenotype indicates that the SCF–LIN-23 <sup>$\beta$ -TrCP</sup> E3 ligase promotes the degradation of positive cell cycle regulators. Indeed, strong genetic evidence points to the CDC-25.1 and CDC-25.2 phosphatases as critical *in vivo* targets (Hebeisen and Roy 2008; Segref *et al.* 2010; Son *et al.* 2016). Substrate recognition by  $\beta$ -TrCP often involves interaction with a phosphorylated recognition site known as a phosphodegron. Gain-of-function mutations of CDC-25.1 and CDC-25.2 that induce extra intestinal divisions during embryogenesis appear to disrupt a  $\beta$ -TrCP phosphodegron. Similar to the threonine–tyrosine residues in CDK-1 and CDK-2, CDK-4 contains a conserved tyrosine residue that is probably a WEE-1/CDC-25-regulated inhibitory phosphorylation site. However, it is not known whether increased activity of CDK-4, CDK-2, or CDK-1 drives intestinal overproliferation in *cdc-25* gain-of-function mutants. The fact that loss of *lin-23 <sup>$\beta$ -TrCP</sup>* or *cul-1<sup>Cul1</sup>* results in much more extensive hyperplasia, compared to *cdc-25.1/cdc-25.2* gain-of-function mutants, indicates that the SCF–LIN-23 <sup>$\beta$ -TrCP</sup> E3 ligase has critical substrates in addition to the CDC-25 phosphatase. CYD-1 remains a prime candidate, as this G1 cyclin contains a conserved  $\beta$ -TrCP phosphodegron, and GSK-3 $\beta$  phosphorylation-dependent degradation of human cyclin D1 has been reported (Diehl *et al.* 1998).

Candidate substrates for CDK-4/CYD-1<sup>cyclin D</sup> phosphorylation in the regulation of cell cycle entry have also been identified. One of these targets is the *C. elegans* Rb-related protein LIN-35, in agreement with the well-established Cdk4/6–cyclin D regulation of mammalian Rb family proteins (Leng *et al.* 2002; Rubin 2013). LIN-35<sup>Rb</sup> is a substrate for CDK-4/CYD-1<sup>cyclin D</sup> phosphorylation *in vitro*, and loss of *lin-35<sup>Rb</sup>* alleviates the requirement for *cdk-4* and *cyd-1* *in vivo* (Boxem and van den Heuvel 2001; The *et al.* 2015). Of note,

CDK-4/cyclin D phosphorylates LIN-35 at residues that correspond to the CDK phosphorylation sites of Rb that disrupt E2F binding (The *et al.* 2015). However, most cell divisions still depend on CDK-4/cyclin D even in *lin-35<sup>Rb</sup>* null mutants. This demonstrates that the CDK-4/cyclin D kinase has essential functions in addition to inactivating *lin-35<sup>Rb</sup>*. A genetic suppressor screen identified FZR-1<sup>FZR1/Cdh1</sup> as a second critical CDK-4/cyclin D target. The phosphorylation of the FZR-1 N-terminus by CDK-4/cyclin D resembles inhibitory phosphorylation of mammalian FZR1/Cdh1 (The *et al.* 2015). Substantial additional evidence supports the idea that CDK-4/cyclin D promotes cell cycle entry through the combined inhibition of LIN-35<sup>Rb</sup>-mediated transcriptional repression and APC/C–FZR-1-mediated protein degradation. These mechanisms for G1 regulation may be conserved in mammalian cells (The *et al.* 2015).

### Progression through larval cell divisions

In addition to the CDK-4/cyclin D regulator of cell cycle entry, several additional cell cycle genes are required for the proliferation of postembryonic precursor cells. Homozygous *cdk-1* null mutants complete embryogenesis due to the presence of maternal product, and their somatic blast cells go through S phase but arrest in G2 in the first stage larvae (Boxem *et al.* 1999). Such animals grow substantially, molt, and continue intestinal endoreplication cycles during postembryonic development. Thus, CDK-1 is specifically required for the G2/M transition. The lack of proliferation of precursor cells for the ventral nerve cord creates a typical uncoordinated (Unc) phenotype, while the absence of germline proliferation causes sterility. Several mutants with similar Sterile-Unc (Stu) phenotypes defined additional cell cycle genes, such as *air-2* Aurora B and *zyg-1* PLK4 (O’Connell *et al.* 1998; Woollard and Hodgkin 1999). The molecular genetic analysis of additional *stu* mutants could provide further insight in cell cycle regulation.

### The timing of larval development and cell division

Genetic studies have resulted in a model in which larval stage-determining transcription factors (LIN-14 for L1 and HBL-1 Hunchback-like for L2) are downregulated by specific miRNAs (*lin-4* and micro-RNAs of the *let-7* family) to allow transitioning to the next larval stage. These transcription factors and miRNAs have been defined by “heterochronic” mutations that either cause reiteration or skipping of larval stages. Thus, the specific pattern of cell division is not only determined by the cell lineage, but also by the larval stage. For instance, the vulval precursor cells are formed in the L1 larval stage, and undergo a prolonged G1 arrest before entering S phase and cell division in the L3 stage (Figure 2C). In heterochronic mutants that skip the L1 or L2 stage, the VPCs enter S phase and cell division one larval stage earlier (Euling and Ambros 1996) (Figure 2C). However, it is unclear whether heterochronic transcription factors directly control cell cycle regulators. The *let-7* miRNA is needed for the L4-to-adult transition and indirectly upregulates the heterochronic

transcription factor LIN-29. Cells in the epidermis arrest proliferation and terminally differentiate in late L4. This coincides with high expression of CKI-1<sup>CIP/KIP</sup> and reduced expression of CDK-1, as well as other cell cycle regulators (Hong *et al.* 1998; Rausch *et al.* 2015). *let-7* has been implicated in the repression of CDK-1, while upregulation of *cki-1* was shown to depend on LIN-29 and its transcriptional cofactor MAB-10<sup>NAB</sup> (Harris and Horvitz 2011). However, direct transcriptional control of *cki-1* by LIN-29/MAB-10 has not been demonstrated.

### Dauer development

*C. elegans* can adjust its development and cell division pattern dependent on environmental conditions. In addition to L1 arrest in the absence of food, conditions of high population density and limited food can induce an alternative third larval stage, with animals arresting as stress-resistant and long-lived dauer larvae (Riddle and Albert 1997; Antebi 2013). Dauer development coincides with a prolonged cell cycle arrest, through mechanisms that are only partly understood. Sensory neurons in the head perceive the presence of nutrients and dauer pheromone in the milieu, and release insulin-like peptides and TGF- $\beta$ -related ligands when conditions are favorable. The effects of these pathways on cell division are reasonably well understood for germ precursor cells (discussed below); however, it remains unclear how dauer-inducing systemic signals connect to the cell cycle of somatic cells. As in L1 arrest (see above), *daf-16*<sup>FoxO</sup>-dependent *cki-1*<sup>CIP/KIP</sup> expression, counteracted by IGF signaling, is probably at least partly responsible for the cell cycle arrest during dauer induction (Hong *et al.* 1998; Baugh and Sternberg 2006).

## The Control of DNA Replication

### Replication licensing in larvae

To ensure genome stability, it is essential that all genomic DNA is replicated fully, but only once per cell cycle. The replication licensing system ensures that each replication origin is activated only a single time in each S phase. Activating replication origins multiple times would result in the creation of an unstable “honeycomb” pattern of replicated DNA that has to be resolved by recombination, leading to genomic instability and gene amplification (Alexander and Orr-Weaver 2016).

The licensing system works by restricting the licensing of DNA replication origins to late M or G1 phase. The license is equivalent to the loading of the replicative helicase onto DNA replication origins. This loading is temporally separated from the activation of the helicase in S phase. Thus, the licensing system ensures that each origin can only be “fired” once per S phase because new replicative helicases cannot be loaded onto origins in S phase to allow the refiring of origins (Tanaka and Araki 2013). In animals and yeast, the regulation of replication licensing primarily involves two replication


licensing factors, Cdt1 and Cdc6, that load the MCM2-7 replicative helicase complex onto replication origins (Riera *et al.* 2014) (Figure 5). The replication origin is bound by the origin recognition complex (ORC). The core ORC is composed of subunits ORC2-5 (ORC-2, -3, -4, -5 in *C. elegans*), and it associates with ORC1 (ORC-1) and ORC6. While the core ORC complex and ORC1 are required for replication in animals, ORC6 is not required (DePamphilis *et al.* 2006). *C. elegans* ORC-1 through ORC-5 have been characterized (Sonneville *et al.* 2012). However, a *C. elegans* homolog of ORC6 has not been reported and is not readily apparent in homology searches.

In the process of DNA replication licensing, the licensing factor Cdc6 loads first to the ORC. Cdt1 is bound to the MCM2-7 replicative helicase and is required to load MCM2-7 onto the Cdc6–ORC complex. The ORC–Cdc6–MCM2-7 complex can recruit more replicative helicase complexes to the origin, so that each origin has several MCM2-7 complexes loaded (two of which will be utilized for bidirectional DNA replication). The loading of MCM2-7 forms the prereplicative complex (pre-RC).

The MCM2-7 helicase remains inactive during G1 phase and is activated at origins to initiate DNA replication during S phase. No further MCM2-7 complexes can be loaded onto replication origins in S phase because the licensing factors Cdt1 and Cdc6 are inactivated. In animal cells, Cdt1 is degraded and Cdc6 is exported from the nucleus. The regulation of the replication licensing system thus prevents DNA rereplication by temporally separating the loading of the replicative helicases (in late M or G1 phases) from their activation in S phase.

An additional licensing protein found in animals is the protein Geminin, which binds and inhibits Cdt1 in the S and G2 phases (Pozo and Cook 2017). In normal (full) somatic cell cycles, most Cdt1 is degraded at the initiation of S phase, and Geminin then binds to newly synthesized Cdt1 to prevent its activity in the S and G2 phases. However, other cell cycles, such as rapid embryonic cleavage divisions, may not invoke Cdt1 degradation, and so the inhibition of Cdt1 by Geminin is more important (Kermi *et al.* 2017). Geminin contains a destruction-box degron and its degradation is catalyzed by the APC/C ubiquitin ligase in mitosis. Geminin degradation releases Cdt1 to allow its participation in replication licensing in late M or G1 phases.

A critical regulator of replication licensing in animals and fission yeast is the CRL ubiquitin ligase CRL4–CDT-2, which contains CUL-4 as the scaffold and uses CDT-2 as the substrate receptor. The role of the CRL4 complex in this process was first identified in *C. elegans*. The *C. elegans cul-4* mutant has a dramatic rereplication phenotype, with somatic larval cells containing up to 100C DNA content (Zhong *et al.* 2003). Notably, the rereplication phenotype is not observed in embryos or germ cells, suggesting that replication licensing is regulated differently in those tissues. CUL-4 was shown to be required for the degradation of the replication licensing factor CDT-1 during S phase in larvae (Zhong *et al.* 2003) (Figure 5). It was subsequently shown that Cdt2 is the specific



**Figure 5** The regulation of DNA replication licensing in larval somatic cells. The prereplicative complex forms in G1 phase. In S phase, replication licensing is prevented by the nuclear export of CDC-6, which is initiated by the phosphorylation of CDC-6 on consensus CDK sites, and the ubiquitin-mediated degradation of CDT-1. Note that both processes are regulated by the ubiquitin ligase CRL4-CDT-2, which directly targets the degradation of CDT-1 and indirectly promotes CDC-6 nuclear export by targeting the degradation of the CDK-inhibitor CKI-1. See text for details.

CRL4 substrate receptor component for CDT-1 degradation in fission yeast and multiple animals (Havens and Walter 2011), including *C. elegans* (Kim and Kipreos 2007). Cdt1 degradation is restricted to S phase because CRL4-Cdt2 recognizes Cdt1 only when it is physically associated with proliferating cell nuclear antigen (PCNA), which is present in the DNA replication complex at replication forks (Havens and Walter 2011). Consistent with this mechanism, mutating *C. elegans* CDT-1 to remove the PCNA-binding site stabilizes the protein during S phase (Kim *et al.* 2007).

The CRL4-CDT-1 complex is a master regulator that restricts the activity of multiple replication licensing factors. In

addition to directly targeting CDT-1 for degradation, CUL-4 is also required for the nuclear export of the CDC-6 replication licensing factor (Kim *et al.* 2007). As in mammals, the nuclear localization of *C. elegans* CDC-6 is controlled by phosphorylation flanking its nuclear localization sequences during S phase to prevent its nuclear import (Kim *et al.* 2007). Because the putative nuclear export sequence is still active during S phase, CDC-6 becomes cytoplasmically localized. To promote the nuclear export of CDC-6, CRL4-CDT-2 targets the degradation of the CDK-inhibitor CKI-1. During a normal S phase, reducing CKI-1 levels allows sufficient CDK activity to phosphorylate CDC-6 on multiple CDK consensus sites clustered around the three putative nuclear localization sequences, which promotes CDC-6 nuclear export (Kim *et al.* 2007). If CUL-4 is inactivated, then CDC-6 does not become phosphorylated on these sites during S phase. Consequently, CDC-6 remains nuclear-localized, where it can function to reload the MCM2-7 helicase onto replication origins if CDT-1 is also present. The presence of nuclear CDT-1 and CDC-6 in *cul-4* mutants during S phase allows the refiring of DNA replication origins to induce rereplication. This CRL4-CDT-2 pathway was subsequently shown to be conserved in human cells, with CRL4-CDT-2 targeting the CKI-1 ortholog p21<sup>Cip1</sup> to regulate Cdc6 nuclear localization (Kim *et al.* 2008) (Figure 5).

#### **Tissue-specific differences in regulating replication: germ cells and embryos**

The observation that inactivating CRL4-CDT-2 does not result in an increase in the DNA content of germ cells or early embryonic cells, in contrast to larval somatic cells, suggests that different regulatory mechanisms control replication licensing in these tissues (Zhong *et al.* 2003; Kim *et al.* 2007). One potential difference is the contribution of the Cdt1-regulatory protein Geminin. The *C. elegans* Geminin homolog GMN-1 binds the CDT-1 protein, as expected, and *gmn-1* RNAi reduces the numbers of germ cells (Yanagi *et al.* 2005). However, the level of DNA in the germ cells of *gmn-1* null mutants is not increased relative to wild-type, implying that there is no rereplication (Kim and Kipreos 2007). Co-inactivating CRL4-CDT-2 and GMN-1 also does not increase DNA levels in germ cells, suggesting that there are other regulatory mechanisms to restrain replication licensing in germ cells (Kim and Kipreos 2007).

In the early embryo, which has S-M cycles that lack gap phases, replication licensing occurs during late M phase, with MCM2-7 loading onto chromosomes during metaphase and anaphase of mitosis (Korzelius *et al.* 2011a; Sonnevile *et al.* 2012). This early licensing is required to allow DNA replication to commence immediately after M phase. In contrast, replication licensing occurs during G1 phase in somatic cells of larvae (Zhong *et al.* 2003). GMN-1<sup>Geminin</sup> depletion in the early embryo allows MCM2-7 to bind earlier in meiosis (anaphase of meiosis I rather than anaphase of meiosis II) and earlier in mitosis (prometaphase rather than metaphase and anaphase). This indicates that GMN-1 contributes to the timing of replication licensing in the early embryo. However,



*gmn-1* RNAi produced no phenotypes in the early embryo beyond the earlier initiation of replication licensing (Sonneville *et al.* 2012).

Analysis of fluorescently tagged replication components indicates that the nuclear export of multiple pre-RC components prevents origin refiring during S phase in the early embryo (Sonneville *et al.* 2012). During S phase, the core ORC2-5 complex and the separate ORC-1 subunit are excluded from the nucleus, and will only rebind to chromatin in early mitosis prior to the loading of the MCM2-7 complex for the subsequent round of DNA replication (Figure 6). As in somatic larval cell cycles, CDC-6 is also exported from the nucleus during S phase. Additionally, CDT-1 appears to be excluded from S phase nuclei, although this conclusion is based on relatively weak antibody staining of the endogenous protein and should be further substantiated.

To determine if the observed nuclear exclusion of the pre-RC components is functionally important for replication licensing, Sonneville *et al.* (2012) depleted the nuclear export factor Exportin-1 (*XPO-1*) by RNAi. In *xpo-1*(RNAi) embryos, there was a significant delay in the nuclear export of the ORC subunits and CDC-6. Tellingly, this led to the rereplication of genomic DNA, indicating that the nuclear export of pre-RC components forms the major control preventing replication licensing during S phase in the early embryo (Sonneville *et al.* 2012).

These results indicate that different tissues use different replication licensing regulations to ensure that DNA is not overreplicated. The truncated cell cycle of the early embryo, comprising S–M–S–M cycles, presumably is not compatible with the degradation of CDT-1 in every cell cycle, as occurs in larval somatic cells, which then requires the translation of new CDT-1 in the G1 phase prior to the next S phase. Why germ cells require a different DNA replication licensing control is not clear, but it may similarly result from a truncated or absent G1 phase (see below). The exact replication licensing regulatory pathway operating in germ cells remains to be determined.

### **Bypassing the DNA replication checkpoint**

The *mcm-4/lin-6* gene encodes the MCM4 component of the MCM2-7 replicative helicase. *mcm-4* mutant larvae are defective in replicating their genomic DNA, as would be expected for inactivation of the replicative helicase complex (Korzelius *et al.* 2011a). Strikingly, despite not replicating their genomic DNA, *mcm-4* mutant somatic cells nevertheless undergo mitosis at the appropriate time to create daughter cells with reduced DNA levels (Sulston and Horvitz 1981; Korzelius *et al.* 2011a). This phenotype implies the absence of the DNA replication checkpoint that normally would prevent mitotic entry in response to unreplicated genomic DNA. *C. elegans* larval somatic cells do have a functioning DNA replication checkpoint because treatment with hydroxyurea (which stalls DNA replication fork movement) inhibits entry into mitosis (Euling and Ambros 1996). Most likely, the replicative helicase cannot form in the absence of MCM-4 and

therefore DNA replication never initiates, and because there is no DNA replication the DNA replication checkpoint never engages (Korzelius *et al.* 2011a).


The signal for the DNA replication checkpoint includes single-stranded DNA and stalled replication forks (Smith *et al.* 2010). However, if DNA replication does not initiate then there would be no stalled replication forks and the DNA replication checkpoint would not be expected to be activated. Loss of other replication components that prevent DNA replication from initiating would be expected to similarly bypass the DNA replication checkpoint. Consistently, loss of the replication licensing factors CDT-1 or CDC-6, or the helicase component MCM-5, produces a similar phenotype in embryos: continued cell division in the absence of DNA replication (Zhong *et al.* 2003; Kim *et al.* 2007; Korzelius *et al.* 2011a). This supports the model that the bypass of the DNA replication checkpoint results from the failure to initiate DNA replication.

While the *mcm-4(e1466)* homozygous mutant shows no observable DNA replication in somatic larval cells, there is a surprising level of germ cell proliferation. Germ cells still incorporate the thymidine analog 5-bromo-2'-deoxyuridine (BrdU) in *mcm-4(e1466)* larval mutants, while all somatic cells fail to incorporate BrdU (Korzelius *et al.* 2011a). This difference between somatic and germline tissues could result from *mcm-4* maternal product (mRNA or protein) in the germline that is sufficient for the continued DNA replication in the larval stages, or it may reflect a surprising lack of requirement for the full-length MCM-4 protein (as the *e1466* mutation introduces a premature stop codon at amino acid 88). Overall, this provides further evidence for differences in the regulation of DNA replication between germ cells and larval somatic tissues.

### **Regulation of the Cell Cycle in Germ Cells**

The cell cycle regulation of germ cells differs substantially from that of somatic cells. Most notably, germ cells are the only tissue that divides continuously in larval stages and adults. During the larval stages, the total number of germline cells increases rapidly, with germ cells dividing faster in larval stages than in the adult (Roy *et al.* 2016). Based on cell counts, the doubling time of germ cells in the L2 larval stage was reported to be ~4 hr (Kipreos *et al.* 1996). Cell counts for the L3-to-adult larval period produced an average doubling time of ~9 hr, but this includes a percentage of germ cells that enter the meiotic program and thus would no longer be actively cycling (Roy *et al.* 2016).

In adults, germ cell proliferation is restricted to the distal stem cell niche, which is localized within the region of the gonad encompassed by the somatic DTC (Figure 7A). The DTC forms a cap around the first 3–4 germ cell diameters, but also has extensive intercalating protrusions that extend 6–8 cell diameters from the distal end (Byrd *et al.* 2014). These DTC intercalations define the extent of the DTC “plexus” (Figure 7A). The DTC plexus is considered to be



**Figure 6** The regulation of DNA replication licensing differs between the early embryo and larval somatic cells. (A) The control of replication licensing in larval somatic cells (see Figure 3 for components). (B) The regulation of replication licensing in the early embryo. Replication licensing occurs in mitosis. Replication licensing is prevented from reinitiating in S phase by the export of CDC-6 and the ORC complex (ORC1 and ORC2-5 exported separately). It is not known if CDC-6 remains associated with ORC components during S phase or if CDC-6 nuclear export is regulated by CDK phosphorylation (as in larval somatic cells). See text for details.


the true stem cell niche that encloses the GSC pool (Byrd *et al.* 2014); while in more proximal regions of the proliferative zone, germ cells are in various states of meiotic differentiation (Cinquin *et al.* 2010; Fox and Schedl 2015; Brenner and Schedl 2016). An example of more proximal cells having increased meiotic tendencies is that upon arresting cells in mitosis by inactivating the APC/C ubiquitin ligase, cells in the DTC plexus niche remain undifferentiated, while cells that are more proximal enter meiosis (Cinquin *et al.* 2010).

Within the adult proliferative zone, all germ cells incorporate BrdU, suggesting that they are all either mitotically cycling or completing premeiotic S phase (Crittenden *et al.* 2006). The length of germ cell cycles has been inferred from analysis of 5-ethynyl-2'-deoxyuridine (EdU) incorporation, labeling cells in S phase, and phosphohistone H3 staining of mitotic cells. The reported doubling times of germ cells in the proliferative zone of adult hermaphrodites vary from ~6.15 to 16–24 hr (Crittenden *et al.* 2006; Fox *et al.* 2011; Fox and Schedl 2015; Rosu and Cohen-Fix 2017). However, as the proliferative zone contains at least two distinct populations, the distal GSC niche and a more proximal premeiotic region, which regions are analyzed could have a significant impact on the cell cycle timing. Live imaging of germ cells using a photoconvertible marker showed that the region of the GSC niche, the most distal five to seven rows of germ cells, divided at the fastest rate, with on average one division every 6 hr (Rosu and Cohen-Fix 2017). The middle five to seven rows divided at an intermediate rate, and the most proximal five to seven rows showed hardly any cell division, suggesting that these cells are premeiotic. These data support the idea that, within the most distal region, all GSCs are equivalent for proliferation and stochastically enter meiosis as they are displaced proximally from the DTC plexus (Rosu

and Cohen-Fix 2017). Thus, it appears that the stem cell character of GSCs is defined by their location within the niche.

GSC cell cycles differ from somatic larval-stage cell cycles in having a very short or absent G1 phase. When analyzing the entire proliferative zone, relatively few cells with 2C DNA content are observed (Fox *et al.* 2011; Roy *et al.* 2016); however, within one to five cell diameters of the distal end, a notable percentage of cells have 2C DNA content. It is possible that a G1 phase is only present in the most distal region. However, an argument against a G1 phase follows from the observation that inactivation of **CDK-4** does not affect germ cell proliferation, even though it is essential for G1 phase progression in somatic cells (Park and Krause 1999; Boxem and van den Heuvel 2001). In contrast, inactivation of the G1/S cyclin **CYE-1** and its binding partner **CDK-2** inhibits germ cell proliferation (Fox *et al.* 2011). However, as we discussed previously, it has not been excluded that the **CDK-4/cyclin D** complex is only required when cells are transitioning into the cell cycle from the G0 phase. Under normal laboratory growth conditions, GSCs do not have a detectable G0 phase. Thus, it is possible that distal GSCs have a G1 phase despite not requiring **CDK-4/cyclin D** activity.

Another major difference between germ cells and other tissues is their syncytial structure, in which germ cells are in contact with a shared cytoplasm (the “rachis”) through an opening in their plasma membrane (Figure 7A). Germ cells begin to acquire their syncytial structure in the L2 larval stage and all germ cells are syncytial in adults (Amini *et al.* 2014). The most obvious function of the syncytial opening to the rachis is to allow maternal mRNA and protein that is created by germ cells to flow to the developing oocytes. Maternal product is observed over the entire length of the rachis (Gibert *et al.* 1984).



**Figure 7** The hermaphrodite gonad and the regulation of mitotic proliferation. (A) Diagram of a cross section of one arm of the hermaphrodite gonad. Germ cell nuclei are green circles. The somatic distal tip cell (DTC) at the distal end of the gonad is colored yellow. The germ cells line the side of the gonad and only one layer is shown in the cross section. The plasma membranes around the nuclei (black lines) include an opening to the central cytoplasm (rachis). (B) Simplified diagram of the interplay of several regulators of mitotic proliferation within the proliferative zone. The Delta ligands LAG-2 and APX-1 originate from the DTC, while other components are in the distal germ cells or in the transition zone as germ cells transition from mitotic proliferation to meiosis. See text for details.

The syncytium does not appear to transmit cell cycle-state information. Active mitotic CDK1–cyclin B has been shown, when introduced into vertebrate cells, to induce the activation of CDK1–cyclin B in the recipient cell that then drives the cell into mitosis (Kishimoto 2015). However, in the *C. elegans* gonad, the entry of one germ cell into mitosis does not trigger mitosis in the surrounding germ cells. This suggests that active CDK1–cyclin B does not transfer from mitotic germ cells to nearby germ cells through their openings to the rachis.

#### Cell cycle regulators in the mitosis vs. meiosis transition

There are two major levels of control of germ cell proliferation: (1) regulation of the size of the GSC pool, which is controlled by regulating the mitosis vs. meiosis transition, and (2) regulation of the rate of germ cell proliferation. The regulation of GSC differentiation to enter meiosis has been extensively reviewed elsewhere (Hansen and Schedl 2013; Kimble and Seidel 2013). Therefore, we will focus on those aspects of the mitosis vs. meiosis decision that are known to interface with cell cycle regulators.

Notch signaling by the DTC is the major regulator of the mitosis vs. meiosis decision [reviewed in Hansen and Schedl (2013) and Kimble and Seidel (2013)]. The DTC expresses the DSL (Delta, Serrate, LAG-2) ligand proteins LAG-2 and APX-1, which are bound by the Notch receptor GLP-1 on germ cells. After GLP-1 activation, the intracellular domain of GLP-1 is cleaved to release an intracellular domain (GLP-1-ICD), which acts as a transcription factor in complex with the CSL DNA-binding protein LAG-1 and the transcriptional coactivator LAG-3/SEL-8 (Figure 7B). GLP-1 activity leads to the

inhibition of GLD-1, a K homology (KH) domain RNA-binding protein that inhibits mitotic proliferation. GLP-1 also leads to the inhibition of GLD-2 and GLD-3, a cytoplasmic poly(A) polymerase and KH-domain RNA-binding protein, respectively, which promote meiosis.

The critical target genes for GLP-1-ICD/LAG-1 transcription are *lst-1* and *sygl-1*, which are redundantly required to maintain GSCs in the mitotic state (Kershner *et al.* 2014) (Figure 7B). *lst-1* and *sygl-1* encode novel proteins, LST-1 and SYGL-1, which inhibit GLD-1 activity by acting in association with the translational repressor FBF-1 (a PUF, Pumilio and FBF, family protein) (Brenner and Schedl 2016). In addition, LST-1 also inhibits GLD-1 activity independent of FBF-1 (Figure 7B). GLP-1 signaling therefore controls the meiosis vs. mitosis cell fate decision, at least in part, by inhibiting GLD-1. The mitotic index is unchanged in *glp-1* loss-of-function mutants that have reduced sizes of the proliferative zone (Fox and Schedl 2015; Roy *et al.* 2016). Thus, while GLP-1 signaling controls the size of the GSC niche, GLP-1 does not appear to regulate cell cycle progression directly.

A failure of germ cells to enter meiosis and instead continue mitotic division produces germline tumors, in which mitotic germ cells are present throughout the gonad. Germline tumors can arise from constitutive activation of Notch signaling, e.g., through GLP-1 gain-of-function mutations, which promotes mitotic proliferation and inhibits meiosis. Germline tumors can also arise from loss of the RNA regulators that promote meiosis and block mitotic proliferation. A complete failure to enter meiosis occurs upon loss of both the GLD-2 and GLD-3 pathway, and the GLD-1 and NOS-3 pathway (with NOS-3 a Nanos-related translational regulator). In this situation, germ cells divide mitotically even when distant from the Notch signal in the stem cell niche.

**Cyclin E expression to maintain germ cells in a mitotic state:** The G1/S cyclin CYE-1 protein is present in the proliferative zone but not in the adjacent meiotic regions of the germline (Brodigan *et al.* 2003). Within the proliferative zone, CYE-1 expression is high in all germ cells, indicating that its expression is not cell cycle regulated (Fox *et al.* 2011). It is possible that the constant CYE-1 expression contributes to the apparent lack of a G1 phase in most germ cells, as other cells that have constant cyclin E expression, such as mammalian stem cells, also lack G1 phases (Orford and Scadden 2008).

CYE-1 expression in germ cells is important not only to allow proliferation within the proliferative zone but also to prevent entry into meiosis. This was shown by RNAi of *cye-1* in L4 larvae, which blocked cell proliferation and induced germ cells to prematurely enter meiosis (Fox *et al.* 2011). In contrast, inactivation of other cell cycle regulators that are important for S and G2 phases does not induce meiotic entry (Fox *et al.* 2011). This suggests that either the decision to enter meiosis is directly regulated by CDK-2/CYE-1, or that it occurs early in the cell cycle (pre-S phase) and loss of CYE-1 causes arrest at that decision point.

GLD-1, which inhibits mitotic proliferation, and CDK-2/CYE-1, which promotes mitotic proliferation, mutually repress each other (Figure 7B). GLD-1 inhibits CYE-1 expression in the meiotic regions (Biedermann *et al.* 2009). The repression is direct and is mediated by GLD-1 binding to its target sequence in the *cye-1* mRNA 3'-UTR (Biedermann *et al.* 2009). In *gld-1* mutants, meiotic cells reenter the mitotic cell cycle, and this mitotic proliferation requires the derepression of CYE-1 that occurs in these mutants (Biedermann *et al.* 2009). Conversely, CYE-1 negatively represses GLD-1 protein expression in the distal region of the gonad. GLD-1 is normally absent from the most distal region of the gonad, but becomes expressed there when CYE-1 is inactivated (Jeong *et al.* 2011). This negative regulation of GLD-1 by CDK-2/CYE-1 also appears to be direct. CDK-2/CYE-1 can phosphorylate GLD-1 *in vitro*, and inactivation of CYE-1 causes the loss of GLD-1 phosphorylation *in vivo* (Jeong *et al.* 2011). Replacement of the CDK-target sites in GLD-1 with nonphosphorylatable residues extends GLD-1 protein expression to more distal regions, thereby reducing the size of the proliferative zone. Therefore, CDK-2/CYE-1 phosphorylation of GLD-1 is important for GLD-1 regulation and determining the size of the stem cell niche.

The expression of CYE-1 is also subject to negative regulation by the SCF-PROM-1 E3 ubiquitin ligase (Fox *et al.* 2011) (Figure 7B). Inactivation of the substrate receptor component of the E3 complex, PROM-1, allows CYE-1 expression to extend beyond its usual sharply defined region in the proliferative zone. This suggests that CYE-1 is subject to ubiquitin-mediated degradation as germ cells leave the proliferative zone, in addition to its translational regulation by GLD-1. Despite ectopic CYE-1 expression, the *prom-1* mutant germ cells still cease mitotic proliferation and enter meiosis. This indicates that CYE-1 expression alone is not sufficient to maintain mitotic proliferation in germ cells. Double mutants of *prom-1*, with either *gld-1/nos-3* or *gld-2/gld-3* pathway genes, reveals that the three pathways function redundantly to promote entry into meiosis and the cessation of mitosis, with additional SCF-PROM-1 targets, beyond CYE-1, that impact this transition (Mohammad *et al.* 2018).

#### **FBF represses CKI-2 expression to prevent germ cell arrest:**

FBF-1 and FBF-2 are closely related PUF family RNA-binding proteins. FBF-1 and FBF-2 show the highest level of expression in the proliferative zone (Lamont *et al.* 2004). Together, FBF-1 and FBF-2 are required to maintain germ cells in the mitotic state (Crittenden *et al.* 2002). FBF represses the translation of the CDK inhibitor CKI-2 in the proliferative zone through direct interaction with the *cki-2* 3'-UTR (Kalchhauser *et al.* 2011). Inactivation of both FBF-1 and FBF-2 causes the loss of all mitotic germ cells, at least in part through the expression of CKI-2. If CKI-2 is co-inactivated with FBF-1/2 then the loss of mitotic germ cells is delayed. Thus, FBF promotes germ cell proliferation in part by repressing CKI-2 expression.

**CRL2-LRR-1 promotes mitotic proliferation and prevents checkpoint activation:** Inactivation of the ubiquitin ligase CRL2-LRR-1 leads to the accumulation of CKI-1 in all germ cells (Feng *et al.* 1999; Starostina *et al.* 2010). In the absence of CRL2-LRR-1 activity, CKI-1 protein accumulates and germ cells arrest with a 2C DNA content. CRL2-LRR-1 appears to directly target CKI-1, as LRR-1 binds CKI-1 and mediates its degradation when the two proteins are coexpressed in mammalian cells (Starostina *et al.* 2010).

Loss of LRR-1 also activates the DNA replication checkpoint, as indicated by the presence of the phosphorylated active form of the CHK-1 kinase in embryos (Merlet *et al.* 2010). The *lrr-1* mutant germ cell arrest can be abrogated by inactivation of CHK-1 or its upstream activator, ATL-1<sup>ATR/ATM</sup> kinase. The activation of the DNA replication checkpoint in *lrr-1* mutants is associated with the accumulation of single-strand DNA-binding replication protein A (RPA-1) on chromatin, suggesting a defect in completing DNA replication (Merlet *et al.* 2010).

#### **Regulation of the rate of germ cell proliferation**

The germline in an adult hermaphrodite contains approximately twice as many cells (~2000) as the entire soma (~1000) (Kimble and Seidel 2013). A substantial percentage of total energy expenditure in *C. elegans* adults goes toward germ cell proliferation and the creation of oocytes that are loaded with maternal product. Given the scale of the energy consumption required for reproduction, it is not surprising that the rate of germ cell proliferation is closely linked to the level of food intake. A surprisingly large number of regulatory pathways control the rate of germ cell proliferation in the adult hermaphrodite. We will describe these pathways in this section.

**TGF- $\beta$  promotes the mitotic germ cell fate:** Environmental conditions can impact the rate of germ cell proliferation. These include the availability of food and signals that indicate that the food supply will soon be exhausted by the presence of many other nearby nematodes, each of which is capable of generating a steep exponential increase of 250–300 progeny approximately every 6 days. The TGF- $\beta$  signaling pathway is activated in response to favorable environmental conditions, including plentiful food, and is repressed in response to unfavorable conditions that includes crowding or a lack of food (Gumienny and Savage-Dunn 2013). The chemosensory ASI neurons generate the TGF- $\beta$  ligand (DAF-7) in response to low pheromone levels (signifying noncrowded conditions) and plentiful food (Dalfó *et al.* 2012).

The TGF- $\beta$  signaling pathway is required for creating the full complement of larval germ cells (Dalfó *et al.* 2012). Inactivation of TGF- $\beta$  signaling reduces germ cell numbers by half and produces a smaller proliferative zone (Dalfó *et al.* 2012). However, the mitotic index within the proliferative zone is similar to wild-type (Dalfó *et al.* 2012). This suggests that TGF- $\beta$  signaling regulates the mitosis-to-meiosis decision to control the size of the GSC pool, rather than the rate of germ cell division.

TGF- $\beta$  signaling functions in the somatic DTC to indirectly regulate germ cells (Dalfó *et al.* 2012). Intriguingly, the regulation of germ cells by TGF- $\beta$  is independent of GLP-1 Notch signaling, as loss of TGF- $\beta$  signaling still reduces germ cell numbers in a tumorous mutant that lacks the GLP-1 receptor (Dalfó *et al.* 2012). The TGF- $\beta$  pathway functions in DTCs to increase the size of the GSC pool by inhibiting entry into meiosis. TGF- $\beta$  acts through its canonical pathway to inhibit the DAF-3/DAF-5 transcriptional repressor complex (Dalfó *et al.* 2012), but the relevant downstream gene targets have not been reported.

**Insulin/IGF-like signaling promotes germ cell proliferation:** IGF signaling cell-autonomously promotes germ cell proliferation during the larval stages (Michaelson *et al.* 2010). Loss of IGF signaling reduces the mitotic index, indicating that germ cell proliferation is affected rather than the mitosis-to-meiosis transition (Michaelson *et al.* 2010). IGF signaling affects germ cells through its canonical inhibition of the FoxO transcription factor DAF-16 (Michaelson *et al.* 2010). The critical transcriptional targets of DAF-16 that inhibit germ cell proliferation have not been reported.

A partial reduction-of-function allele of the *daf-2* insulin receptor reduces the mitotic index of larval-stage germ cells and adult germline tumors (in which mitotic proliferation of germ cells occurs throughout the gonad as the result of particular mutations) (Pinkston *et al.* 2006; Michaelson *et al.* 2010). In contrast, the adult proliferative zone is not affected by partial *daf-2* mutants. However, stronger *daf-2* mutations do reduce germ cell proliferation in the adult (Narbonne *et al.* 2015). This suggests that IGF signaling regulates germ cell proliferation at all stages, but that larval stages and germline tumors are more sensitive to changes in IGF signaling.

There are 40 *C. elegans* genes that encode insulin-like peptides, *INS-1* through *INS-39* and *DAF-28* (Baugh *et al.* 2011). Interestingly, germ cell proliferation appears to be solely stimulated by *INS-3* and *INS-33* (Michaelson *et al.* 2010). The expression of *ins-3* is largely neuronal and *ins-33* expression is largely hypodermal, suggesting that these are the sources of systemic *INS-3* and *INS-33* peptides (Michaelson *et al.* 2010). Other insulin-like peptides have been shown to control IGF signaling in other tissues, which indicates that insulin-like ligands have specialized tissue functions and that the animal can regulate germline proliferation via IGF signaling independently of other tissues (Michaelson *et al.* 2010).

**DAF-18<sup>PTEN</sup> signaling decreases germ cell proliferation in response to unfertilized oocytes:** Germ cell proliferation decreases in hermaphrodites when their sperm are depleted and this can be reversed by additional sperm provided by mating (Narbonne *et al.* 2015). The inhibition of germ cell proliferation depends on the accumulation of unfertilized oocytes in the spermless gonads. The decrease in germ cell proliferation in response to unfertilized oocytes requires DAF-18<sup>PTEN</sup> but not DAF-16<sup>FoxO</sup>, suggesting that DAF-18 functions in a DAF-16-independent manner (as it does for

the regulation of Z2 and Z3 in the L1 stage) (Narbonne *et al.* 2015).

**Bacterial folates are an exogenous signal to promote germ cell proliferation:** Folates are B-complex vitamins that function to create a subset of nucleosides and amino acids in a metabolic cycle termed “one-carbon metabolism” (Selhub 2002). Animals cannot synthesize folates, in contrast to bacteria, hence *C. elegans* obtain folates from their diet of bacteria. Recently, it was discovered that bacterial folates act as an exogenous signal to cell-autonomously stimulate germ cell proliferation (Chaudhari *et al.* 2016). Bacterial folates stimulate increased DNA replication of isolated germ cells *in vitro*, increased mitotic index and the size of the proliferative zone in wild-type animals, and the growth of germline tumors (Chaudhari *et al.* 2016).

Folates are a family of related molecules that can be interconverted as part of the one-carbon metabolism cycle. Interestingly, only a subset of bacterial folates stimulates germ cell proliferation. Stimulatory folates are 10-formyl-THF-Glu(n) and 5,10-methenyl-THF-Glu(n). The latter folate converts to 10-formyl-THF-Glu(n) at neutral pHs, such as normal culture conditions; this suggests that 10-formyl-THF-Glu(n) is the only stimulatory folate. Other folates that take part in the one-carbon metabolism cycle are unable to stimulate germ cell proliferation under normal (folate replete) growth conditions (Chaudhari *et al.* 2016).

Significantly, the folate stimulation of germ cell proliferation occurs independently of the one-carbon metabolism cycle. Both stimulatory and nonstimulatory folates can rescue folate deficiency, indicating that the ability of stimulatory folates to increase germ cell proliferation rates under normal culture conditions is not linked to their ability to act as a vitamin. The stimulation of germ cell proliferation by folates requires the homolog of the mammalian folate receptor, FOLR-1 (Chaudhari *et al.* 2016). Yet, FOLR-1 is not needed to provide folates as vitamins, which instead depends on the reduced folate carrier FOLT-1, whose inactivation produces severe folate deficiency phenotypes (Balamurugan *et al.* 2007; Austin *et al.* 2010). Additionally, a folate-related molecule, dihydropteroate, which cannot take part in one-carbon metabolism, also stimulates germ cell proliferation in a FOLR-1-dependent manner (Chaudhari *et al.* 2016). Interestingly, the stimulatory folate, 10-formyl-THF-Glu(n), and dihydropteroate are both relatively unstable relative to other folates. It is possible that the use of these unstable folates for signaling allows a tighter linkage between the availability of high-quality bacterial food (*e.g.*, live bacteria) and the rate of germ cell proliferation. The intracellular pathway for FOLR-1-dependent folate signaling is currently not known.

**Volatile bacterial odors increase reproductive rate via neuronal signals:** Volatile odors, which can transmit through the air from specific bacteria, can affect the rate of egg production (Sowa *et al.* 2015). Odors from the *Escherichia coli* strain HB101, even when physically separated from the

animals, causes hermaphrodites to lay a normal-sized brood over a shorter period of time compared to the rate with a normal diet of *OP50* bacteria. The accelerated rate of egg production when grown with *HB101* odors is correlated with increased proliferation of germ cells in the proliferative zone. The AWB chemosensory neurons mediate the response to the volatile odors via release of neuropeptides that are required for the stimulation of germ cell proliferation. Genetic interactions suggest that the AWB neuropeptide signaling is independent of insulin and TGF- $\beta$  signaling pathways. Currently, the neuropeptides and their target tissues have not been reported.

**The regulation of germ cell numbers upon starvation:** The starvation of L4-stage hermaphrodite larvae causes a dramatic reorganization of the adult germline, wherein the germ cells are dramatically reduced in number (Angelo and Van Gilst 2009). Upon starvation, mitotic germ cells in the proliferative zone undergo a G2-phase cell cycle arrest (Seidel and Kimble 2015). During starvation, adult hermaphrodites create one oocyte at a time over the course of many hours, in contrast to the situation in fed adult hermaphrodites where multiple developing oocytes are present in each gonad arm (Seidel and Kimble 2011). Embryos created during starvation either hatch in the hermaphrodite parent, leading to the death of the parent through “bagging” (eating the insides of the parent), or the embryos die *in situ*, presumably as a result of defects associated with the starvation (Seidel and Kimble 2011). This starvation state has been termed “adult reproductive diapause” (ARD) (Angelo and Van Gilst 2009) and “oogenic germline starvation response” (Seidel and Kimble 2011). The latter term is in reference to the fact that the shrinkage of the gonad corresponds to the creation of oocytes, presumably as gonadal cytoplasm is transferred to the developing oocytes (Seidel and Kimble 2011). A percentage of GSCs survive starvation in the DTC plexus. When starved animals are provided with food, these remaining GSCs repopulate the germline, which enlarges to the same size as the germlines of continuously fed animals (Angelo and Van Gilst 2009).

**Dafachronic acid inhibits germ cell proliferation in adults and mediates the adult starvation response:** Placing young adult hermaphrodites into starvation conditions results in an increase in the bile acid-like steroid hormone dafachronic acid (DA) as well as the cytochrome P450 enzyme *DAF-9*, which is required to create DA (Thondamal *et al.* 2014). Adult hermaphrodites that are starved have reductions in the number of germ cells in the proliferative zone within 1 day. This reduction of the proliferative zone requires DA; in the absence of DA, the proliferative zone maintains the same number of germ cells over the first 1–2 days of starvation (Thondamal *et al.* 2014) (H. Aguilaniu, personal communication). Addition of DA inhibits germ cell proliferation *in vivo* in both wild-type animals and tumorous germline mutants. The inhibition depends on the canonical DA-steroid hormone receptor *DAF-*

12 (Mukherjee *et al.* 2017). DA can inhibit the proliferation of isolated germ cells *in vitro* in a *DAF-12*-dependent manner, indicating a direct effect of DA on germ cells (Mukherjee *et al.* 2017). *DAF-12* is also required in germ cells for *in vivo* inhibition, again suggesting that DA acts directly on germ cells. Inactivation of *DAF-9 in vivo* (which would block DA production) increases the rate of germ cell proliferation in wild-type and tumorous germline mutants, suggesting that DA normally functions to restrain germ cell proliferation.

DA has multiple roles in *C. elegans*, including inhibiting entry into the dauer pathway, regulating heterochrony, and promoting longevity in animals that lack germlines (Antebi 2013). Significantly, the larval role of DA in blocking entry into the dauer pathway has the indirect effect of increasing germ cell proliferation by preventing the dauer-stage arrest of all cell divisions. This potential functional paradox is solved by the finding that DA only inhibits germ cell proliferation in adults but not in larvae (Mukherjee *et al.* 2017). Thus, DA indirectly promotes the proliferation of germ cells in larvae by blocking dauer entry, and directly inhibits germ cell proliferation in adults.

Adult germ cells are more sensitive to starvation than larval germ cells, with the mitotic index in adults dropping significantly within 30 min and a complete shut down by 3 hr. In contrast, in larval stages, the germ cell mitotic index is only partially reduced after 4–5 hr of starvation, with a complete shut down only by 7–8 hr (Seidel and Kimble 2015). The adult-specific role of DA in inhibiting germ cell proliferation may contribute to the more rapid cessation of germ cell proliferation upon starvation in adults.

Further research will be required to link the distinct regulatory signals that control the rate of germ cell proliferation (described in this section) to cell division regulators that directly impact the cell cycle and, in the case of *CDK-2/CYE-1*, also regulate the mitosis-to-meiosis decision.

## The Interplay Between Cell Proliferation, Arrest, and Differentiation

Developmentally induced signal transduction pathways and asymmetric cell divisions provide spatiotemporal control of gene expression. Transcriptional regulation of cell cycle genes is thought to form an important connection between developmental signals and the cell cycle. However, only a few molecular connections have been firmly established, and it remains poorly understood what mechanisms underlie the lineage-specific patterns of cell division and timing of terminal differentiation. In this section, we summarize insight in how transcription factors and chromatin regulators interact with the cell cycle to control the temporal pattern of cell division, and coordinate proliferation with differentiation in a lineage-specific manner.

### *Rb/E2F-mediated transcriptional repression*

The best known transcriptional regulators of the cell cycle are the heterodimeric E2F/DP transcription factors and their

binding partners of the Rb-corepressor family (van den Heuvel and Dyson 2008). Rb/E2F complexes act in concert with other transcriptional regulators and exert important functions in developmental control that go well beyond cell cycle regulation. Studies in *Drosophila* highlighted the contribution of repressive E2Fs in association with a fly Rb member in restricting gene expression outside the appropriate cellular context (Dimova *et al.* 2003). *C. elegans* *efl-1*<sup>E2F</sup>, *dpl-1*<sup>DP</sup>, and *lin-35*<sup>Rb</sup> match this paradigm, as follows from their activity as class B synthetic multivulva (synMuv) genes. Genetic analyses have shown that class B synMuv genes act redundantly with class A genes in the transcriptional repression of *lin-3*<sup>EGF</sup> throughout the animal (Cui *et al.* 2006; Saffer *et al.* 2011). Consequently, synMuv A,B double mutation leads to ectopic expression of *lin-3*<sup>EGF</sup> in hyp-7, which induces neighboring VPCs to adopt a vulval fate, and causes the formation of multiple vulvas.

Remarkably, at least seven genetically identified synMuvB genes encode proteins that form a conserved repressor complex (Harrison *et al.* 2006). This DRM (DP, Rb, MuvB) complex is closely related to DREAM (*Drosophila* RBF, E2F, and Myb), which was identified through biochemical purification of Rb/E2F complexes from *Drosophila* embryos (Korenjak *et al.* 2004). A similar mammalian DREAM complex contains repressive E2F4/E2F5 and Rb-related proteins p130 or p107, as well as homologs of five synMuvB proteins (Sadashivam and DeCaprio 2013). Interestingly, mammalian DREAM acts to repress cell cycle genes in quiescent cells. In contrast, the fly and worm complexes appear most critical for restricting tissue-specific gene expression (van den Heuvel and Dyson 2008). In addition to its contribution in *lin-3* repression, DRM acts to prevent expression of germline-specific genes in somatic cells. Moreover, at least a subset of synMuv B genes, including *lin-35*, *efl-1*, and *efl-2*, also inhibits cell cycle entry (Boxem and van den Heuvel 2002).

Transcriptional studies confirmed the dual roles for LIN-35 and DRM in repressing developmental as well as cell cycle genes (Kirienko and Fay 2007; Goetsch *et al.* 2017). In addition, ectopic expression of G1 CDK-cyclins in differentiated muscle cells induced the expression of a distinct set of > 200 genes with a strong cell cycle signature and enriched for promoters with E2F-binding sites (Korzelius *et al.* 2011b). These data suggest that LIN-35/E2F-mediated repression normally contributes to the postmitotic state of muscle cells and that active CDKs primarily antagonize the repression of cell cycle genes. EFL-1 genetically behaves as a transcriptional repressor and its predicted amino acid sequence is most similar to repressive E2F4/5 (Ceol and Horvitz 2001).

Surprisingly, activating E2Fs have not unequivocally been identified in *C. elegans*. EFL-3 resembles atypical E2F7/8, which repress a subset of E2F targets in mammalian systems (van den Heuvel and Dyson 2008). EFL-2 is somewhat closer to activating E2Fs in predicted amino acid sequence, but RNAi studies have not revealed cell cycle or synMuv B functions (Ceol and Horvitz 2001; Boxem and van den Heuvel

2002). E2F (the E2F/DP heterodimer) may potentially act as a transcriptional activator in the gonad, as *efl-1* and *dpl-1* promote the expression of oogenesis and embryogenesis-related genes, independently of *lin-35* (Chi and Reinke 2006).

The strongest indication that activating E2Fs exist in *C. elegans* is provided by the dual phenotype associated with loss of *dpl-1*<sup>DP</sup>. RNAi of *dpl-1* suppresses the cell cycle arrest of *cyd-1* mutants, similar to *efl-1*, which indicates that *dpl-1* normally acts to inhibit cell cycle entry. At the same time, *dpl-1* loss interferes with the expression of an S phase reporter and the proliferation of ventral cord precursor cells, which implies a cell cycle-promoting role (Ceol and Horvitz 2001; Boxem and van den Heuvel 2002). A cell cycle-promoting function of DPL-1 would be expected to depend on association with an E2F partner, but an activating partner has not been identified. Incomplete inactivation of *efl-2* or redundancies among the three *C. elegans* E2F-related genes may have masked this role.

### Lineage-dependent regulation of the cell cycle

In other model organisms, transcriptional regulation of D-type cyclins and CKIs is an important mechanism for connecting developmental and environmental signals to the cell cycle. Similarly, *C. elegans* *cyd-1* cyclin D and *cdk-4* are transcriptionally activated coincident with cell cycle entry, and in a lineage-specific manner (Park and Krause 1999). Several critical DNA sequences that are evolutionarily conserved have been identified in the enhancer/promoter regions (Brodigan *et al.* 2003). While these sequences are putative transcription factor-binding sites, essential *trans*-activating factors have not been described.

In contrast to *cyd-1* and *cdk-4*, genetic experiments have identified transcription factors as upstream *cki-1* regulators. DAF-16<sup>FoxO</sup> promotes *cki-1* expression during L1 and dauer arrest, and transcription factors have been implicated in lineage-specific cell cycle arrest through *cki-1* induction (described above; Baugh and Sternberg 2006). For example, the VPCs are formed during the L1 stage and remain quiescent until the induction of vulva formation in L3 larvae (Sternberg 2005) (Figure 2C). This arrest depends on *cki-1*, as RNAi of *cki-1* causes all VPCs to undergo an extra cell division in L2 (Hong *et al.* 1998). A screen for extra VPC cell divisions in L2 identified the CDC-14 phosphatase, as well as the LIN-1<sup>Ets</sup> and LIN-31<sup>FoxB</sup> transcription factors, as positive regulators of *cki-1* expression in the VPCs (Saito *et al.* 2004; Clayton *et al.* 2008).

The LIN-1 and LIN-31 transcription factors are downstream targets of the EGFR-RAS-MAPK-related pathway that induces vulva formation (Sternberg 2005). MAPK-mediated inhibition of LIN-1 and LIN-31 in the L3 larval stage induces one of the VPCs (P6.p) to adopt a primary vulval fate, which coincides with three full rounds of cell division (Figure 2C). Thus, the transcription factors that induce *cki-1* expression in the L2 stage are downregulated at the time of cell cycle re-entry. This indicates a potential direct link between EGFR-RAS-MAPK-signaling and the cell cycle (Clayton *et al.* 2008).

Indeed, genetic experiments support the notion that LIN-31 induces *cki-1* transcription, and represses *cdk-4* and *cye-1* transcription, until phosphorylation by MAPK inhibits LIN-31 in L3 larvae (Roiz *et al.* 2016). In addition, the homeodomain transcription factor LIN-39 promotes VPC proliferation by preventing their fusion with the general hypodermis, and activating the transcription of *cye-1* and other cell cycle genes in the VPCs (Roiz *et al.* 2016).

In addition to EGFR-RAS-MAPK signaling, Notch and Wnt-signaling pathways coordinate induction of the vulval cell fates with multiple rounds of cell division. VPCs that do not adopt a vulval fate go through a single division cycle before fusing with the general hypodermis (Figure 2C; P3.p, P4.p, P8.p). Thus, the signals that trigger vulval induction appear to expand the intrinsic cell proliferation potential of the VPCs. Some components of the mediator complex, which couples sequence-specific transcription factors to the RNA Polymerase II complex, are required for the VPC quiescence in L2 and vulval induction in L3 (Sternberg 2005; Clayton *et al.* 2008). While the mechanism is not understood, these contributions specifically involve a CDK8 module of the mediator complex, which has been implicated in transcriptional repression.

The LIN-3<sup>EGF</sup> signal that activates the LET-23<sup>EGFR</sup>-LET-60<sup>Ras</sup>-MPK-1<sup>MAPK</sup> pathway and induces a primary vulval cell fate comes from a specific cell in the somatic gonad. This “anchor cell” (AC) also organizes the connection between the gonad and vulva, which involves breaking down the basement membranes between these organs. AC signaling and invasion depend on the nuclear hormone receptor transcription factor NHR-67 (Matus *et al.* 2015). Interestingly, loss of *nhr-67* results in continued cell division of ACs that fail to invade, while *cki-1* induction arrests cell division and restores basement membrane invasion. The data are consistent with a model in which NHR-67 induces *cki-1* transcription to arrest cell division of the AC, while the arrest in G1 is critical for the fully differentiated AC state (Matus *et al.* 2015).

### **Coordinating cell cycle arrest and terminal differentiation**

Cell proliferation and terminal differentiation in general appear to be mutually exclusive processes, but the underlying regulatory mechanisms are still poorly understood. The fully differentiated state is acquired through a gradual process in which sequential binary decisions progressively restrict developmental potential (Kaletta *et al.* 1997; Bertrand and Hobert 2010). Initially, this lineage restriction occurs coincident with proliferation and results in the formation of cell type-specific precursor cells, which eventually exit the cell cycle to acquire a permanent postmitotic state. While still poorly understood for neuronal differentiation, recent studies provide insight in the coordination between terminal differentiation and cell cycle arrest during postembryonic muscle formation in *C. elegans*.

**Neuronal differentiation:** In the *C. elegans* nervous system, the generation of specific neuronal cell types follows acquisition of the postmitotic state. The “terminal selector”


transcription factors that induce a specific neural identity continue to be needed to maintain this identity, but not to retain the postmitotic state (Deneris and Hobert 2014). Some of these terminal selectors are first expressed in postmitotic cells, and it is possible that the combinatorial transcription factors that induce pan-neuronal properties are responsible for cell cycle withdrawal. The redundancy among such factors, which include the HOX genes and TCF/ $\beta$ -catenin combinations, may explain the lack of reported mutations that allow continued cell division at the expense of neuronal differentiation. Possible exceptions are *egl-44* and *egl-46* mutants, in which Q neuroblast descendants undergo an extra terminal division (Feng *et al.* 2013). Moreover, certain mutations result in neuronal lineage reiteration. Most prominently, in *unc-86*<sup>POU</sup> mutants, several neuroblasts adopt a repetitive stem cell-like division pattern. This phenotype appears to reflect a defect in cell identity rather than a specific defect in cell cycle exit.

**Muscle formation:** Mammalian skeletal muscle development involves a cascade of myogenic regulatory factors (MRFs) that induce muscle differentiation as well as cell cycle arrest. *C. elegans* and other invertebrates express a single MyoD-related MRF to induce contractile muscle formation (Krause *et al.* 1990). Mutation of *C. elegans* *hlh-1*<sup>MyoD</sup> does not prevent striated muscle formation, as additional factors, in particular *unc-120*<sup>SRF</sup> and *hnd-1*<sup>HAND</sup>, act partially redundantly (Fukushige *et al.* 2006). In addition to embryonic muscle, muscle cells are formed during larval development in the lineage of the mesoblast (M) (Figure 2C). This includes 16 striated body wall muscles, as well as 16 nonstriated muscles required for egg laying. HLH-1<sup>MyoD</sup> is present specifically in the striated muscle and appears well before the terminal division (Harfe *et al.* 1998). Remarkably, loss of *hlh-1*<sup>MyoD</sup> results in the formation of extra nonstriated uterine and vulval muscles, which normally do not express HLH-1. This discrepancy was traced to the formation of extra precursor cells of the egg-laying muscle (known as “sex myoblasts”) at the expense of differentiated body wall muscle (Harfe *et al.* 1998). Recent observations indicate that mesoblast-specific *hlh-1* knockout interferes with the proper cell cycle arrest of body wall muscle precursor cells (M. Godfrey and S. van den Heuvel, unpublished data). Thus, HLH-1<sup>MyoD</sup> contributes to both muscle cell differentiation and cell cycle arrest.

### **Coordinating cell cycle exit with differentiation: SWI/SNF chromatin remodelers**

Studies of the *C. elegans* mesoblast lineage have highlighted the role of chromatin regulators in the coordination between cell cycle arrest and differentiation. Specifically, SWI/SNF chromatin remodeling complexes were found to contribute critically to the transition from proliferating precursor cells to postmitotic muscle cells (Ruijtenberg and van den Heuvel 2015). Analyzing body wall muscle and egg-laying muscle formation in M lineage-specific mutants revealed that the cell cycle arrest during differentiation uses highly redundant





**Figure 8** Model for cell cycle arrest during terminal differentiation of body wall muscle precursor cells. APC/C<sup>FZR-1</sup>- and SCF<sup>LIN-23</sup>-mediated ubiquitin-dependent protein degradation, LIN-35<sup>Rb</sup>-mediated transcriptional repression, and association of CDK-inhibitory proteins with CDK/cyclin complexes promotes cell-cycle arrest. During terminal differentiation of muscle precursor cells, SWI/SNF complexes in cooperation with lineage-specific transcription factors provide an additional level of control. The different G1/S inhibitors and SWI/SNF chromatin remodelers cooperate by providing alternative levels of control over the basic cell cycle regulators, with each level antagonizing CDK-4/CYD-1 cyclin D and/or CDK-2/CYE-1 cyclin E kinase activity. Together, these regulators provide a highly robust control network for cell cycle exit.

controls. Double mutation of *lin-35* and *fzr-1*, which deregulates cell divisions in some other lineages, did not alter the M lineage division pattern. A few extra M descendants were formed following RNAi of *cki-1*, *lin-23*, or *cul-1*, as expected, and also upon RNAi of genes encoding SWI/SNF core subunits. However, testing multiple mutant and RNAi combinations revealed that only simultaneous inhibition of SWI/SNF and G1/S inhibitor functions severely disrupts cell cycle arrest and terminal differentiation, resulting in the tumorous overproliferation of M daughter cells (Ruijtenberg and van den Heuvel 2015). These data show that canonical G1/S inhibitors and SWI/SNF chromatin remodeling complexes substantially overlap in promoting cell cycle exit and the differentiation of muscle precursor cells.

Whole-worm chromatin immunoprecipitation (ChIP) experiments have revealed thousands of genomic SWI/SNF-binding sites (Riedel *et al.* 2013). This includes association with the promoter/enhancer regions of nearly all negative regulators of cell division, as well as the positive regulators *cdk-4* and *cye-1*. Similarly, analysis of ChIP data demonstrated HLH-1<sup>MYOD</sup> binding at overlapping locations in the *cki-1*, *cdk-4*, and *cye-1* gene regulatory regions (Ruijtenberg and van den Heuvel 2015; ChIP data from the Krause and Snyder laboratories). Combined with single-molecule FISH studies of gene expression, the data support a model in which HLH-1<sup>MYOD</sup> acts in concert with SWI/SNF chromatin remodelers to induce expression of muscle-specific genes as well as cell cycle inhibitors. At the same time, SWI/SNF complexes directly or indirectly appear to repress *cdk-4* and *cye-1* (Figure 8).

In *Drosophila* and mammals, SWI/SNF chromatin remodelers antagonize Polycomb repressor complexes (PRCs) and act as important tumor suppressor genes (Kadoch and Crabtree 2015). Interestingly, downregulation of *mes-2*, related to the PRC2 EZH2 methyltransferase, strongly suppressed the extra divisions of muscle precursor cells in lineage-specific

*swn-1* mutants (Ruijtenberg and van den Heuvel 2015). Thus, it is conceivable that loss of SWI/SNF leads to the recruitment of PRC2-related H3K27 methylation complexes that repress the transcription of genes that are normally activated during terminal differentiation. Recent results indicate that the SWI/SNF complex is highly dosage-dependent, with complete loss-of-function preventing, rather than promoting, cell proliferation (A. van der Vaart and van den Heuvel, unpublished data). These data resemble those obtained in human cancer studies (Kadoch and Crabtree 2015), indicating that the studies of SWI/SNF remodelers in cell cycle arrest during the differentiation of *C. elegans* muscle cells offers insight into the frequent mutation of SWI/SNF genes in human cancers.

#### Combining cell proliferation and differentiation through asymmetric cell division

In *C. elegans*, many cell divisions create an anterior and a posterior daughter cell, which differ in nuclear levels of the POP-1<sup>TCF/LEF</sup> transcription factor and cell fate (Park and Priess 2003). Such asymmetric cell divisions may segregate the potential to proliferate and commitment to differentiate to different daughter cells. A Wnt/ $\beta$ -catenin asymmetry pathway controls the nuclear level of POP-1, and thereby determines whether POP-1 acts as a transcriptional repressor with UNC-37<sup>Groucho</sup> or as a transcriptional activator with SYS-1 <sup>$\beta$ -catenin</sup>. Thus, POP-1 is an important determinant of cell fate, which shows complex interactions with the cell cycle machinery. For instance, in the somatic gonad, Cyclin D and other G1 regulators appear to regulate axis formation and sex determination, by acting as upstream regulators of the POP-1 and FKH-6 transcription factors, respectively (Tilman and Kimble 2005).

The Wnt/ $\beta$ -catenin asymmetry pathway also controls an asymmetry in CKI-1 and CYE-1 levels during specific cell divisions in the somatic gonad (Fujita *et al.* 2007). The second

asymmetric divisions of the Z1.a and Z4.p gonadal precursor cells each form a DTC. During the asymmetric division, the DTC daughter receives a higher level of *CKI-1*, while the sister cell maintains more *CYE-1*. Reducing *CYE-1* or increasing *CKI-1* in the sister cell causes it to adopt a DTC fate, instead of its normal quiescence followed by continued mitotic division in the L3 stage. Based on these observations, maintenance of *CDK-2/cyclin E* activity at a low level has been proposed to prevent the differentiation of the sister cells (Fujita *et al.* 2007).

Stem cell-like seam cells in the epidermis undergo asymmetric cell divisions during each larval stage (Figure 2, A and B). These asymmetric divisions create a new seam cell, and a cell that either fuses with the general epidermis (*hyp7*) or forms neurons (Sulston and Horvitz 1977). Around the L1/L2 molt, six of the seam cells (V1-4p-V6p and Tp) also undergo a symmetric cell division that expands the seam cell number (Figure 2A). The extra divisions substantially depend on a Runx family transcription factor, *RNT-1*, in association with the cofactors *BRO-1*<sup>CBF $\beta$</sup>  and *UNC-37*<sup>Groucho</sup> (Nimmo *et al.* 2005; Kagoshima *et al.* 2007; Xia *et al.* 2007). Based on genetic interactions, this repressor complex has been proposed to antagonize the *cki-1*, *lin-35*, and *fzr-1* cell cycle inhibitor genes. This regulation, in combination with Wnt/ $\beta$  catenin signaling and the heterochronic network, establishes the proper cell division–differentiation patterns in the seam cell lineages.

## Perspectives

The use of *C. elegans* as a model system has yielded considerable insight into how this simple animal regulates its cell divisions in the context of its physiology and development. The analysis of *C. elegans* has broken new ground in areas that include the regulation of stem cell divisions in the hermaphrodite germline and asymmetric cell divisions in the early embryo. Cell cycle regulatory functions have been described for many different CRL/SCF ubiquitin ligases, and links have been established between transcriptional regulators and cell cycle genes in the decision between proliferation and terminal differentiation. Moreover, the studies of cell cycle regulation in *C. elegans* have uncovered differences between tissues and developmental stages in the regulation of DNA replication and cell division.

Much research has been focused on how external conditions are associated with systemic factors, such as insulin or other hormonal signals, in the regulation of cell division. External signals that come from the environment impact how *C. elegans* responds in its boom and bust life cycle. In response to replete food conditions without crowding, the animal rapidly develops and reproduces. In the absence of food, animals undergo regulated quiescence in the L1 and dauer larval stages, and as adults reorganize the germline to conserve GSCs while shrinking the remainder of the germline to generate limited numbers of oocytes. Thus, cell cycle regulation in *C. elegans* is more tightly linked to its surroundings

than, for example, the typical cell within a mammalian tissue.

Studies of the early embryo have yielded many insights into how the cell cycle is coupled to the regulation of cell polarity and positioning of the mitotic spindle, the regulation of spindle forces, and cell division asymmetry. The early embryo has also provided extensive insight into the control of centrosome duplication. The biology of the spindle and centrosomes were not covered extensively in this review, and are instead extensively described in another WormBook chapter on cell division and other reviews (Rose and Gönczy 2014).

When considering the field as a whole, what remains less well understood is the question how cell proliferation is regulated to produce the invariant cell lineage. Cell cycle regulators that are required to allow the proper cell lineage have been identified, and substantial redundancies among such regulators have been detected. The inactivation of specific negative regulators interferes with the normal program of cell cycle exit resulting in supernumerary cell divisions, and the inactivation of positive regulators prevents normal cell divisions. However, what is still largely missing is an understanding of how these cell cycle regulators are controlled within normal development to produce the stereotypical cell division pattern where, for example, a vulval lineage will undergo three rounds of cell division and then exit the cell cycle. Global regulators of these cell division patterns, such as heterochronic genes that control the timing of larval stage programs, have not yet been linked at the molecular level to the control of specific cell cycle regulators. Therefore, the promise of using *C. elegans* to understand how an invariant cell lineage is generated still remains largely an outstanding question.

In addition to the number and timing of cell divisions, the coordination between cell cycle arrest and terminal differentiation remains an important topic for future studies. Such studies will continue to benefit from the transparency of *C. elegans*, by allowing detailed observations in intact living animals. Many recent technical developments will help future analyses through advanced genome engineering, time-lapse fluorescence microscopy, and single-cell analysis methods. The possibilities for reverse genetics continue to expand, for instance with the addition of lineage-specific gene knockout and protein degradation strategies, and whole-genome sequencing has significantly accelerated conventional mutational analysis. Individual cells, proteins, and protein complexes can be followed *in vivo*, and can be purified efficiently from stage-synchronized animals. Many insights are expected to be obtained from genomic, transcriptomic, and proteomic analyses of individual cells, or pools of cells, extracted at well-defined developmental times. With the current knowledge of the central cell cycle regulators, the stage is now set for the connection of a large variety of cell cycle phenotypes to molecular pathways and the acquisition of a more complete understanding of the regulatory networks that control the cell cycle in animal development.

## Acknowledgments

We thank all of our colleagues for their support and inspiration over the years, and apologize for all of the good research that could not be included. S.v.d.H. thanks J. Teapal for help with the figures. This work was financed by a grant from the National Institutes of Health, National Institute of General Medical Sciences (R01 GM 074212) to E.T.K. and by the Netherlands Organization for Scientific Research (Nederlandse Organisatie voor Wetenschappelijk Onderzoek), in part through a ZonMW TOP project (91216058) to S.v.d.H.

## Literature Cited

- Alexander, J. L., and T. L. Orr-Weaver, 2016 Replication fork instability and the consequences of fork collisions from rereplication. *Genes Dev.* 30: 2241–2252. <https://doi.org/10.1101/gad.288142.116>
- Amini, R., E. Goupil, S. Labella, M. Zetka, A. S. Maddox *et al.*, 2014 *C. elegans* anillin proteins regulate intercellular bridge stability and germline syncytial organization. *J. Cell Biol.* 206: 129–143 [corrigenda: *J. Cell Biol.* 209: 467 (2015)]. <https://doi.org/10.1083/jcb.201310117>
- Angelo, G., and M. R. Van Gilst, 2009 Starvation protects germline stem cells and extends reproductive longevity in *C. elegans*. *Science* 326: 954–958. <https://doi.org/10.1126/science.1178343>
- Antebi, A., 2013 Steroid regulation of *C. elegans* diapause, developmental timing, and longevity. *Curr. Top. Dev. Biol.* 105: 181–212. <https://doi.org/10.1016/B978-0-12-396968-2.00007-5>
- Ashcroft, N., and A. Golden, 2002 CDC-25.1 regulates germline proliferation in *Caenorhabditis elegans*. *Genesis* 33: 1–7. <https://doi.org/10.1002/gene.10083>
- Ashcroft, N. R., M. Srayko, M. E. Kosinski, P. E. Mains, and A. Golden, 1999 RNA-mediated interference of a *cdc25* homolog in *Caenorhabditis elegans* results in defects in the embryonic cortical membrane, meiosis, and mitosis. *Dev. Biol.* 206: 15–32. <https://doi.org/10.1006/dbio.1998.9135>
- Austin, M. U., W. S. Liao, K. Balamurugan, B. Ashokkumar, H. M. Said *et al.*, 2010 Knockout of the folate transporter *folt-1* causes germline and somatic defects in *C. elegans*. *BMC Dev. Biol.* 10: 46. <https://doi.org/10.1186/1471-213X-10-46>
- Balachandran, R. S., C. S. Heighington, N. G. Starostina, J. W. Anderson, D. L. Owen *et al.*, 2016 The ubiquitin ligase CRL2ZYG11 targets cyclin B1 for degradation in a conserved pathway that facilitates mitotic slippage. *J. Cell Biol.* 215: 151–166. <https://doi.org/10.1083/jcb.201601083>
- Balamurugan, K., B. Ashokkumar, M. Moussaif, J. Y. Sze, and H. M. Said, 2007 Cloning and functional characterization of a folate transporter from the nematode *Caenorhabditis elegans*. *Am. J. Physiol. Cell Physiol.* 293: C670–C681. <https://doi.org/10.1152/ajpcell.00516.2006>
- Bashir, T., N. V. Dorrello, V. Amador, D. Guardavaccaro, and M. Pagano, 2004 Control of the SCF(Skp2-Cks1) ubiquitin ligase by the APC/C(Cdh1) ubiquitin ligase. *Nature* 428: 190–193. <https://doi.org/10.1038/nature02330>
- Baugh, L. R., 2013 To grow or not to grow: nutritional control of development during *Caenorhabditis elegans* L1 arrest. *Genetics* 194: 539–555. <https://doi.org/10.1534/genetics.113.150847>
- Baugh, L. R., and P. W. Sternberg, 2006 DAF-16/FOXO regulates transcription of *cki-1*/Cip/Kip and repression of *lin-4* during *C. elegans* L1 arrest. *Curr. Biol.* 16: 780–785. <https://doi.org/10.1016/j.cub.2006.03.021>
- Baugh, L. R., N. Kurhanewicz, and P. W. Sternberg, 2011 Sensitive and precise quantification of insulin-like mRNA expression in *Caenorhabditis elegans*. *PLoS One* 6: e18086. <https://doi.org/10.1371/journal.pone.0018086>
- Bertrand, V., and O. Hobert, 2010 Lineage programming: navigating through transient regulatory states via binary decisions. *Curr. Opin. Genet. Dev.* 20: 362–368. <https://doi.org/10.1016/j.gde.2010.04.010>
- Bethke, A., N. Fielenbach, Z. Wang, D. J. Mangelsdorf, and A. Antebi, 2009 Nuclear hormone receptor regulation of micro-RNAs controls developmental progression. *Science* 324: 95–98. <https://doi.org/10.1126/science.1164899>
- Biedermann, B., J. Wright, M. Senften, I. Kalchauer, G. Sarathy *et al.*, 2009 Translational repression of cyclin E prevents precocious mitosis and embryonic gene activation during *C. elegans* meiosis. *Dev. Cell* 17: 355–364. <https://doi.org/10.1016/j.devcel.2009.08.003>
- Boxem, M., and S. van den Heuvel, 2001 *lin-35* Rb and *cki-1* Cip/Kip cooperate in developmental regulation of G1 progression in *C. elegans*. *Development* 128: 4349–4359.
- Boxem, M., and S. van den Heuvel, 2002 *C. elegans* class B synthetic multivulva genes act in G1 regulation. *Curr. Biol.* 12: 906–911. [https://doi.org/10.1016/S0960-9822\(02\)00844-8](https://doi.org/10.1016/S0960-9822(02)00844-8)
- Boxem, M., D. G. Srinivasan, and S. van den Heuvel, 1999 The *Caenorhabditis elegans* gene *ncc-1* encodes a *cdc2*-related kinase required for M phase in meiotic and mitotic cell divisions, but not for S phase. *Development* 126: 2227–2239.
- Brauchle, M., K. Baumer, and P. Gönczy, 2003 Differential activation of the DNA replication checkpoint contributes to asynchrony of cell division in *C. elegans* embryos. *Curr. Biol.* 13: 819–827. [https://doi.org/10.1016/S0960-9822\(03\)00295-1](https://doi.org/10.1016/S0960-9822(03)00295-1)
- Brenner, J. L., and T. Schedl, 2016 Germline stem cell differentiation entails regional control of cell fate regulator GLD-1 in *Caenorhabditis elegans*. *Genetics* 202: 1085–1103. <https://doi.org/10.1534/genetics.115.185678>
- Brodigan, T. M., J. Liu, M. Park, E. T. Kipreos, and M. Krause, 2003 Cyclin E expression during development in *Caenorhabditis elegans*. *Dev. Biol.* 254: 102–115. [https://doi.org/10.1016/S0012-1606\(02\)00032-5](https://doi.org/10.1016/S0012-1606(02)00032-5)
- Buck, S. H., D. Chiu, and R. M. Saito, 2009 The cyclin-dependent kinase inhibitors, *cki-1* and *cki-2*, act in overlapping but distinct pathways to control cell cycle quiescence during *C. elegans* development. *Cell Cycle* 8: 2613–2620. <https://doi.org/10.4161/cc.8.16.9354>
- Budirahardja, Y., and P. Gönczy, 2008 PLK-1 asymmetry contributes to asynchronous cell division of *C. elegans* embryos. *Development* 135: 1303–1313. <https://doi.org/10.1242/dev.019075>
- Burrows, A. E., B. K. Scurman, M. E. Kosinski, C. T. Richie, P. L. Sadler *et al.*, 2006 The *C. elegans* Myt1 ortholog is required for the proper timing of oocyte maturation. *Development* 133: 697–709. <https://doi.org/10.1242/dev.02241>
- Buttitta, L. A., A. J. Katzaroff, C. L. Perez, A. de la Cruz, and B. A. Edgar, 2007 A double-assurance mechanism controls cell cycle exit upon terminal differentiation in *Drosophila*. *Dev. Cell* 12: 631–643. <https://doi.org/10.1016/j.devcel.2007.02.020>
- Byrd, D. T., K. Knobel, K. Affeldt, S. L. Crittenden, and J. Kimble, 2014 A DTC niche plexus surrounds the germline stem cell pool in *Caenorhabditis elegans*. *PLoS One* 9: e88372. <https://doi.org/10.1371/journal.pone.0088372>
- Castro, P. V., S. Khare, B. D. Young, and S. G. Clarke, 2012 *Caenorhabditis elegans* battling starvation stress: low levels of ethanol prolong lifespan in L1 larvae. *PLoS One* 7: e29984. <https://doi.org/10.1371/journal.pone.0029984>
- Ceol, C. J., and H. R. Horvitz, 2001 *dpl-1* DP and *efl-1* E2F act with *lin-35* Rb to antagonize Ras signaling in *C. elegans* vulval development. *Mol. Cell* 7: 461–473. [https://doi.org/10.1016/S1097-2765\(01\)00194-0](https://doi.org/10.1016/S1097-2765(01)00194-0)
- Chalfie, M., Y. Tu, G. Euskirchen, W. W. Ward, and D. C. Prasher, 1994 Green fluorescent protein as a marker for gene

- expression. *Science* 263: 802–805. <https://doi.org/10.1126/science.8303295>
- Chase, D., C. Serafinas, N. Ashcroft, M. Kosinski, D. Longo *et al.*, 2000 The polo-like kinase PLK-1 is required for nuclear envelope breakdown and the completion of meiosis in *Caenorhabditis elegans*. *Genesis* 26: 26–41. [https://doi.org/10.1002/\(SICI\)1526-968X\(200001\)26:1<26::AID-GENE6>3.0.CO;2-O](https://doi.org/10.1002/(SICI)1526-968X(200001)26:1<26::AID-GENE6>3.0.CO;2-O)
- Chaudhari, S. N., M. Mukherjee, A. S. Vagasi, G. Bi, M. M. Rahman *et al.*, 2016 Bacterial folates provide an exogenous signal for *C. elegans* germline stem cell proliferation. *Dev. Cell* 38: 33–46. <https://doi.org/10.1016/j.devcel.2016.06.013>
- Cheng, K. C.-C., R. Klancer, A. Singson, and G. Seydoux, 2009 Regulation of MBK-2/DYRK by CDK-1 and the pseudophosphatases EGG-4 and EGG-5 during the oocyte-to-embryo transition. *Cell* 139: 560–572. <https://doi.org/10.1016/j.cell.2009.08.047>
- Chi, W., and V. Reinke, 2006 Promotion of oogenesis and embryogenesis in the *C. elegans* gonad by EFL-1/DPL-1 (E2F) does not require LIN-35 (pRB). *Development* 133: 3147–3157. <https://doi.org/10.1242/dev.02490>
- Cinquin, O., S. L. Crittenden, D. E. Morgan, and J. Kimble, 2010 Progression from a stem cell-like state to early differentiation in the *C. elegans* germ line. *Proc. Natl. Acad. Sci. USA* 107: 2048–2053. <https://doi.org/10.1073/pnas.0912704107>
- Clayton, J. E., S. J. L. van den Heuvel, and R. M. Saito, 2008 Transcriptional control of cell-cycle quiescence during *C. elegans* development. *Dev. Biol.* 313: 603–613. <https://doi.org/10.1016/j.ydbio.2007.10.051>
- Cowan, C. R., and A. A. Hyman, 2006 Cyclin E-Cdk2 temporally regulates centrosome assembly and establishment of polarity in *Caenorhabditis elegans* embryos. *Nat. Cell Biol.* 8: 1441–1447. <https://doi.org/10.1038/ncb1511>
- Crittenden, S. L., D. S. Bernstein, J. L. Bachorik, B. E. Thompson, M. Gallegos *et al.*, 2002 A conserved RNA-binding protein controls germline stem cells in *Caenorhabditis elegans*. *Nature* 417: 660–663. <https://doi.org/10.1038/nature754>
- Crittenden, S. L., K. A. Leonhard, D. T. Byrd, and J. Kimble, 2006 Cellular analyses of the mitotic region in the *Caenorhabditis elegans* adult germ line. *Mol. Biol. Cell* 17: 3051–3061. <https://doi.org/10.1091/mbc.e06-03-0170>
- Cui, M., J. Chen, T. R. Myers, B. J. Hwang, P. W. Sternberg *et al.*, 2006 SynMuv genes redundantly inhibit *lin-3*/EGF expression to prevent inappropriate vulval induction in *C. elegans*. *Dev. Cell* 10: 667–672. <https://doi.org/10.1016/j.devcel.2006.04.001>
- Dalfó, D., D. Michaelson, and E. J. Hubbard, 2012 Sensory regulation of the *C. elegans* germline through TGF- $\beta$ -dependent signaling in the niche. *Curr. Biol.* 22: 712–719. <https://doi.org/10.1016/j.cub.2012.02.064>
- Davis, E. S., L. Wille, B. A. Chestnut, P. L. Sadler, D. C. Shakes *et al.*, 2002 Multiple subunits of the *Caenorhabditis elegans* anaphase-promoting complex are required for chromosome segregation during meiosis I. *Genetics* 160: 805–813.
- Deneris, E. S., and O. Hobert, 2014 Maintenance of postmitotic neuronal cell identity. *Nat. Neurosci.* 17: 899–907. <https://doi.org/10.1038/nn.3731>
- DePamphilis, M. L., J. J. Blow, S. Ghosh, T. Saha, K. Noguchi *et al.*, 2006 Regulating the licensing of DNA replication origins in metazoa. *Curr. Opin. Cell Biol.* 18: 231–239. <https://doi.org/10.1016/j.ceb.2006.04.001>
- Deppe, U., E. Schierenberg, T. Cole, C. Krieg, D. Schmitt *et al.*, 1978 Cell lineages of the embryo of the nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 75: 376–380. <https://doi.org/10.1073/pnas.75.1.376>
- Detwiler, M. R., M. Reuben, X. Li, E. Rogers, and R. Lin, 2001 Two zinc finger proteins, OMA-1 and OMA-2, are redundantly required for oocyte maturation in *C. elegans*. *Dev. Cell* 1: 187–199. [https://doi.org/10.1016/S1534-5807\(01\)00026-0](https://doi.org/10.1016/S1534-5807(01)00026-0)
- Deyter, G. M. R., T. Furuta, Y. Kurasawa, and J. M. Schumacher, 2010 *Caenorhabditis elegans* cyclin B3 is required for multiple mitotic processes including alleviation of a spindle checkpoint-dependent block in anaphase chromosome segregation. *PLoS Genet.* 6: e1001218. <https://doi.org/10.1371/journal.pgen.1001218>
- Dickinson, D. J., and B. Goldstein, 2016 CRISPR-based methods for *Caenorhabditis elegans* genome engineering. *Genetics* 202: 885–901.
- Diehl, J. A., M. Cheng, M. F. Roussel, and C. J. Sherr, 1998 Glycogen synthase kinase-3 $\beta$  regulates cyclin D1 proteolysis and subcellular localization. *Genes Dev.* 12: 3499–3511. <https://doi.org/10.1101/gad.12.22.3499>
- Dimova, D. K., O. Stevaux, M. V. Frolov, and N. J. Dyson, 2003 Cell cycle-dependent and cell cycle-independent control of transcription by the *Drosophila* E2F/RB pathway. *Genes Dev.* 17: 2308–2320. <https://doi.org/10.1101/gad.1116703>
- Dorée, M., and T. Hunt, 2002 From Cdc2 to Cdk1: when did the cell cycle kinase join its cyclin partner? *J. Cell Sci.* 115: 2461–2464.
- Dunphy, W. G., L. Brizuela, D. Beach, and J. Newport, 1988 The Xenopus cdc2 protein is a component of MPF, a cytoplasmic regulator of mitosis. *Cell* 54: 423–431. [https://doi.org/10.1016/0092-8674\(88\)90205-X](https://doi.org/10.1016/0092-8674(88)90205-X)
- Edgar, L. G., and J. D. McGhee, 1988 DNA synthesis and the control of embryonic gene expression in *C. elegans*. *Cell* 53: 589–599. [https://doi.org/10.1016/0092-8674\(88\)90575-2](https://doi.org/10.1016/0092-8674(88)90575-2)
- Encalada, S. E., P. R. Martin, J. B. Phillips, R. Lyczak, D. R. Hamill *et al.*, 2000 DNA replication defects delay cell division and disrupt cell polarity in early *Caenorhabditis elegans* embryos. *Dev. Biol.* 228: 225–238. <https://doi.org/10.1006/dbio.2000.9965>
- Encalada, S. E., J. Willis, R. Lyczak, and B. Bowerman, 2005 A spindle checkpoint functions during mitosis in the early *Caenorhabditis elegans* embryo. *Mol. Biol. Cell* 16: 1056–1070. <https://doi.org/10.1091/mbc.e04-08-0712>
- Espeut, J., P. Lara-Gonzalez, M. Sassine, A. K. Shiau, A. Desai *et al.*, 2015 Natural loss of Mps1 kinase in nematodes uncovers a role for Polo-like Kinase 1 in spindle checkpoint initiation. *Cell Rep.* 12: 58–65. <https://doi.org/10.1016/j.celrep.2015.05.039>
- Euling, S., and V. Ambros, 1996 Heterochronic genes control cell cycle progress and developmental competence of *C. elegans* vulva precursor cells. *Cell* 84: 667–676. [https://doi.org/10.1016/S0092-8674\(00\)81045-4](https://doi.org/10.1016/S0092-8674(00)81045-4)
- Fay, D. S., and M. Han, 2000 Mutations in *cye-1*, a *Caenorhabditis elegans* cyclin E homolog, reveal coordination between cell-cycle control and vulval development. *Development* 127: 4049–4060.
- Fay, D. S., S. Keenan, and M. Han, 2002 *fzr-1* and *lin-35*/Rb function redundantly to control cell proliferation in *C. elegans* as revealed by a nonbiased synthetic screen. *Genes Dev.* 16: 503–517. <https://doi.org/10.1101/gad.952302>
- Feng, G., P. Yi, Y. Yang, Y. Chai, D. Tian *et al.*, 2013 Developmental stage-dependent transcriptional regulatory pathways control neuroblast lineage progression. *Development* 140: 3838–3847. <https://doi.org/10.1242/dev.098723>
- Feng, H., W. Zhong, G. Punkosdy, S. Gu, L. Zhou *et al.*, 1999 CUL-2 is required for the G1-to-S-phase transition and mitotic chromosome condensation in *Caenorhabditis elegans*. *Nat. Cell Biol.* 1: 486–492. <https://doi.org/10.1038/70272>
- Fire, A., S. Xu, M. K. Montgomery, S. A. Kostas, S. E. Driver *et al.*, 1998 Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391: 806–811. <https://doi.org/10.1038/35888>
- Fisher, R. P., and D. O. Morgan, 1994 A novel cyclin associates with MO15/CDK7 to form the CDK-activating kinase. *Cell* 78: 713–724. [https://doi.org/10.1016/0092-8674\(94\)90535-5](https://doi.org/10.1016/0092-8674(94)90535-5)

- Flemming, A. J., Z. Z. Shen, A. Cunha, S. W. Emmons, and A. M. Leroi, 2000 Somatic polyploidization and cellular proliferation drive body size evolution in nematodes. *Proc. Natl. Acad. Sci. USA* 97: 5285–5290. <https://doi.org/10.1073/pnas.97.10.5285>
- Fox, P. M., and T. Schedl, 2015 Analysis of germline stem cell differentiation following loss of GLP-1 Notch activity in *Caenorhabditis elegans*. *Genetics* 201: 167–184. <https://doi.org/10.1534/genetics.115.178061>
- Fox, P. M., V. E. Vought, M. Hanazawa, M. H. Lee, E. M. Maine *et al.*, 2011 Cyclin E and CDK-2 regulate proliferative cell fate and cell cycle progression in the *C. elegans* germline. *Development* 138: 2223–2234. <https://doi.org/10.1242/dev.059535>
- Fujita, M., H. Takeshita, and H. Sawa, 2007 Cyclin E and CDK2 repress the terminal differentiation of quiescent cells after asymmetric division in *C. elegans*. *PLoS One* 2: e407. <https://doi.org/10.1371/journal.pone.0000407>
- Fukushige, T., T. M. Brodigan, L. A. Schriefer, R. H. Waterston, and M. Krause, 2006 Defining the transcriptional redundancy of early bodywall muscle development in *C. elegans*: evidence for a unified theory of animal muscle development. *Genes Dev.* 20: 3395–3406. <https://doi.org/10.1101/gad.1481706>
- Fukuyama, M., S. B. Gendreau, W. B. Derry, and J. H. Rothman, 2003 Essential embryonic roles of the CKI-1 cyclin-dependent kinase inhibitor in cell-cycle exit and morphogenesis in *C. elegans*. *Dev. Biol.* 260: 273–286. [https://doi.org/10.1016/S0012-1606\(03\)00239-2](https://doi.org/10.1016/S0012-1606(03)00239-2)
- Fukuyama, M., A. E. Rougvie, and J. H. Rothman, 2006 *C. elegans* DAF-18/PTEN mediates nutrient-dependent arrest of cell cycle and growth in the germline. *Curr. Biol.* 16: 773–779. <https://doi.org/10.1016/j.cub.2006.02.073>
- Galli, M., and D. O. Morgan, 2016 Cell size determines the strength of the spindle assembly checkpoint during embryonic development. *Dev. Cell* 36: 344–352. <https://doi.org/10.1016/j.devcel.2016.01.003>
- Gautier, J., C. Norbury, M. Lohka, P. Nurse, and J. Maller, 1988 Purified maturation-promoting factor contains the product of a *Xenopus* homolog of the fission yeast cell cycle control gene *cdc2+*. *Cell* 54: 433–439. [https://doi.org/10.1016/0092-8674\(88\)90206-1](https://doi.org/10.1016/0092-8674(88)90206-1)
- Gibert, M. A., J. Starck, and B. Beguet, 1984 Role of the gonad cytoplasmic core during oogenesis of the nematode *Caenorhabditis elegans*. *Biol. Cell* 50: 77–85. <https://doi.org/10.1111/j.1768-322X.1984.tb00254.x>
- Goetsch, P. D., J. M. Garrigues, and S. Strome, 2017 Loss of the *Caenorhabditis elegans* pocket protein LIN-35 reveals MuvB's innate function as the repressor of DREAM target genes. *PLoS Genet.* 13: e1007088. <https://doi.org/10.1371/journal.pgen.1007088>
- Golden, A., P. L. Sadler, M. R. Wallenfang, J. M. Schumacher, D. R. Hamill *et al.*, 2000 Metaphase to anaphase (mat) transition-defective mutants in *Caenorhabditis elegans*. *J. Cell Biol.* 151: 1469–1482. <https://doi.org/10.1083/jcb.151.7.1469>
- Grallert, A., E. Boke, A. Hagting, B. Hodgson, Y. Connolly *et al.*, 2015 A PP1–PP2A phosphatase relay controls mitotic progression. *Nature* 517: 94–98. <https://doi.org/10.1038/nature14019>
- Griffin, E. E., D. J. Odde, and G. Seydoux, 2011 Regulation of the MEX-5 gradient by a spatially segregated kinase/phosphatase cycle. *Cell* 146: 955–968. <https://doi.org/10.1016/j.cell.2011.08.012>
- Gumienny, T. L., and C. Savage-Dunn, 2013 TGF- $\beta$  signaling in *C. elegans* (July 10, 2013), *WormBook*, ed. The *C. elegans* Research Community *WormBook*, doi/10.1895/wormbook.1.22.2, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.22.2>
- Guo, S., and K. J. Kemphues, 1995 *par-1*, a gene required for establishing polarity in *C. elegans* embryos, encodes a putative Ser/Thr kinase that is asymmetrically distributed. *Cell* 81: 611–620. [https://doi.org/10.1016/0092-8674\(95\)90082-9](https://doi.org/10.1016/0092-8674(95)90082-9)
- Guven-Ozkan, T., Y. Nishi, S. M. Robertson, and R. Lin, 2008 Global transcriptional repression in *C. elegans* germline precursors by regulated sequestration of TAF-4. *Cell* 135: 149–160. <https://doi.org/10.1016/j.cell.2008.07.040>
- Hall, D. H., and Z. F. Altun, 2008 *C. elegans Atlas*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Hansen, D., and T. Schedl, 2013 Stem cell proliferation vs. meiotic fate decision in *Caenorhabditis elegans*. *Adv. Exp. Med. Biol.* 757: 71–99. [https://doi.org/10.1007/978-1-4614-4015-4\\_4](https://doi.org/10.1007/978-1-4614-4015-4_4)
- Harfe, B. D., C. S. Branda, M. Krause, M. J. Stern, and A. Fire, 1998 MyoD and the specification of muscle and non-muscle fates during postembryonic development of the *C. elegans* mesoderm. *Development* 125: 2479–2488.
- Harris, D. T., and H. R. Horvitz, 2011 MAB-10/NAB acts with LIN-29/EGR to regulate terminal differentiation and the transition from larva to adult in *C. elegans*. *Development* 138: 4051–4062. <https://doi.org/10.1242/dev.065417>
- Harrison, M. M., C. J. Ceol, X. Lu, and H. R. Horvitz, 2006 Some *C. elegans* class B synthetic multivulva proteins encode a conserved LIN-35 Rb-containing complex distinct from a NuRD-like complex. *Proc. Natl. Acad. Sci. USA* 103: 16782–16787. <https://doi.org/10.1073/pnas.0608461103>
- Hartwell, L. H., 2002 Nobel Lecture. *Yeast and cancer*. *Biosci. Rep.* 22: 373–394. <https://doi.org/10.1023/A:1020918107706>
- Havens, C. G., and J. C. Walter, 2011 Mechanism of CRL4(Cdt2), a PCNA-dependent E3 ubiquitin ligase. *Genes Dev.* 25: 1568–1582. <https://doi.org/10.1101/gad.2068611>
- Hebeisen, M., and R. Roy, 2008 CDC-25.1 stability is regulated by distinct domains to restrict cell division during embryogenesis in *C. elegans*. *Development* 135: 1259–1269. <https://doi.org/10.1242/dev.014969>
- Hedgcock, E. M., and J. G. White, 1985 Polyploid tissues in the nematode *Caenorhabditis elegans*. *Dev. Biol.* 107: 128–133. [https://doi.org/10.1016/0012-1606\(85\)90381-1](https://doi.org/10.1016/0012-1606(85)90381-1)
- Hong, Y., R. Roy, and V. Ambros, 1998 Developmental regulation of a cyclin-dependent kinase inhibitor controls postembryonic cell cycle progression in *Caenorhabditis elegans*. *Development* 125: 3585–3597.
- Horvitz, H. R., and J. E. Sulston, 1980 Isolation and genetic characterization of cell-lineage mutants of the nematode *Caenorhabditis elegans*. *Genetics* 96: 435–454.
- Jeffrey, P. D., A. A. Russo, K. Polyak, E. Gibbs, J. Hurwitz *et al.*, 1995 Mechanism of CDK activation revealed by the structure of a cyclinA-CDK2 complex. *Nature* 376: 313–320. <https://doi.org/10.1038/376313a0>
- Jeong, J., J. M. Verheyden, and J. Kimble, 2011 Cyclin E and Cdk2 control GLD-1, the mitosis/meiosis decision, and germline stem cells in *Caenorhabditis elegans*. *PLoS Genet.* 7: e1001348. <https://doi.org/10.1371/journal.pgen.1001348>
- Kadoch, C., and G. R. Crabtree, 2015 Mammalian SWI/SNF chromatin remodeling complexes and cancer: mechanistic insights gained from human genomics. *Sci. Adv.* 1: e1500447. <https://doi.org/10.1126/sciadv.1500447>
- Kagoshima, H., R. Nimmo, N. Saad, J. Tanaka, Y. Miwa *et al.*, 2007 The *C. elegans* CBF $\beta$  homologue BRO-1 interacts with the Runx factor, RNT-1, to promote stem cell proliferation and self-renewal. *Development* 134: 3905–3915. <https://doi.org/10.1242/dev.008276>
- Kalchauer, I., B. M. Farley, S. Pauli, S. P. Ryder, and R. Ciosk, 2011 FBF represses the Cip/Kip cell-cycle inhibitor CKI-2 to promote self-renewal of germline stem cells in *C. elegans*. *EMBO J.* 30: 3823–3829. <https://doi.org/10.1038/emboj.2011.263>
- Kaletta, T., H. Schnabel, and R. Schnabel, 1997 Binary specification of the embryonic lineage in *Caenorhabditis elegans*. *Nature* 390: 294–298. <https://doi.org/10.1038/36869>
- Kasuga, H., M. Fukuyama, A. Kitazawa, K. Kontani, and T. Katada, 2013 The microRNA miR-235 couples blast-cell quiescence to the nutritional state. *Nature* 497: 503–506. <https://doi.org/10.1038/nature12117>

- Kermi, C., E. Lo Furno, and D. Maiorano, 2017 Regulation of DNA replication in early embryonic cleavages. *Genes (Basel)* 8: 42. <https://doi.org/10.3390/genes8010042>
- Kershner, A. M., H. Shin, T. J. Hansen, and J. Kimble, 2014 Discovery of two GLP-1/Notch target genes that account for the role of GLP-1/Notch signaling in stem cell maintenance. *Proc. Natl. Acad. Sci. USA* 111: 3739–3744. <https://doi.org/10.1073/pnas.1401861111>
- Kim, J., H. Feng, and E. T. Kipreos, 2007 *C. elegans* CUL-4 prevents rereplication by promoting the nuclear export of CDC-6 via a CKI-1-dependent pathway. *Curr. Biol.* 17: 966–972. <https://doi.org/10.1016/j.cub.2007.04.055>
- Kim, J., I. Kawasaki, and Y.-H. Shim, 2010 *cdc-25.2*, a *C. elegans* ortholog of *cdc25*, is required to promote oocyte maturation. *J. Cell Sci.* 123: 993–1000. <https://doi.org/10.1242/jcs.060442>
- Kim, S., C. Spike, and D. Greenstein, 2013 Control of oocyte growth and meiotic maturation in *Caenorhabditis elegans*. *Adv. Exp. Med. Biol.* 757: 277–320. [https://doi.org/10.1007/978-1-4614-4015-4\\_10](https://doi.org/10.1007/978-1-4614-4015-4_10)
- Kim, Y., and E. T. Kipreos, 2007 The *Caenorhabditis elegans* replication licensing factor CDT-1 is targeted for degradation by the CUL-4/DDB-1 complex. *Mol. Cell. Biol.* 27: 1394–1406. <https://doi.org/10.1128/MCB.00736-06>
- Kim, Y., N. G. Starostina, and E. T. Kipreos, 2008 The CRL4<sup>Cdt2</sup> ubiquitin ligase targets the degradation of p21<sup>Cip1</sup> to control replication licensing. *Genes Dev.* 22: 2507–2519. <https://doi.org/10.1101/gad.1703708>
- Kimble, J., and S. L. Crittenden, 2005 Germline proliferation and its control (August 15, 2005), *WormBook*, ed. The *C. elegans* Research Community WormBook, doi/10.1895/wormbook.1.13.1, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.13.1>
- Kimble, J., and D. Hirsh, 1979 The postembryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. *Dev. Biol.* 70: 396–417. [https://doi.org/10.1016/0012-1606\(79\)90035-6](https://doi.org/10.1016/0012-1606(79)90035-6)
- Kimble, J., and H. Seidel, 2014 *C. elegans* germline stem cells and their niche. *StemBook* [Internet]. *Stem Book*. Available at: <http://dx.doi.org/10.3824/stembook.1.95.1>.
- Kipreos, E. T., 2005 *C. elegans* cell cycles: invariance and stem cell divisions. *Nat. Rev. Mol. Cell Biol.* 6: 766–776. <https://doi.org/10.1038/nrm1738>
- Kipreos, E. T., L. E. Lander, J. P. Wing, W. W. He, and E. M. Hedgecock, 1996 *cul-1* is required for cell cycle exit in *C. elegans* and identifies a novel gene family. *Cell* 85: 829–839. [https://doi.org/10.1016/S0092-8674\(00\)81267-2](https://doi.org/10.1016/S0092-8674(00)81267-2)
- Kipreos, E. T., S. P. Gohel, and E. M. Hedgecock, 2000 The *C. elegans* F-box/WD-repeat protein LIN-23 functions to limit cell division during development. *Development* 127: 5071–5082.
- Kirienko, N. V., and D. S. Fay, 2007 Transcriptome profiling of the *C. elegans* Rb ortholog reveals diverse developmental roles. *Dev. Biol.* 305: 674–684. <https://doi.org/10.1016/j.ydbio.2007.02.021>
- Kishimoto, T., 2015 Entry into mitosis: a solution to the decades-long enigma of MPF. *Chromosoma* 124: 417–428. <https://doi.org/10.1007/s00412-015-0508-y>
- Kitagawa, R., E. Law, L. Tang, and A. M. Rose, 2002 The Cdc20 homolog, FZY-1, and its interacting protein, IFY-1, are required for proper chromosome segregation in *Caenorhabditis elegans*. *Curr. Biol.* 12: 2118–2123. [https://doi.org/10.1016/S0960-9822\(02\)01392-1](https://doi.org/10.1016/S0960-9822(02)01392-1)
- Korenjak, M., B. Taylor-Harding, U. K. Binné, J. S. Satterlee, O. Stevaux *et al.*, 2004 Native E2F/RBF complexes contain Myb-interacting proteins and repress transcription of developmentally controlled E2F target genes. *Cell* 119: 181–193. <https://doi.org/10.1016/j.cell.2004.09.034>
- Korzelius, J., I. The, S. Ruijtenberg, V. Portegijs, H. Xu *et al.*, 2011a *C. elegans* MCM-4 is a general DNA replication and checkpoint component with an epidermis-specific requirement for growth and viability. *Dev. Biol.* 350: 358–369. <https://doi.org/10.1016/j.ydbio.2010.12.009>
- Korzelius, J., I. The, S. Ruijtenberg, M. B. W. Prinsen, V. Portegijs *et al.*, 2011b *Caenorhabditis elegans* cyclin D/CDK4 and cyclin E/CDK2 induce distinct cell cycle re-entry programs in differentiated muscle cells. *PLoS Genet.* 7: e1002362. <https://doi.org/10.1371/journal.pgen.1002362>
- Krause, M., A. Fire, S. W. Harrison, J. Priess, and H. Weintraub, 1990 CeMyoD accumulation defines the body wall muscle cell fate during *C. elegans* embryogenesis. *Cell* 63: 907–919. [https://doi.org/10.1016/0092-8674\(90\)90494-Y](https://doi.org/10.1016/0092-8674(90)90494-Y)
- Kreutzer, M. A., J. P. Richards, M. N. De Silva-Udawatta, J. J. Temenak, J. A. Knoblich *et al.*, 1995 *Caenorhabditis elegans* cyclin A- and B-type genes: a cyclin A multigene family, an ancestral cyclin B3 and differential germline expression. *J. Cell Sci.* 108: 2415–2424.
- Lamont, L. B., S. L. Crittenden, D. Bernstein, M. Wickens, and J. Kimble, 2004 FBF-1 and FBF-2 regulate the size of the mitotic region in the *C. elegans* germline. *Dev. Cell* 7: 697–707. <https://doi.org/10.1016/j.devcel.2004.09.013>
- Lee, Y. U., M. Son, J. Kim, Y. H. Shim, and I. Kawasaki, 2016 CDC-25.2, a *C. elegans* ortholog of *cdc25*, is essential for the progression of intestinal divisions. *Cell Cycle* 15: 654–666. <https://doi.org/10.1080/15384101.2016.1146839>
- Leng, X., M. Noble, P. D. Adams, J. Qin, and J. W. Harper, 2002 Reversal of growth suppression by p107 via direct phosphorylation by cyclin D1/cyclin-dependent kinase 4. *Mol. Cell. Biol.* 22: 2242–2254. <https://doi.org/10.1128/MCB.22.7.2242-2254.2002>
- EHernault, S. W., D. C. Shakes, and S. Ward, 1988 Developmental genetics of chromosome I spermatogenesis-defective mutants in the nematode *Caenorhabditis elegans*. *Genetics* 120: 435–452.
- Lilly, M. A., and R. J. Duronio, 2005 New insights into cell cycle control from the *Drosophila* endocycle. *Oncogene* 24: 2765–2775. <https://doi.org/10.1038/sj.onc.1208610>
- Liu, J., and E. T. Kipreos, 2000 Evolution of cyclin-dependent kinases (CDKs) and CDK-activating kinases (CAKs): differential conservation of CAKs in yeast and metazoa. *Mol. Biol. Evol.* 17: 1061–1074. <https://doi.org/10.1093/oxfordjournals.molbev.a026387>
- Liu, J., S. Vasudevan, and E. T. Kipreos, 2004 CUL-2 and ZYG-11 promote meiotic anaphase II and the proper placement of the anterior-posterior axis in *C. elegans*. *Development* 131: 3513–3525. <https://doi.org/10.1242/dev.01245>
- Lorson, M. A., H. R. Horvitz, and S. van den Heuvel, 2000 LIN-5 is a novel component of the spindle apparatus required for chromosome segregation and cleavage plane specification in *Caenorhabditis elegans*. *J. Cell Biol.* 148: 73–86. <https://doi.org/10.1083/jcb.148.1.73>
- Lu, Z., and T. Hunter, 2010 Ubiquitylation and proteasomal degradation of the p21<sup>Cip1</sup>, p27<sup>Kip1</sup> and p57<sup>Kip2</sup> CDK inhibitors. *Cell Cycle* 9: 2342–2352. <https://doi.org/10.4161/cc.9.12.11988>
- Maruyama, R., N. V. Velarde, R. Klancer, S. Gordon, P. Kadandale *et al.*, 2007 EGG-3 regulates cell-surface and cortex rearrangements during egg activation in *Caenorhabditis elegans*. *Curr. Biol.* 17: 1555–1560. <https://doi.org/10.1016/j.cub.2007.08.011>
- Masui, Y., and C. L. Markert, 1971 Cytoplasmic control of nuclear behavior during meiotic maturation of frog oocytes. *J. Exp. Zool.* 177: 129–145. <https://doi.org/10.1002/jez.1401770202>
- Matus, D. Q., L. L. Lohmer, L. C. Kelley, A. J. Schindler, A. Q. Kohrman *et al.*, 2015 Invasive cell fate requires G1 cell-cycle arrest and histone deacetylase-mediated changes in gene expression. *Dev. Cell* 35: 162–174. <https://doi.org/10.1016/j.devcel.2015.10.002>
- McCarter, J., B. Bartlett, T. Dang, and T. Schedl, 1997 Soma-germ cell interactions in *Caenorhabditis elegans*: multiple events of hermaphrodite germline development require the somatic

- sheath and spermathecal lineages. *Dev. Biol.* 181: 121–143. <https://doi.org/10.1006/dbio.1996.8429>
- McNally, K. L., and F. J. McNally, 2005 Fertilization initiates the transition from anaphase I to metaphase II during female meiosis in *C. elegans*. *Dev. Biol.* 282: 218–230. <https://doi.org/10.1016/j.ydbio.2005.03.009>
- Méchal, M., 2010 Eukaryotic DNA replication origins: many choices for appropriate answers. *Nat. Rev. Mol. Cell Biol.* 11: 728–738. <https://doi.org/10.1038/nrm2976>
- Medema, R. H., G. J. Kops, J. L. Bos, and B. M. Burgering, 2000 AFX-like forkhead transcription factors mediate cell-cycle regulation by Ras and PKB through p27<sup>kip1</sup>. *Nature* 404: 782–787. <https://doi.org/10.1038/35008115>
- Merlet, J., J. Burger, N. Tavernier, B. Richaudeau, J. E. Gomes *et al.*, 2010 The CRL2<sup>LRR-1</sup> ubiquitin ligase regulates cell cycle progression during *C. elegans* development. *Development* 137: 3857–3866. <https://doi.org/10.1242/dev.054866>
- Michael, W. M., 2016 Cyclin CYB-3 controls both S-phase and mitosis and is asymmetrically distributed in the early *C. elegans* embryo. *Development* 143: 3119–3127. <https://doi.org/10.1242/dev.141226>
- Michaelson, D., D. Z. Korta, Y. Capua, and E. J. Hubbard, 2010 Insulin signaling promotes germline proliferation in *C. elegans*. *Development* 137: 671–680. <https://doi.org/10.1242/dev.042523>
- Miller, D. L., and M. B. Roth, 2009 *C. elegans* are protected from lethal hypoxia by an embryonic diapause. *Curr. Biol.* 19: 1233–1237. <https://doi.org/10.1016/j.cub.2009.05.066>
- Miller, M. A., V. Q. Nguyen, M. H. Lee, M. Kosinski, T. Schedl *et al.*, 2001 A sperm cytoskeletal protein that signals oocyte meiotic maturation and ovulation. *Science* 291: 2144–2147.
- Miller, M. A., P. J. Ruest, M. Kosinski, S. K. Hanks, and D. Greenstein, 2003 An Eph receptor sperm-sensing control mechanism for oocyte meiotic maturation in *Caenorhabditis elegans*. *Genes Dev.* 17: 187–200.
- Mohammad, A., K. V. Broek, C. Wang, A. Daryabeigi, V. Jantsch *et al.*, 2018 Initiation of meiotic development is controlled by three posttranscriptional pathways in *Caenorhabditis elegans*. *Genetics* 209: 1197–1224. <https://doi.org/10.1534/genetics.118.300985>
- Mukherjee, M., S. N. Chaudhari, R. S. Balachandran, A. S. Vagasi, and E. T. Kipreos, 2017 Dafachronic acid inhibits *C. elegans* germ cell proliferation in a DAF-12-dependent manner. *Dev. Biol.* 432: 215–221. <https://doi.org/10.1016/j.ydbio.2017.10.014>
- Murphy, C. T., and P. J. Hu, 2013 Insulin/insulin-like growth factor signaling in *C. elegans* (December 26, 2013), *WormBook*, ed. The *C. elegans* Research Community WormBook, doi/10.1895/wormbook.1.164.1, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.164.1>
- Narbonne, P., P. S. Maddox, and J. C. Labbe, 2015 DAF-18/PTEN locally antagonizes insulin signalling to couple germline stem cell proliferation to oocyte needs in *C. elegans*. *Development* 142: 4230–4241. <https://doi.org/10.1242/dev.130252>
- Nasmyth, K., 2005 How do so few control so many? *Cell* 120: 739–746. <https://doi.org/10.1016/j.cell.2005.03.006>
- Nimmo, R., A. Antebi, and A. Woollard, 2005 *mab-2* encodes RNT-1, a *C. elegans* Runx homologue essential for controlling cell proliferation in a stem cell-like developmental lineage. *Development* 132: 5043–5054. <https://doi.org/10.1242/dev.02102>
- Nishi, Y., E. Rogers, S. M. Robertson, and R. Lin, 2008 Polo kinases regulate *C. elegans* embryonic polarity via binding to DYRK2-primed MEX-5 and MEX-6. *Development* 135: 687–697. <https://doi.org/10.1242/dev.013425>
- Nurse, P., 2002 Cyclin dependent kinases and cell cycle control (Nobel Lecture). *ChemBioChem* 3: 596–603. [https://doi.org/10.1002/1439-7633\(20020703\)3:7<596::AID-CBIC596>3.0.CO;2-U](https://doi.org/10.1002/1439-7633(20020703)3:7<596::AID-CBIC596>3.0.CO;2-U)
- Nystrom, J., Z. Z. Shen, M. Aili, A. J. Flemming, A. Leroi *et al.*, 2002 Increased or decreased levels of *Caenorhabditis elegans lon-3*, a gene encoding a collagen, cause reciprocal changes in body length. *Genetics* 161: 83–97.
- Nystul, T. G., J. P. Goldmark, P. A. Padilla, and M. B. Roth, 2003 Suspended animation in *C. elegans* requires the spindle checkpoint. *Science* 302: 1038–1041. <https://doi.org/10.1126/science.1089705>
- O’Connell, K. F., C. M. Leys, and J. G. White, 1998 A genetic screen for temperature-sensitive cell-division mutants of *Caenorhabditis elegans*. *Genetics* 149: 1303–1321.
- Orford, K. W., and D. T. Scadden, 2008 Deconstructing stem cell self-renewal: genetic insights into cell-cycle regulation. *Nat. Rev. Genet.* 9: 115–128. <https://doi.org/10.1038/nrg2269>
- Pacquelet, A., 2017 Asymmetric cell division in the one-cell *C. elegans* embryo: multiple steps to generate cell size asymmetry. *Results Probl. Cell Differ.* 61: 115–140. [https://doi.org/10.1007/978-3-319-53150-2\\_5](https://doi.org/10.1007/978-3-319-53150-2_5)
- Park, F. D., and J. R. Priess, 2003 Establishment of POP-1 asymmetry in early *C. elegans* embryos. *Development* 130: 3547–3556. <https://doi.org/10.1242/dev.00563>
- Park, M., and M. W. Krause, 1999 Regulation of postembryonic G1 cell cycle progression in *Caenorhabditis elegans* by a cyclin D/CDK-like complex. *Development* 126: 4849–4860.
- Parry, J. M., N. V. Velarde, A. J. Lefkovith, M. H. Zegarek, J. S. Hang *et al.*, 2009 EGG-4 and EGG-5 link events of the oocyte-to-embryo transition with meiotic progression in *C. elegans*. *Curr. Biol.* 19: 1752–1757. <https://doi.org/10.1016/j.cub.2009.09.015>
- Pellettieri, J., V. Reinke, S. K. Kim, and G. Seydoux, 2003 Coordinate activation of maternal protein degradation during the egg-to-embryo transition in *C. elegans*. *Dev. Cell* 5: 451–462. [https://doi.org/10.1016/S1534-5807\(03\)00231-4](https://doi.org/10.1016/S1534-5807(03)00231-4)
- Pinkston, J. M., D. Garigan, M. Hansen, and C. Kenyon, 2006 Mutations that increase the life span of *C. elegans* inhibit tumor growth. *Science* 313: 971–975. <https://doi.org/10.1126/science.1121908>
- Pozo, P. N., and J. G. Cook, 2017 Regulation and function of Cdt1; a key factor in cell proliferation and genome stability. *Genes (Basel)* 8: 2. <https://doi.org/10.3390/genes8010002>
- Quintin, S., P. E. Mains, A. Zinke, and A. A. Hyman, 2003 The *mbk-2* kinase is required for inactivation of MEI-1/katanin in the one-cell *Caenorhabditis elegans* embryo. *EMBO Rep.* 4: 1175–1181. <https://doi.org/10.1038/sj.embor.7400029>
- Rahman, M. M., S. Rosu, D. Joseph-Strauss, and O. Cohen-Fix, 2014 Down-regulation of tricarboxylic acid (TCA) cycle genes blocks progression through the first mitotic division in *Caenorhabditis elegans* embryos. *Proc. Natl. Acad. Sci. USA* 111: 2602–2607. <https://doi.org/10.1073/pnas.1311635111>
- Rausch, M., M. Ecsedi, H. Bartake, A. Müllner, and H. Grosshans, 2015 A genetic interactome of the *let-7* microRNA in *C. elegans*. *Dev. Biol.* 401: 276–286. <https://doi.org/10.1016/j.ydbio.2015.02.013>
- Riddle, D. L., and P. S. Albert, 1997 Genetic and environmental regulation of dauer larva development in *C. elegans II*, edited by D. L. Riddle, T. Blumenthal, B. J. Meyer, and J. R. Priess. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Riedel, C. G., R. H. Downen, G. F. Lourenco, N. V. Kirienko, T. Heimbucher *et al.*, 2013 DAF-16 employs the chromatin remodeler SWI/SNF to promote stress resistance and longevity. *Nat. Cell Biol.* 15: 491–501. <https://doi.org/10.1038/ncb2720>
- Riera, A., S. Tognetti, and C. Speck, 2014 Helicase loading: how to build a MCM2–7 double-hexamers. *Semin. Cell Dev. Biol.* 30: 104–109. <https://doi.org/10.1016/j.semcdb.2014.03.008>
- Rivers, D. M., S. Moreno, M. Abraham, and J. Ahringer, 2008 PAR proteins direct asymmetry of the cell cycle regulators Polo-like

- kinase and Cdc25. *J. Cell Biol.* 180: 877–885. <https://doi.org/10.1083/jcb.200710018>
- Roiz, D., J. M. Escobar-Restrepo, P. Leu, and A. Hajnal, 2016 The *C. elegans* hox gene *lin-39* controls cell cycle progression during vulval development. *Dev. Biol.* 418: 124–134. <https://doi.org/10.1016/j.ydbio.2016.07.018>
- Rose, L., and P. Gönczy, 2014 Polarity establishment, asymmetric division and segregation of fate determinants in early *C. elegans* embryos (December 30, 2014), *WormBook*, ed. The *C. elegans* Research Community WormBook, doi/10.1895/wormbook.1.30.2, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.30.2>
- Rosu, S., and O. Cohen-Fix, 2017 Live-imaging analysis of germ cell proliferation in the *C. elegans* adult supports a stochastic model for stem cell proliferation. *Dev. Biol.* 423: 93–100. <https://doi.org/10.1016/j.ydbio.2017.02.008>
- Roy, D., D. Michaelson, T. Hochman, A. Santella, Z. Bao *et al.*, 2016 Cell cycle features of *C. elegans* germline stem/progenitor cells vary temporally and spatially. *Dev. Biol.* 409: 261–271. <https://doi.org/10.1016/j.ydbio.2015.10.031>
- Rubin, S. M., 2013 Deciphering the retinoblastoma protein phosphorylation code. *Trends Biochem. Sci.* 38: 12–19. <https://doi.org/10.1016/j.tibs.2012.10.007>
- Ruijtenberg, S., and S. van den Heuvel, 2015 G1/S Inhibitors and the SWI/SNF complex control cell-cycle exit during muscle differentiation. *Cell* 162: 300–313. <https://doi.org/10.1016/j.cell.2015.06.013>
- Sadasivam, S., and J. A. DeCaprio, 2013 The DREAM complex: master coordinator of cell cycle-dependent gene expression. *Nat. Rev. Cancer* 13: 585–595 (erratum: *Nat. Rev. Cancer* 13: 752). <https://doi.org/10.1038/nrc3556>
- Saffer, A. M., D. H. Kim, A. van Oudenaarden, and H. R. Horvitz, 2011 The *Caenorhabditis elegans* synthetic multivulva genes prevent Ras pathway activation by tightly repressing global ectopic expression of *lin-3* EGF. *PLoS Genet.* 7: e1002418. <https://doi.org/10.1371/journal.pgen.1002418>
- Saito, R. M., A. Perreault, B. Peach, J. S. Satterlee, and S. van den Heuvel, 2004 The CDC-14 phosphatase controls developmental cell-cycle arrest in *C. elegans*. *Nat. Cell Biol.* 6: 777–783. <https://doi.org/10.1038/ncb1154>
- Segref, A., J. Cabello, C. Clucas, R. Schnabel, and I. L. Johnstone, 2010 Fate specification and tissue-specific cell cycle control of the *Caenorhabditis elegans* intestine. *Mol. Biol. Cell* 21: 725–738. <https://doi.org/10.1091/mbc.e09-04-0268>
- Seidel, H. S., and J. Kimble, 2011 The oogenic germline starvation response in *C. elegans*. *PLoS One* 6: e28074. <https://doi.org/10.1371/journal.pone.0028074>
- Seidel, H. S., and J. Kimble, 2015 Cell-cycle quiescence maintains *Caenorhabditis elegans* germline stem cells independent of GLP-1/Notch. *Elife* 4: e10832. <https://doi.org/10.7554/eLife.10832>
- Selhub, J., 2002 Folate, vitamin B12 and vitamin B6 and one carbon metabolism. *J. Nutr. Health Aging* 6: 39–42.
- Sherr, C. J., 1996 Cancer cell cycles. *Science* 274: 1672–1677. <https://doi.org/10.1126/science.274.5293.1672>
- Shirayama, M., M. C. Soto, T. Ishidate, S. Kim, K. Nakamura *et al.*, 2006 The conserved kinases CDK-1, GSK-3, KIN-19, and MBK-2 promote OMA-1 destruction to regulate the oocyte-to-embryo transition in *C. elegans*. *Curr. Biol.* 16: 47–55. <https://doi.org/10.1016/j.cub.2005.11.070>
- Singaravelu, G., and A. Singson, 2013 Calcium signaling surrounding fertilization in the nematode *Caenorhabditis elegans*. *Cell Calcium* 53: 2–9. <https://doi.org/10.1016/j.ceca.2012.11.009>
- Singaravelu, G., I. Chatterjee, S. Rahimi, M. K. Druzhinina, L. Kang *et al.*, 2012 The sperm surface localization of the TRP-3/SPE-41 Ca<sup>2+</sup>-permeable channel depends on SPE-38 function in *Caenorhabditis elegans*. *Dev. Biol.* 365: 376–383. <https://doi.org/10.1016/j.ydbio.2012.02.037>
- Slack, F. J., M. Basson, Z. Liu, V. Ambros, H. R. Horvitz *et al.*, 2000 The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the LIN-29 transcription factor. *Mol. Cell* 5: 659–669. [https://doi.org/10.1016/S1097-2765\(00\)80245-2](https://doi.org/10.1016/S1097-2765(00)80245-2)
- Smith, J., L. M. Tho, N. Xu, and D. A. Gillespie, 2010 The ATM-Chk2 and ATR-Chk1 pathways in DNA damage signaling and cancer. *Adv. Cancer Res.* 108: 73–112. <https://doi.org/10.1016/B978-0-12-380888-2.00003-0>
- Son, M., I. Kawasaki, B.-K. Oh, and Y.-H. Shim, 2016 LIN-23, an E3 ubiquitin ligase component, is required for the repression of CDC-25.2 activity during intestinal development in *Caenorhabditis elegans*. *Mol. Cells* 39: 834–840. <https://doi.org/10.14348/molcells.2016.0238>
- Sonneville, R., and P. Gönczy, 2004 *zyg-11* and *cul-2* regulate progression through meiosis II and polarity establishment in *C. elegans*. *Development* 131: 3527–3543. <https://doi.org/10.1242/dev.01244>
- Sonneville, R., M. Querenet, A. Craig, A. Gartner, and J. J. Blow, 2012 The dynamics of replication licensing in live *Caenorhabditis elegans* embryos. *J. Cell Biol.* 196: 233–246. <https://doi.org/10.1083/jcb.201110080>
- Sönnichsen, B., L. B. Koski, A. Walsh, P. Marschall, B. Neumann *et al.*, 2005 Full-genome RNAi profiling of early embryogenesis in *Caenorhabditis elegans*. *Nature* 434: 462–469. <https://doi.org/10.1038/nature03353>
- Sowa, J. N., A. S. Mutlu, F. Xia, and M. C. Wang, 2015 Olfaction modulates reproductive plasticity through neuroendocrine signaling in *Caenorhabditis elegans*. *Curr. Biol.* 25: 2284–2289. <https://doi.org/10.1016/j.cub.2015.07.023>
- Spike, C. A., D. Coetzee, C. Eichten, X. Wang, D. Hansen *et al.*, 2014a The TRIM-NHL protein LIN-41 and the OMA RNA-binding proteins antagonistically control the prophase-to-metaphase transition and growth of *Caenorhabditis elegans* oocytes. *Genetics* 198: 1535–1558. <https://doi.org/10.1534/genetics.114.168831>
- Spike, C. A., D. Coetzee, Y. Nishi, T. Guven-Ozkan, M. Oldenbroek *et al.*, 2014b Translational control of the oogenic program by components of OMA ribonucleoprotein particles in *Caenorhabditis elegans*. *Genetics* 198: 1513–1533. <https://doi.org/10.1534/genetics.114.168823>
- Srinivasan, D. G., R. M. Fisk, H. Xu, and S. van den Heuvel, 2003 A complex of LIN-5 and GPR proteins regulates G protein signaling and spindle function in *C. elegans*. *Genes Dev.* 17: 1225–1239. <https://doi.org/10.1101/gad.1081203>
- Starostina, N. G., J. M. Simpliciano, M. A. McGuirk, and E. T. Kipreos, 2010 CRL2(LRR-1) targets a CDK inhibitor for cell cycle control in *C. elegans* and actin-based motility regulation in human cells. *Dev. Cell* 19: 753–764. <https://doi.org/10.1016/j.devcel.2010.10.013>
- Stegmeier, F., and A. Amon, 2004 Closing mitosis: the functions of the Cdc14 phosphatase and its regulation. *Annu. Rev. Genet.* 38: 203–232. <https://doi.org/10.1146/annurev.genet.38.072902.093051>
- Sternberg, P. W., 2005 Vulval development (June 25, 2005), *WormBook*, ed. The *C. elegans* Research Community WormBook, doi/10.1895/wormbook.1.6.1, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.6.1>
- Stevens, H., A. B. Williams, and W. M. Michael, 2016 Cell-type specific responses to DNA replication stress in early *C. elegans* embryos. *PLoS One* 11: e0164601. <https://doi.org/10.1371/journal.pone.0164601>
- Stitzel, M. L., J. Pellettieri, and G. Seydoux, 2006 The *C. elegans* DYRK kinase MBK-2 marks oocyte proteins for degradation in response to meiotic maturation. *Curr. Biol.* 16: 56–62. <https://doi.org/10.1016/j.cub.2005.11.063>
- Stitzel, M. L., K. C.-C. Cheng, and G. Seydoux, 2007 Regulation of MBK-2/Dyrk kinase by dynamic cortical anchoring during the



- oocyte-to-zygote transition. *Curr. Biol.* 17: 1545–1554. <https://doi.org/10.1016/j.cub.2007.08.049>
- Subramaniam, K., and G. Seydoux, 1999 *nos-1* and *nos-2*, two genes related to *Drosophila* *nanos*, regulate primordial germ cell development and survival in *Caenorhabditis elegans*. *Development* 126: 4861–4871.
- Sulston, J. E., and H. R. Horvitz, 1977 Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev. Biol.* 56: 110–156. [https://doi.org/10.1016/0012-1606\(77\)90158-0](https://doi.org/10.1016/0012-1606(77)90158-0)
- Sulston, J. E., and H. R. Horvitz, 1981 Abnormal cell lineages in mutants of the nematode *Caenorhabditis elegans*. *Dev. Biol.* 82: 41–55. [https://doi.org/10.1016/0012-1606\(81\)90427-9](https://doi.org/10.1016/0012-1606(81)90427-9)
- Sulston, J. E., E. Schierenberg, J. G. White, and J. N. Thomson, 1983 The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* 100: 64–119.
- Tain, L. S., E. Lozano, A. G. Saez, and A. M. Leroi, 2008 Dietary regulation of hypodermal polyploidization in *C. elegans*. *BMC Dev. Biol.* 8: 28. <https://doi.org/10.1186/1471-213X-8-28>
- Tanaka, S., and H. Araki, 2013 Helicase activation and establishment of replication forks at chromosomal origins of replication. *Cold Spring Harb. Perspect. Biol.* 5: a010371. <https://doi.org/10.1101/cshperspect.a010371>
- Tavernier, N., A. Noatynska, C. Panbianco, L. Martino, L. Van Hove *et al.*, 2015 Cdk1 phosphorylates SPAT-1/Bora to trigger PLK-1 activation and drive mitotic entry in *C. elegans* embryos. *J. Cell Biol.* 208: 661–669. <https://doi.org/10.1083/jcb.201408064>
- Tenlen, J. R., J. N. Molk, N. London, B. D. Page, and J. R. Priess, 2008 MEX-5 asymmetry in one-cell *C. elegans* embryos requires PAR-4- and PAR-1-dependent phosphorylation. *Development* 135: 3665–3675. <https://doi.org/10.1242/dev.027060>
- The, I., S. Ruijtenberg, B. P. Bouchet, A. Cristobal, M. B. W. Prinsen *et al.*, 2015 Rb and FZR1/Cdh1 determine CDK4/6-cyclin D requirement in *C. elegans* and human cancer cells. *Nat. Commun.* 6: 5906. <https://doi.org/10.1038/ncomms6906>
- Thondamal, M., M. Witting, P. Schmitt-Kopplin, and H. Aguilaniu, 2014 Steroid hormone signalling links reproduction to lifespan in dietary-restricted *Caenorhabditis elegans*. *Nat. Commun.* 5: 4879. <https://doi.org/10.1038/ncomms5879>
- Tilmann, C., and J. Kimble, 2005 Cyclin D regulation of a sexually dimorphic asymmetric cell division. *Dev. Cell* 9: 489–499. <https://doi.org/10.1016/j.devcel.2005.09.004>
- Tsakamoto, T., M. D. Gearhart, C. A. Spike, G. Huelgas-Morales, M. Mews *et al.*, 2017 LIN-41 and OMA ribonucleoprotein complexes mediate a translational repression-to-activation switch controlling oocyte meiotic maturation and the oocyte-to-embryo transition in *Caenorhabditis elegans*. *Genetics* 206: 2007–2039. <https://doi.org/10.1534/genetics.117.203174>
- van den Heuvel, S., 2005 Cell-cycle regulation (September 21, 2005), *WormBook*, ed. The *C. elegans* Research Community WormBook, doi/10.1895/wormbook.1.28.1, <http://www.wormbook.org>. <https://doi.org/10.1895/wormbook.1.28.1>
- van den Heuvel, S., and N. J. Dyson, 2008 Conserved functions of the pRB and E2F families. *Nat. Rev. Mol. Cell Biol.* 9: 713–724. <https://doi.org/10.1038/nrm2469>
- van der Voet, M., M. A. Lorson, D. G. Srinivasan, K. L. Bennett, and S. van den Heuvel, 2009 *C. elegans* mitotic cyclins have distinct as well as overlapping functions in chromosome segregation. *Cell Cycle* 8: 4091–4102. <https://doi.org/10.4161/cc.8.24.10171>
- Vasudevan, S., N. G. Starostina, and E. T. Kipreos, 2007 The *Caenorhabditis elegans* cell-cycle regulator ZYG-11 defines a conserved family of CUL-2 complex components. *EMBO Rep.* 8: 279–286. <https://doi.org/10.1038/sj.embor.7400895>
- Vidal, A., and A. Koff, 2000 Cell-cycle inhibitors: three families united by a common cause. *Gene* 247: 1–15.
- von Schubert, C., F. Cubizolles, J. M. Bracher, T. Sliedrecht, G. Kops *et al.*, 2015 Plk1 and Mps1 cooperatively regulate the spindle assembly checkpoint in human cells. *Cell Rep.* 12: 66–78. <https://doi.org/10.1016/j.celrep.2015.06.007>
- Von Stetina, J. R., and T. L. Orr-Weaver, 2011 Developmental control of oocyte maturation and egg activation in metazoan models. *Cold Spring Harb. Perspect. Biol.* 3: a005553. <https://doi.org/10.1101/cshperspect.a005553>
- Waaaijers, S., and M. Boxem, 2014 Engineering the *Caenorhabditis elegans* genome with CRISPR/Cas9. *Methods* 68: 381–388.
- Wallenfang, M. R., and G. Seydoux, 2002 *cdk-7* is required for mRNA transcription and cell cycle progression in *Caenorhabditis elegans* embryos. *Proc. Natl. Acad. Sci. USA* 99: 5527–5532. <https://doi.org/10.1073/pnas.082618399>
- Ward, S., and J. S. Carrel, 1979 Fertilization and sperm competition in the nematode *Caenorhabditis elegans*. *Dev. Biol.* 73: 304–321. [https://doi.org/10.1016/0012-1606\(79\)90069-1](https://doi.org/10.1016/0012-1606(79)90069-1)
- Wei, W., N. G. Ayad, Y. Wan, G. J. Zhang, M. W. Kirschner *et al.*, 2004 Degradation of the SCF component Skp2 in cell-cycle phase G1 by the anaphase-promoting complex. *Nature* 428: 194–198. <https://doi.org/10.1038/nature02381>
- Wilson, M. A., R. V. Hoch, N. R. Ashcroft, M. E. Kosinski, and A. Golden, 1999 A *Caenorhabditis elegans wee1* homolog is expressed in a temporally and spatially restricted pattern during embryonic development. *Biochim. Biophys. Acta* 1445: 99–109. [https://doi.org/10.1016/S0167-4781\(99\)00027-5](https://doi.org/10.1016/S0167-4781(99)00027-5)
- Wirt, S. E., A. S. Adler, V. Gebala, J. M. Weimann, B. E. Schaffer *et al.*, 2010 G1 arrest and differentiation can occur independently of Rb family function. *J. Cell Biol.* 191: 809–825. <https://doi.org/10.1083/jcb.201003048>
- Woollard, A., and J. Hodgkin, 1999 *stu-7/air-2* is a *C. elegans* aurora homologue essential for chromosome segregation during embryonic and post-embryonic development. *Mech. Dev.* 82: 95–108. [https://doi.org/10.1016/S0925-4773\(99\)00020-9](https://doi.org/10.1016/S0925-4773(99)00020-9)
- Xia, D., Y. Zhang, X. Huang, Y. Sun, and H. Zhang, 2007 The *C. elegans* CBF $\beta$  homolog, BRO-1, regulates the proliferation, differentiation and specification of the stem cell-like seam cell lineages. *Dev. Biol.* 309: 259–272. <https://doi.org/10.1016/j.ydbio.2007.07.020>
- Yanagi, K., T. Mizuno, T. Tsuyama, S. Tada, Y. Iida *et al.*, 2005 *Caenorhabditis elegans* geminin homologue participates in cell cycle regulation and germ line development. *J. Biol. Chem.* 280: 19689–19694. <https://doi.org/10.1074/jbc.C500070200>
- Yanowitz, J., and A. Fire, 2005 Cyclin D involvement demarcates a late transition in *C. elegans* embryogenesis. *Dev. Biol.* 279: 244–251. <https://doi.org/10.1016/j.ydbio.2004.12.022>
- Yuan, K., and P. H. O'Farrell, 2015 Cyclin B3 is a mitotic cyclin that promotes the metaphase-anaphase transition. *Curr. Biol.* 25: 811–816. <https://doi.org/10.1016/j.cub.2015.01.053>
- Yuan, K., C. A. Sella, A. W. Shermoen, and P. H. O'Farrell, 2016 Timing the *Drosophila* mid-blastula transition: a cell cycle-centered view. *Trends Genet.* 32: 496–507. <https://doi.org/10.1016/j.tig.2016.05.006>
- Zhang, T., S.-T. Qi, L. Huang, X.-S. Ma, Y.-C. Ouyang *et al.*, 2015 Cyclin B3 controls anaphase onset independent of spindle assembly checkpoint in meiotic oocytes. *Cell Cycle* 14: 2648–2654. <https://doi.org/10.1080/15384101.2015.1064567>
- Zhong, W., H. Feng, F. E. Santiago, and E. T. Kipreos, 2003 CUL-4 ubiquitin ligase maintains genome stability by restraining DNA-replication licensing. *Nature* 423: 885–889. <https://doi.org/10.1038/nature01747>
- Zitouni, S., C. Nabais, S. C. Jana, A. Guerrero, and M. Bettencourt-Dias, 2014 Polo-like kinases: structural variations lead to multiple functions. *Nat. Rev. Mol. Cell Biol.* 15: 433–452. <https://doi.org/10.1038/nrm3819>

Communicating editor: G. Seydoux