



Genetic Transformation of Oat Mediated by *Agrobacterium* is enhanced with Sonication and Vacuum Infiltration

Nagesh Dattgonde, Sharad Tiwari *, Swapnil Sapre, Iti Gontia-Mishra

Biotechnology Centre, Jawaharlal Nehru Agriculture University, Jabalpur, India

* Corresponding author: Sharad Tiwari, Biotechnology Centre, Jawaharlal Nehru Agriculture University, Jabalpur, India. Phone: +919424658241; Fax: +917612681021. E-mail: dirbcjbp.jnkvv-mp@nic.in

Background: Oat (*Avena sativa*) with high nutritive value and fiber content is used as the main food grain in many countries for human diet as well as animal feed. Recently, it became difficult to transfer new genes through the conventional breeding due to the lack of desirable traits.

Objectives: The current study aimed at achieving a standardized protocol for *Agrobacterium*-mediated transformation in oat.

Materials and Methods: For oat transformation, mature seeds were sterilized, germinated, and used for explants generation. *Agrobacterium tumefaciens* GV3101 with the binary vector pCAMBIA 1305.1, which carries *gus* as reporter gene, was utilized in the transformation. The co-cultivation treatment assisted with sonication, and vacuum infiltration, and their combination was employed for transformation with different incubation periods of 48, 72, and 96 hours under the dark conditions.

Results: Among the different transformation treatments, the vacuum treatment with 72 hours dark incubation had the best results. Vacuum infiltration of the cultures from leaf base produced a maximum of 25% hygromycin-resistant explants. These explants upon GUS assay and PCR analysis revealed 21.85% and 19.04% transformation efficiency, respectively.

Conclusions: It could be concluded that vacuum infiltration assisted *Agrobacterium*-mediated transformation is the most efficient method to conduct the genetic improvement of the oat using transformation protocol.

Keywords: *Avena*; *Agrobacterium*; Sonication; Vacuum

1. Background

Oat is grown as a multi-purpose crop; it is used as a grain, forage, and a rotation crop. It has rich nutritional value, since it contains the soluble fibers with hypocholesterolemic properties (1). It has excellent growth habit, rapid recovery after cutting, and excellent quality fodder. In India, the yield of oat is quite low, compared with other cereal crops, owing to various environmental stresses. A number of efforts are made towards the development of new cultivars of oat with an improved disease, pest, and herbicide tolerance. The genetic improvement of the commercially important oat cultivars through the conventional breeding is arduous, slow, and costly. A huge improvement is noticed in several monocots by incorporating various agronomical traits using genetic engineering (2, 3). The development of a robust regeneration protocol is necessary to transform cereals (4). *Agrobacterium*-mediated transformation is a complex procedure controlled by diverse parameters of the bacterial, host, and environmental origin. By this technique of genetic

transformation, a wide range of characters such as improvement of the plant tolerance to the environmental stresses, disease resistance, enhanced crop productivity, pest resistance, phytoremediation, and improved nutritional content of the crop plants are developed (5). To increase the transformation efficacy in the plants, sonication and vacuum infiltration play an important role (6, 7). Sonication produces micro-wounds by cavitation throughout the surface of tissue whereas vacuum infiltration provides a rapid entry of the *Agrobacterium* into the plant tissue. There are several reports of the enhanced genetic transformation using these techniques in various crops namely radish (8), citrus (9), cowpea (10), banana (6), lentil (11), and sugarcane (7). Gasparis *et al.*, (12) reported *Agrobacterium*-mediated transformation of the oat using immature embryo and leaf explants. In the current investigation, efforts were made to improve the efficiency of the *Agrobacterium*-mediated

transformation in oat by a conjoint effect of sonication and vacuum infiltration.

2. Objectives

The current study aimed at generating a proficient gene transfer system in oat cv. JO-1 using sonication-assisted *Agrobacterium*-mediated transformation (SAAT) and the vacuum-infiltration-assisted *Agrobacterium*-mediated transformation (VIAAT). The influence of different explants, sonication, and vacuum infiltration were evaluated in *Agrobacterium*-mediated genetic transformation of oat cv. JO-1.

3. Materials and Methods

3.1. Sterilization and Germination of Oat Seeds

Seeds of cultivar JO-1 were procured from the All India Coordinated Forage Research Project (ICAR), Department of Agronomy, JNKVV Jabalpur. Initially, oat seeds were rinsed thoroughly with water. Further, the seeds were kept in 1% Tween-20 solution for 20 minutes and then treated for two minutes with 70% ethanol. The seeds were again treated with 0.5% HgCl₂ solution for five minutes. Finally, seeds were germinated on the Murashige and Skoog (MS) basal medium (13). The six-day-old seedlings were used to obtain leaf base explants, whereas mature embryos were excised from overnight soaked seeds under aseptic conditions.

3.2. Screening of the Explants for Antibiotic Sensitivity

The explants were inoculated into the MS medium as a control, as well as different concentrations of the hygromycin (5 to 50 mg.L⁻¹) and cefotaxime (50 to 750 mg.L⁻¹) to assess antibiotic sensitivity.

3.3. *Agrobacterium* Culture

For genetic transformation studies, *Agrobacterium tumefaciens* strain GV3101 with plasmid pCAMBIA1305.1 (11846 base pair) harboring *gus* (β -glucuronidase) gene and catalytic intron under the regulation of CaMV35S promoter and *nos* terminator was used. The vector also had *hptII* (hygromycin phosphotransferase II) gene for plant selection with CaMV35S promoter and *polyA* terminator (Fig. 1A). *A. tumefaciens* was inoculated into a 50-mL Luria Bertani (LB) medium supplemented with rifampicin and kanamycin (50 mg.L⁻¹ each) and incubated at 28°C for 16 hours at 200 rpm. *Agrobacterium* culture of OD 0.6 (at wavelength of 600 nm) were centrifuged for five minutes at 28000 g and the bacterial cells' pellet was dissolved in the same amount of the co-cultivation broth (1X B₅ medium, 3% sucrose, 100 mg.L⁻¹ MES, 100 μ M acetosyringone, 1 mg.L⁻¹ GA₃, 1.67 mg.L⁻¹ BAP, 400 mg.L⁻¹ L-cysteine, 248 mg.L⁻¹ Na-thiosulphate, and 154 mg.L⁻¹ DTT).

3.4. Sonication-assisted *Agrobacterium*-mediated Transformation

For SAAT (17), calli developed from embryo and leaf base (Fig. 1B) were sonicated in a bath sonicator (HF-frequency: 35 KHz) submerged in 20 mL co-cultivation medium suspended with *Agrobacterium* for 30 seconds. These calli were placed at 25±2°C for 3-4 days in the dark conditions on a medium combination similar to co-cultivation medium with 8 g.L⁻¹ agar, but without *Agrobacterium*.

3.5. Vacuum-infiltration-assisted *Agrobacterium*-mediated transformation

During co-cultivation, explants submerged in co-cultivation medium with *Agrobacterium* were exposed to vacuum infiltration for 15 minutes.

3.6. Sonication and Vacuum-Infiltration-assisted-*Agrobacterium*-mediated Transformation

Conditions optimized in SAAT and VIAAT were applied in combination with achieve maximum *Agrobacterium*-mediated transformation efficiency (8, 9).

3.7. Selection and Plant Regeneration

The calli were washed with MS broth supplemented with cefotaxime (250 mg.L⁻¹). The calli were blot dried on sterile filter paper before placing them on MS medium containing hygromycin 20 mg.L⁻¹ and cefotaxime 500 mg.L⁻¹. Calli showing growth and morphogenesis (Fig. 1C and D) on 20 mg.L⁻¹ hygromycin supplemented MS medium were considered as putatively transformed plants. These calli were three times sub-cultured on hygromycin-containing medium at 10-12 days intervals.

3.8. GUS Histochemical Assay

Histochemical GUS analysis in explants was assayed after the third round of selection on antibiotics as described by Jefferson (14). Briefly, 10 explants with shoots were incubated in X-Gluc solution for overnight at room temperature under dark conditions, and washed with ethanol to remove chlorophyll. Histochemical localization of GUS activity was examined under stereomicroscope (Fig. 1E and F).

3.9. PCR Analysis of the Putative Transgenic Plants

The genomic DNA was isolated from the putative transformed plant tissue as per Saghai-Marooof et al., (15). PCR amplification of the marker genes: *hptII* and the CAMV 35S promoter region were performed to confirm the transformation as described by Sapre et al. (16) The amplification products of the *hptII* gene (499 bp) and CaMV 35S promoter (500 bp) were subjected to the electrophoresis and compared with those of the positive control (plant transformation vector; i.e., pCAMBIA plasmid) and negative control (DNA from non-transformed plants) (Fig. 2A and B).




Figure 1. Regeneration and transgenic plant selection of oat; (A) pCambia 1305 vector map; (B) Embryogenic callus from mature embryo explants; (C) Organogenesis from callus; (D) Green plantlets; (E, F) Expression of the *gus* gene in different tissues; (G, H) Plant with roots




Figure 2. PCR Analysis of the putative transformed oat plants; A) PCR amplification of *hptII* gene; B) PCR amplification of the *gus* gene; lane M, 100 bp DNA ladder; P, positive control; lanes 1-5, putative transformants; N, non-transformed control plant

4. Results

4.1. Effect of Hygromycin and Cefotaxime

The sensitivity of the explants to the antibiotic hygromycin was established prior to the actual transformation experiments by finding the minimal inhibitory concentration (MIC) to select presumed transformants. In the absence of antibiotics, the explants regenerated normally and produced calli and shoots; however, morphogenic ability of the cultures was inhibited with the addition of 20 mg.L⁻¹ hygromycin, since adventitious shoots exhibited very slow growth. While, 25 mg.L⁻¹ hygromycin inhibited shoot regeneration and resulted in the whitening of the cultures within 1-2 weeks. Explants dried due to necrosis within one week of the culture, when cultured on higher

concentrations of hygromycin (30-40 mg.L⁻¹). Hence, MS medium fortified with 20 mg.L⁻¹ of hygromycin was used to select the transformants in the subsequent transformation experiments.

4.2. Influence of co-Cultivation Duration

Duration of co-cultivation was influential on transformation efficiency. The 72-hour duration explants produced 13.87% (embryo) and 14.53% (leaf base) hygromycin-resistant cultures. The transformation efficiency at 72 hours was significantly higher compared with those of the 48 and 96 hours. These antibiotic-resistant cultures revealed the transformation efficiency of 7.36% and 6.71% in GUS histochemical assay and 7.30% and 6.09% in the PCR analysis of the transformation efficiency from the embryo and leaf-base explants, respectively (Table 1).

Table 1. Overall effects of co-cultivation period and different treatments on transformation efficiency

Total Number of Cultures	Explant	Transformation Efficiency, %		
		Hygromycin-resistant Culture	GUS Putative Culture	PCR Putative Culture
Sonication time (hour)				
48				
100	Embryo	22.59 ± 1.23 ^d	15.56 ± 0.87 ^e	14.78 ± 1.28 ^e
102	Leafbase	21.00 ± 0.98 ^e	14.81 ± 1.28 ^f	13.11 ± 1.61 ^{sh}
72				
75	Embryo	24.19 ± 0.87 ^b	19.78 ± 1.16 ^b	19.26 ± 1.59 ^b
79	Leafbase	23.22 ± 1.03 ^c	18.70 ± 1.05 ^c	17.58 ± 1.29 ^c
96				
68	Embryo	18.11 ± 1.76 ^j	14.59 ± 0.93 ^{fg}	13.67 ± 1.11 ^f
73	Leafbase	19.89 ± 0.76 ^f	14.33 ± 0.73 ^{fg}	12.89 ± 1.26 ^{gh}
Vacuum infiltration				
48				
102	Embryo	15.93 ± 0.98 ^{hi}	11.00 ± 0.81 ^k	10.15 ± 1.18 ^j
110	Leafbase	14.44 ± 0.91 ^k	11.89 ± 0.99 ^j	10.67 ± 1.02 ^j
72				
104	Embryo	24.33 ± 0.83 ^b	21.96 ± 1.18 ^a	20.74 ± 1.08 ^a
85	Leafbase	25.00 ± 0.96 ^a	21.85 ± 1.65 ^a	19.04 ± 0.96 ^b
96				
96	Embryo	17.37 ± 1.21 ⁱ	13.67 ± 1.41 ^{hi}	12.78 ± 0.84 ^b
93	Leafbase	18.89 ± 1.17 ^g	13.56 ± 1.29 ^{hi}	11.33 ± 0.92 ^j
Sonication + vacuum infiltration				
48				
89	Embryo	20.22 ± 0.88 ^f	14.30 ± 1.16 ^{fg}	13.85 ± 0.94 ^f
84	Leafbase	20.00 ± 0.71 ^f	14.11 ± 0.88 ^{gh}	13.33 ± 0.78 ^g
72				
86	Embryo	21.41 ± 1.10 ^e	17.67 ± 0.75 ^d	16.30 ± 0.71 ^d
102	Leafbase	21.44 ± 0.89 ^e	18.70 ± 0.87 ^c	17.44 ± 1.21 ^c
96				
114	Embryo	17.78 ± 1.21 ^{hi}	13.22 ± 0.82 ⁱ	13.07 ± 0.65 ^{sh}
112	Leafbase	16.30 ± 1.56 ^j	12.11 ± 1.20 ^j	11.22 ± 0.98 ^j
Control				
48				
103	Embryo	11.30 ± 1.37 ^m	9.09 ± 0.78 ^l	8.38 ± 1.05 ^k
95	Leafbase	10.69 ± 0.98 ⁿ	8.78 ± 0.72 ^l	7.03 ± 0.89 ^l
72				
99	Embryo	13.87 ± 1.05 ^l	7.36 ± 0.65 ^m	7.30 ± 0.71 ^l
98	Leafbase	14.53 ± 1.13 ^k	6.71 ± 0.54 ⁿ	6.09 ± 0.69 ^m
96				
96	Embryo	9.02 ± 0.86 ^{op}	4.90 ± 0.43 ^{oP}	3.94 ± 0.94 ⁿ
94	Leafbase	7.48 ± 0.73 ^q	4.04 ± 0.68 ^P	3.56 ± 0.75 ⁿ

Data are expressed as mean values of three independent experiments ± standard deviation; means followed by the same letters are not significant at 5% level.

4.3. Efficacy of SAAT

Under SAAT technique, cultures were subjected to ultrasound and inoculated in the *Agrobacterium* culture. With this technique, embryos and leaf base explants generated 24.19% and 23.22% hygromycin-resistant cultures after 72 hours of the co-cultivation, respectively. Analysis of the transformation efficiency on the embryo cultures showed the efficiency rates of 19.78% and 19.26% using GUS assay and gene specific PCR, respectively. Similarly, leaf base cultures responded to 18.70% and 17.58% transformation efficiency, according to GUS assay and PCR analysis, respectively (Table 1).

4.4. Effect of VIAAT on Transformation Efficiency

Vacuum pressure for 15 minutes at 750 mmHg produced 24.33% and 25% hygromycin-resistant cultures in the embryo and leaf bases, respectively. The embryo explants exhibited 21.96% and 20.74% of the transformation efficiencies when analyzed on the basis of GUS assay and gene specific PCR analysis, respectively. Whereas, the leaf base explants upon GUS and PCR analysis produced 21.85% and 19.04% transformation efficiency, respectively (Table 1).

4.5 Effect of SVAAAT on Transformation Efficiency

Upon the employment of both sonication and vacuum techniques, 21.4% of the cultures from embryos and leaf base survived on hygromycin after 72 hours of co-cultivation. On the basis of histochemical GUS assay and gene specific PCR, the embryo explants exhibited 17.67% and 16.30% transformation efficiency, respectively. Whereas, 18.70% and 17.44% transformation efficiency was conferred by the leaf explants on the basis of GUS histochemical assay and PCR analysis, respectively (Table 1).

5. Discussion

The genetic and physiological status of the cells varies in different explants. However, these features affect the transformation efficiency. Along with explant, the co-cultivation period of the cultures with *Agrobacterium* to assist transfer of T-DNA into plant cell also influences the efficiency of transformation. Determination of competence for transformation at different co-cultivation duration, two explants namely mature embryos and leaf base were utilized. In the current study, the maximum transformation efficiency was obtained for the mature embryo explants at 72 hours of co-cultivation duration. The study results were comparable with those of the transformation studies conducted on oat cultivars-cv. Bajka showing 12.3% transformation efficiency using immature embryo as the explant (12). Co-cultivation duration of three days was sufficient for an efficient transformation. The longer co-cultivation duration resulted in a higher transient expression of the transgene in the soybean (17).

Agrobacterium-mediated transformation assisted with sonication was effective in the transfer of foreign DNA

into cells in a number of species, especially those where the host was recalcitrant to the *Agrobacterium*-mediated transformation (17).

Employment of embryos and leaf base explants produced 24.19% and 23.22% hygromycin-resistant cultures respectively, by the employment of SAAT followed by 72 hours of the co-cultivation period. The improved transformation via SAAT could possibly be due to micro-wounding on the surface of tissue, which aided in the entry of *Agrobacterium* into the plant tissue. The enhanced level of transient expression of the *gus* gene in soybean was observed by treating embryogenic cultures with sonication for 30 seconds (18).

The duration of the co-cultivation period after sonication had enhanced effects on the transformation efficiency.

Creation of vacuum may cause the release of gases from the culture probably through wounds instead of stomata. This phenomenon exposes more plant cells to the *Agrobacterium*, making it more susceptible to the transformation compared with the one where *Agrobacterium* is present in the culture superficially. The current study results suggested that the utilization of vacuum infiltration remarkably increased the transformation efficacy than the other treatments. VIAAT technique is reported for the efficient transformation of rice (19) and wheat (20), as well.

When both sonication and vacuum were applied simultaneously, more than 21% of the hygromycin-resistant cultures were obtained in both explant types. The lower transformation efficiencies of SVIAAT method in comparison with those of the individual effect of processes (i.e., SAAT and VAAT) may be attributed to the damage of the callus's tissue due to sonication followed by vacuum infiltration, hence the plant cells could not recover the incurred damage. The current study results were in congruence with those of the Subramanyam *et al.* (6), where the transformation of banana was conducted using SVAAAT.

6. Conclusions

The current study results demonstrated a promising protocol for *Agrobacterium*-mediated genetic transformation in the oat. To date, this is the only maiden report of the oat transformation using SAAT and VIAAT. Duration of co-cultivation, sonication, and vacuum infiltration positively influence the oat transformation. The developed transformation protocol involves co-cultivation of *A. tumefaciens* along with cultures developed from leaf base under vacuum (15 inch Hg) for 10-15 minutes in the presence of 100 µM acetosyringone. The selection of transformant was conducted on MS medium supplemented with 20 mg/L hygromycin. This transformation protocol could be used efficiently to improve oat considering different agronomical and nutritive traits via genetic engineering.

Acknowledgement

Sharad Tiwari appreciates the research grant from Madhya Pradesh Council of Science & Technology, Bhopal, India.

References

- Ferdman B, Singh G. Persistent Truncus Arteriosus. *Curr Treat Options Cardiovasc Med.* 2003;**5**(5):429-438. [pmid: 12941211](#)
- Jakob K, Zhou F, Paterson AH. Genetic improvement of C4 grasses as cellulosic biofuel feedstocks. Biofuels. New York: Springer; 2009. p. 113-138.
- Takahashi W, Takamizo T. Molecular Breeding of Grasses by Transgenic Approaches for Biofuel Production. In: Ciftci Y, editor. Transgenic Plants-Advances and Limitations. Rijeka: InTech; 2012. p. 91-116.
- Gless C, Lorz H, Jahne-Gartner A. Establishment of a highly efficient regeneration system from leaf base segments of oat (*Avena sativa* L.). *Plant Cell Rep.* 1998;**17**(6-7):441-445. [doi: 10.1007/s002990050422](#) [pmid: 30736616](#)
- Ziemienowicz A. Agrobacterium-mediated plant transformation: Factors, applications and recent advances. *Biocatal Agricult Biotechnol.* 2014;**3**(4):95-102. [doi: 10.1016/j.bcab.2013.10.004](#)
- Subramanyam K, Subramanyam K, Sailaja KV, Srinivasulu M, Lakshmidevi K. Highly efficient Agrobacterium-mediated transformation of banana cv. Rasthali (AAB) via sonication and vacuum infiltration. *Plant Cell Rep.* 2011;**30**(3):425-436. [doi: 10.1007/s00299-010-0996-4](#) [pmid: 21212957](#)
- Mayavan S, Subramanyam K, Arun M, Rajesh M, Kapil Dev G, Sivanandhan G, et al. Agrobacterium tumefaciens-mediated in planta seed transformation strategy in sugarcane. *Plant Cell Rep.* 2013;**32**(10):1557-1574. [doi: 10.1007/s00299-013-1467-5](#) [pmid: 23749098](#)
- Park BJ, Liu Z, Kanno A, Kameya T. Transformation of radish (*Raphanus sativus* L.) via sonication and vacuum infiltration of germinated seeds with Agrobacterium harboring a group 3 LEA gene from *B. napus*. *Plant Cell Rep.* 2005;**24**(8):494-500. [doi: 10.1007/s00299-005-0973-5](#) [pmid: 15843933](#)
- de Oliveira ML, Febres VJ, Costa MG, Moore GA, Otoni WC. High-efficiency Agrobacterium-mediated transformation of citrus via sonication and vacuum infiltration. *Plant Cell Rep.* 2009;**28**(3):387-395. [doi: 10.1007/s00299-008-0646-2](#) [pmid: 19048258](#)
- Bakshi S, Sadhukhan A, Mishra S, Sahoo L. Improved Agrobacterium-mediated transformation of cowpea via sonication and vacuum infiltration. *Plant Cell Rep.* 2011;**30**(12):2281-2292. [doi: 10.1007/s00299-011-1133-8](#) [pmid: 21853337](#)
- Chopra R, Aparna, Saini R. Use of sonication and vacuum infiltration for Agrobacterium – mediated transformation of an Indian lentil (*Lens culinaris* Medik.) cultivar. *Sci Horticult.* 2012;**143**:127-134. [doi: 10.1016/j.scienta.2012.06.019](#)
- Gasparis S, Bregier C, Orczyk W, Nadolska-Orczyk A. Agrobacterium-mediated transformation of oat (*Avena sativa* L.) cultivars via immature embryo and leaf explants. *Plant Cell Rep.* 2008;**27**(11):1721-1729. [doi: 10.1007/s00299-008-0593-y](#) [pmid: 18690445](#)
- Murashige T, Skoog F. A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiol Plant.* 1962;**15**(3):473-497. [doi: 10.1111/j.1399-3054.1962.tb08052.x](#)
- Jefferson RA. Assaying chimeric genes in plants: The GUS gene fusion system. *Plant Mol Biol Rep.* 1987;**5**(4):387-405. [doi: 10.1007/bf02667740](#)
- Saghai-Maroo MA, Soliman KM, Jorgensen RA, Allard RW. Ribosomal DNA spacer-length polymorphisms in barley: mendelian inheritance, chromosomal location, and population dynamics. *Proc Natl Acad Sci U S A.* 1984;**81**(24):8014-8018. [doi: 10.1073/pnas.81.24.8014](#) [pmid: 6096873](#)
- Sapre S, Kumar S, Tiwari S. Transient GUS expression and efficient transformation in tobacco (*Nicotiana tabacum*). *JNKVV Res J.* 2011;**45**(1):96-100.
- Trick HN, Finer JJ. Sonication-assisted Agrobacterium-mediated transformation of soybean [*Glycine max* (L.) Merrill] embryogenic suspension culture tissue. *Plant Cell Rep.* 1998;**17**(6-7):482-488. [doi: 10.1007/s002990050429](#) [pmid: 30736623](#)
- Gaba V, Kathiravan K, Amutha S, Singer S, Xiaodi X, Ananthakrishnan G. The Uses Of Ultrasound In Plant Tissue Culture. In: Dutta Gupta S, Ybaraki Y, editors. Plant tissue culture engineering. Netherlands: Springer; 2006. p. 417-426.
- Dong J, Kharb P, Teng W, Hall TC. Characterization of rice transformed via an Agrobacterium-mediated inflorescence approach. *Mol Breed.* 2001;**7**(3):187-194. [doi: 10.1023/a:1011357709073](#)
- Amoah BK, Wu H, Sparks C, Jones HD. Factors influencing Agrobacterium-mediated transient expression of uidA in wheat inflorescence tissue. *J Exp Bot.* 2001;**52**(358):1135-1142. [doi: 10.1093/jexbot/52.358.1135](#) [pmid: 11432931](#)