

POSTER PRESENTATION

Open Access

# Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis

S Saunier<sup>1\*</sup>, AA Bizet<sup>1</sup>, F Silbermann<sup>1</sup>, E Filhol<sup>1</sup>, T Blisnick<sup>2</sup>, A Henneveu<sup>3</sup>, E Montenont<sup>1</sup>, I Perrault<sup>4</sup>, C Boyle-Feysot<sup>5</sup>, J-M Rozet<sup>4</sup>, P Bastin<sup>2</sup>, HH Arts<sup>6</sup>, C Antignac<sup>1</sup>, AR Benmerah<sup>3</sup>

From First International Cilia in Development and Disease Scientific Conference (2012)  
London, UK. 16-18 May 2012

Nephronophthisis (NPH) is a kidney ciliopathy often associated with extra-renal defects and for which 12 genes (*NPHP1-12*) have been identified. *NPHP1* and *NPHP4* control the ciliary access at the transition zone and the velocity of some intraflagellar transport (IFT)/BBS proteins in *C.elegans*. Recently, in a collaborative effort, we have identified, in families with isolated NPH, mutations in *TTC21B* as well as in *WDR19*, which encode the retrograde IFT-A proteins IFT139 and IFT144, respectively. By ciliome sequencing of 1600 candidate genes from 14 NPH patients followed by Sanger sequencing of a cohort of 52 patients, we have found respectively 8 and 7 patients carrying pathogenic missense mutations in genes coding IFT-A proteins, including *WDR35*, *TTC21B* and *IFT140*, which could partially affect their function. Together, these results indicate that IFT-A are involved in nephronophthisis. Moreover, alteration of cilia length was observed in patient kidney, *Nphp4*<sup>-/-</sup> mice kidney tubules and *NPHP1* or *NPHP4* knockdown IMCD3 cell lines. In these cells, primary cilia present swellings at the distal region accompanied by an accumulation of IFT-B at the base and the tip, similar to what was observed in IFT-A mutants, suggesting a possible alteration of retrograde transport. Additionally, ARL13B, a small GTPase required for proper cilium shape and IFT stability, is absent along the axoneme of *NPHP4*-KD-IMCD cells. By controlling the entry of ciliary components at the transition zone, *NPHP1* and *NPHP4* may modulate IFT-A cargos thus participating in the same pathway (i.e. Wnt/PCP), alteration of which would lead to renal lesions observed in nephronophthisis.

#### Author details

<sup>1</sup>Inserm U983, France. <sup>2</sup>Pasteur Institute, CNRS UMR2581, France. <sup>3</sup>Inserm U1016, Cochin Institute, France. <sup>4</sup>Inserm U781, France. <sup>5</sup>Fondation Imagine, Paris, France. <sup>6</sup>Radboud University Nijmegen Medical Center, the Netherlands.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P99

Cite this article as: Saunier *et al.*: Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis. *Cilia* 2012 **1**(Suppl 1):P99.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [sophie.saunier@inserm.fr](mailto:sophie.saunier@inserm.fr)

<sup>1</sup>Inserm U983, France

Full list of author information is available at the end of the article