

1 **Mechanical forces inducing oxaliplatin resistance in pancreatic cancer can be targeted by**
2 **autophagy inhibition**

3 Maria Kalli^{1*}, Fotios Mpekris¹, Antonia Charalambous¹, Christina Michael¹, Chrystalla Stylianou¹,
4 Chrysovalantis Voutouri¹, Andreas G. Hadjigeorgiou¹, Antonia Papoui¹, John D. Martin² and
5 Triantafyllos Stylianopoulos^{1*}

6

7 ¹ Cancer Biophysics Laboratory, Department of Mechanical and Manufacturing Engineering,
8 University of Cyprus, Nicosia, Cyprus










9 ² Materia Therapeutics, Las Vegas, NV, United States

10 * Correspondence to Maria Kalli and Triantafyllos Stylianopoulos, Email: mkalli02@ucy.ac.cy;
11 tstylian@ucy.ac.cy

12

13

14 **Supplementary Figures**




15

16

17 **Suppl. Fig. 1.** Principal component analysis of RPPA dataset from MIA PaCa-2 cells which subjected to either
 18 zero or four mmHg pressure for 16 hours or 48 hours. **a,** Representation of the RPPA dataset into a 3D space

19 defined by the first three principal components. **b**, Percent of total variance explained by each principal
 20 component. **c, e, g**, Representative immunoblotting showing the levels of LC3BI, LC3BII and p62 in KPC (c),
 21 Pan02 (e) and PANC-1 (g) cells exposed to different levels of compression for 24 hours. Antibody against β -
 22 actin was used as a loading control. Quantification of each antibody compared with loading control was
 23 quantified by ImageJ. Numbers in gray font represent the ratio of LC3BII/I (indicator of autophagic vacuoles)
 24 and p62 (indicator of autophagic degradation) expression normalized to β -actin. **d, f, h** LC3B and p62 mRNA
 25 expression was quantified by qPCR in control and compressed KPC (d), Pan02 (f) and PANC-1 (h) cells. Each
 26 bar indicates the mean fold change \pm SE (n=3 independent samples). Differences were statistically significant
 27 when $P < 0.05$ in two-way ANOVA analysis (Tukey's multiple comparisons test). **i**, Representative
 28 immunoblotting showing the levels of LC3BI, LC3BII and p62 in MIA PaCa-2 cells exposed to 4 mmHg of
 29 compression for 24 hours in the presence or not of 10 μ M HCQ. Antibody against β -actin was used as a
 30 loading control. Quantification of each antibody compared with loading control was quantified by ImageJ.
 31 Numbers in gray font represent the ratio of LC3BII/I and p62 expression normalized to β -actin.

32




33

34 **Suppl. Fig. 2.** Identification of working concentrations of Oxaliplatin and HCQ in MIA PaCa-2 and KPC cells
 35 through IC50 curves and western blotting. **a-b**, MIA PaCa-2 (a) or KPC (b) cells were treated with indicated
 36 concentrations of each drug for 24 hours. Resazurin assay was then performed, and graphs represent the

37 average fold change of % difference \pm SE in Resazurin reduction using as a reference the untreated cells (n=3
38 independent samples). **c**, Representative immunoblotting showing the levels of caspase 3 in KPC cells
39 treated with the indicated concentrations of oxaliplatin for 24 hours. Antibody against β -actin was used as
40 a loading control. **d-e**, Representative immunoblotting showing the levels of LC3BI, LC3BII and p62 in MIA
41 PaCa-2 (d) and KPC (e) cells treated with the indicated concentrations of HCQ for 24 hours. Antibody against
42 GAPDH (d) and β -actin (e) was used as a loading control.

43

44




45

46 **Suppl. Fig. 3. a,** Representative immunoblotting showing LC3BI, LC3BII and p62 protein levels in control and
 47 compressed MIA PaCa-2 cells treated with oxaliplatin and/ or HCQ for 24 hours. Antibody against β -actin
 48 was used as a loading control. Quantification of each antibody compared with loading control was quantified

49 by ImageJ. Numbers in gray font represent the ratio of LC3BII/I and p62 expression in each condition
 50 normalized to β -actin. **b**, Representative immunoblotting showing Caspase 3 and cleaved Caspase 3 protein
 51 levels in control and compressed KPC cells treated with HCQ (10 μ M) combined or not with Oxaliplatin (100
 52 μ g/mL) for 24 hours. Antibody against β -tubulin was used as a loading control. Quantification of each
 53 antibody compared to loading control was quantified by ImageJ. Numbers in gray font represent the average
 54 ratio of cleaved Caspase 3 / β -actin protein expression in each condition (n=2 replicates from 2 independent
 55 experiments). **c**, Gating strategy for Annexin V+ and PI+ cells. The first two rows represent the Fluorescent
 56 Minus One (FMOs) control samples, for the selection of Annexin V+ and PI+ cells. The other two rows
 57 represent the selection strategy of the Annexin V+/PI+ untreated control and compressed cells, respectively.


58



59

60 **Suppl. Fig. 4. a**, Study treatment protocol for the KPC tumor model. Losartan and HCQ were administered
 61 at 40- and 50 mg/kg daily, respectively, via intraperitoneal injection (i.p.) once tumor size reached an

62 average volume of 50 mm³. Two cycles of Oxaliplatin treatment (5 mg/kg) were administered via intravenous
63 injection (i.v.) starting on day 7, allowing Losartan to prime the TME. **b**, Images of MIA PaCa-2 tumors derived
64 from mice of indicated treatments that were excised at day 22. **c-d**, Graphs represent relative mice mass
65 changes in MIA PaCa-2 and KPC bearing mice treated as indicated (n=7 mice per group; n=6 mice in the
66 HCQ+Oxal.-treated group). Data are presented as mean ± SE. Statistical analyses were performed by
67 comparing means between two independent groups using the two-way ANOVA test with Tukey's multiple
68 comparison test. P-values less than 0.05 are denoted on the bar graph. **e**, The effect of HCQ, Losartan or
69 their combination on the relative tumor growth of KPC models (n=6 or 7 mice per group). Data are presented
70 as mean ± SE. Statistical analyses were performed by comparing means between two independent groups
71 using the two-way ANOVA test (Tukey's multiple comparisons test). No statistically significant differences
72 were observed. **f**, The effect of HCQ, Losartan or their combination on the relative tumor perfusion as
73 measured by CEUS (between Day 5 and Day 12) of KPC model (n=6 mice per treatment group). Data are
74 presented as mean ± SE. Statistical analyses were performed by comparing means between two
75 independent groups using the two-way ANOVA test (Tukey's multiple comparisons test). P-values less than
76 0.05 are denoted on the bar graph. Panel (**a**) was created with BioRender.com.



77

78 **Suppl. Fig. 5. a-b**, Quantification of the fraction of area positive for HABP1 (a) and Collagen I (b) stain
 79 normalized to DAPI stain in KPC tumor tissue sections. **c**, Quantification of vessel pericyte coverage as

80 indicated by CD31 and a-SMA overlapping staining normalized to DAPI stain in KPC tumors. **d-f**,
 81 Quantification of vessel open lumen fraction as indicated by CD31 staining in MIA PaCa-2 (d) and KPC (e,f)
 82 tumors of the indicated treatments. **g**, Quantification of a-SMA stain normalized to DAPI stain in KPC tumors.
 83 Data are presented as mean \pm SE (n=2-3 tumors per condition; at least 5 image fields per condition).
 84 Statistical analyses were performed by comparing means between two independent groups using the two-
 85 way ANOVA multiple comparison test (Tukey's correction). P-values less than 0.05 are denoted on the bar
 86 graphs.

87

88

89

Supplementary Table 1. Primer sequences for qPCR				
Target gene	Human		Murine	
	Forward	Reverse	Forward	Reverse
<i>LC3B</i>	GAGAAGCAGCTTCTGTTCTGG	GTGTCCGTTACCAACAGGAAG	CCACGTGAACATGAGCGAG	AACCATGTACAGGAAGCCGT
<i>p62</i>	CCAGAGAGTTCCAGCACA	CCGACTCCATCTGTTCTT	GGACAGCCAGAGGAACAGAT	GGGAGATGTGGGTATAGGGC
<i>actin</i>	CGAGCACAGAGCCTCGCCTTTGCC	TGTCGACGACGAGCGCGCGCA	GACGGCCAGGTCATCACTAT	AAGGAAGGCTGGAAAAGAGC

90

Supplementary Table 2. p-value with multiple way ANOVA		
	Different time points	Different pressures
1 st principal component	0.14	0.096
2 nd principal component	0.39	0.0081*
3 rd principal component	0.48	0.77

91


92

93

94 **Western Blotting Original Images**


95 **Figure 1g.**

96 **Blotting with a-LC3B only**




97

98 **Blotting with anti-p62 only**



99


100 **Blotting with anti-b-actin on p62**



101

102 **Figure 1h**

103 **Blotting with anti-Bcl-2**



104

105 **Blotting with anti-b-actin on Bcl-2**



106

107 **Blotting with Bcl-xL**




109 **Blotting with anti-b-actin on Bcl-xL**




111

112 **Suppl. Fig. 1c**


113 **Blotting with anti-LC3B only**



Blotting with anti-p62 only




115 **Blotting with anti-beta-actin on p62**



116

117 **Suppl. Fig. 1e**

118 **Blotting with anti-LC3B only**



119

120

121

122

123 **Blotting with anti-p62**



124


125 **Blotting with anti-b-actin**



126


127 **Supplementary Fig. 1g**

128 **Blotting with anti-LC3B only**




129

130 **Blotting with anti-p62 only**



131

132 **Blotting with anti-b-actin on p62**




133

134


135 **Suppl. Fig. 1i**

136 **Blotting with anti-LC3B** (*enhanced contrast/brightness for LC3BI*)




137

138 **Blotting with anti-p62**



139


140 **Blotting with anti-b-actin on p62**



141

142 **Figure 2c**

143 **Blotting with anti-LC3B only**




144

145

146 **Blotting with anti-p62 only**


147



148

149 **Blotting with anti-b-actin on p62**

150




151

152


153 **Supplementary Fig. 2d**

154 **Blotting with anti-LC3B**




155

156 **Blotting with anti-p62**



157

158 **Blotting with anti-GAPDH on anti-p62**




159

160


161 **Supplementary Fig.2e**

162 **Blotting with anti-LC3B only**




163

164 **Blotting with anti-p62**



165

166 **Blotting with anti-beta-actin on anti-p62**




167

168


169 **Supplementary Fig. 2c**

170 **Blotting with anti-Caspase 3**



171

172 **Blotting with actin on anti-Caspase 3**



173


174

175 **Figure 3b (2 blots from 2 biological replicates)**


176 **#1**

#2

Compression	-	+	-	+	-	+	-	+
Oxaliplatin	-	-	+	+	-	-	+	+
HCQ	-	-	-	-	+	+	+	+



Compression	-	+	-	+	-	+	-	+
HCQ	-	-	+	+	-	-	+	+
Oxaliplatin	-	-	-	-	+	+	+	+



177

178

179

180

181

182

183


184

185

186

187 **Suppl. Fig. 3a**

188 **Blotting with anti-LC3B only**



189

190 **Blotting with anti-p62 only**



191

192 **Blotting with anti-b-actin on p62**




193

194

195 **Suppl. Fig. 3b**


196 **Blotting with caspase 3**

197



198

199 **Blotting with b-tubulin**



200


201

202

203 **Figure 7c. (MIA PaCa-2)**

204

Blotting with anti-caspase 3




205

206

207


Blotting with anti-PARP1



208

209

Blotting with anti-b-actin on Caspase 3




210

211

212

Blotting with anti-Bcl-2



213

214

Blotting with anti-b-tubulin on Bcl-2




215

216

217

Blotting with anti-Bcl-xL



218

219 **Blotting with anti-b-actin on Bcl-xL**

-55 kDa
-42 kDa

220
221


222 **Figure 7c (KPC)**

223 **Blotting with anti-caspase 3**

-48 kDa
-17 kDa
-14 kDa

224

225 **Blotting with anti- PARP1 (signal was then enhanced using ImageJ for all samples)**

226 
-89 kDa

227 **Blotting with anti-b-tubulin on PARP1**

-55 kDa

228
229
230

231 **Blotting with Bcl-2**

-26 kDa

232

233 **Blotting with anti-b-actin on Bcl-2**

-42 kDa

234

235

236 **Blotting with anti Bcl-xL**

-26 kDa

237

238

Blotting with b-actin on Bcl-xL



239

240

241