



Impact of COVID-19 related home confinement measures on the lifestyle, body weight, and perceived glycemic control of diabetics

Muna Abed Alah^{a,*}, Sami Abdeen^a, Vahe Kehyayan^b, Iheb Bougmiza^{c,d}

^a Community Medicine Department, Hamad Medical Corporation (HMC), Doha, Qatar

^b University of Calgary in Qatar, Doha, Qatar

^c Community Medicine Department, Primary Health Care Corporation (PHCC), Doha, Qatar

^d Community Medicine Department, College of Medicine, Sousse University, Tunisia

ARTICLE INFO

Keywords:

COVID-19
Diabetes
Lifestyle
Glycemic control
Diet
Physical activity

ABSTRACT

Background: People with Diabetes Mellitus are at high risk of encountering COVID-19 infection and are more vulnerable to the negative repercussions of this infection. In this study we aimed to explore the impact of COVID-19 related home confinement measures on physical activity, dietary habits, body weight and perceived glycemic control of adults with type 2 Diabetes Mellitus (T2DM) in Qatar.

Methods: A cross sectional web-based survey was conducted between January and February 2021 targeting adults ≥ 18 years with T2DM.

Results: Over 40% of the participants reported unhealthy dietary changes. We found a significant increase in the sitting/reclining, and screen times. One third of the participants reported weight gain, while one fifth reported poorer glycemic control since the start of home confinement measures. We found that reporting at least one unhealthy dietary change ($p < 0.001$) and being a female ($p = 0.002$) were significantly associated with reporting greater weight gain. Participants who reported five unhealthy dietary behaviours were more than seven times more likely to perceive poorer glycemic control during home confinement measures compared to those who did not report any unhealthy changes (OR: 7.27, 95%CI 1.60–33.5, $p = 0.011$).

Conclusion: Adults with T2DM experienced adverse lifestyle changes during COVID-19 related home confinement measures. Further research is needed to investigate the persistence of such changes in the post pandemic era.

1. Introduction

A growing body of evidence has shown that people with chronic diseases such as Diabetes Mellitus (DM) are at high risk of getting infected with SARS-CoV2 [1–3]. DM is associated with increased mortality and severity of disease in COVID-19 pneumonia [4]. On the other hand, movement restrictions and lockdown measures imposed during the COVID-19 pandemic resulted in deteriorated glycemic control among diabetic patients [5,6]. Reasons behind such deterioration include interruption of medical services with limited access to health-care, treatment delays, low patients' compliance with medications and adopting unhealthy lifestyle habits during the lockdown measures including reduced physical activity especially with closures of gyms and shopping malls, shifting to work from home, and unhealthy dietary habits resulting in weight gain [6–8]. Nutritional and behavioral

interventions are of paramount importance and maybe more important than medical interventions to achieve better glycemic control during the unprecedented crises of the COVID-19 pandemic. Improving the metabolic health is crucial to win the fight against COVID-19 and any other potential future outbreaks [9]. In this study we aimed to explore the impact of COVID-19 related home confinement measures on physical activity, dietary habits, body weight and perceived glycemic control of adults with type 2 Diabetes Mellitus (T2DM) in Qatar.

2. Methods

We conveniently invited adults ≥ 18 years with T2DM to take part in an online web-based survey that was adapted from other validated and reliable questionnaires, which we then translated into three other languages (Arabic, Urdu, and Malayalam) [10,11], between January and

* Corresponding author.

E-mail addresses: MAbedAlah@hamad.qa, muna_talal6@yahoo.com (M. Abed Alah), SAbdeen@hamad.qa (S. Abdeen), vkehaya@ucalgary.ca (V. Kehyayan), mbougmiza@phcc.gov.qa (I. Bougmiza).

<https://doi.org/10.1016/j.metop.2021.100144>

Received 16 October 2021; Received in revised form 26 October 2021; Accepted 26 October 2021

Available online 29 October 2021

2589-9368/© 2021 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

February 2021. The link to the online version of the questionnaire was posted on social media platforms and was circulated through WhatsApp groups (Supplementary file 1).

We collected sociodemographic characteristics of participants such as age, gender, nationality, marital status, employment status, educational level, and comorbidities. The questionnaire also assessed changes in dietary habits (a list of healthy and unhealthy dietary behaviours), physical activity including changes in time spent in exercise (regardless of the type or intensity of exercise), sitting/reclining, and screen times (expressed as hours/day before and during home confinement), weight gain (amount of weight gain during home confinement expressed as weight gain categories (no change, >3 kg, 3–6 kg, 7–10 kg, > 10 kg), the perceived glycemic control (expressed as less controlled, more controlled, or the same) and the reasons behind poorer glycemic control. An ethical approval was obtained from the institutional review board (IRB) of Hamad Medical Corporation (MRC 01-20-838). Taking the survey implied informed consent.

2.1. Data analysis

We used the Statistical Package for the Social Sciences (SPSS) version 26 to analyse the data. Descriptive statistics were presented as frequencies and percentages for categorical variables. The normality of continuous variables was tested using Shapiro Wilk test, and the univariable analysis using nonparametric Mann-Whitney U, and Kruskal Wallis tests was used to compare ordinal and not normally distributed continuous variables between different subgroups. We used the Wilcoxon Signed Rank test to test the differences in screen, sitting/reclining, and exercise times as expressed before and during the home confinement measures. We carried out an ordinal logistic regression to determine the predictors of weight gain considering weight gain categories as a dependent ordinal variable, and a multiple logistic regression analysis to determine the predictors of glycemic control among participants taking into consideration the glycemic control status as binary dependent variable (less controlled vs more controlled or the same combined). Goodness of Fit was assessed using Pearson and Deviance tests for ordinal regression and Hosmer-Lemeshow test for logistic regression. The variables included in the regression models were selected based on clinical relevance (based on risk factors established in the literature), and statistical significance (variables with P values of <0.25 on the univariable analysis were included in the model). The associations between risk factors and outcomes are presented as adjusted odds ratios (ORs) and 95% confidence intervals (95% CIs). P-values of less than 0.05 were considered significant.

3. Results

A total of 171 diabetics completed the survey. The majority (40.4%) were between 35 and 44 years of age, males (68.4%), non-Arabs (90.1%). A total of 148 (86.5%) participants were employed with 78 (52.7%) of them worked from home during home confinement measures. About one quarter (25.7%) of the participants had other comorbidities beside T2DM, most commonly hypertension (Table 1).

The participants reported several unhealthy dietary changes most commonly eating larger quantities of food (46.2%), eating more fat rich food (43.3%), and more sugar and/or sweetened food or beverages (40.4%). On the other hand, 90.6%, and 82.5% of the participants reported healthier changes such as depending more on home cooking and eating more fruits and vegetables, respectively. Upon assessing the changes in physical activity, we found a significant increase in each of the sitting/reclining (1.66 h/day mean increase, 95%CI:1.06–2.26), and screen times (1.83 h/day mean increase, 95%CI:1.31–2.35), with $p < 0.001$. On the other hand, we did not find any significant change in the exercise time. About one third (38.6%) of the participants reported some weight gain since the start of home confinement measures with more than half of them (57.6%) reported an increase of 3–6 kg in their body

Table 1
Sociodemographic profiles and primary outcomes of the participants.

Variable		No (%)
Age	18–34	39 (22.8)
	35–44	69 (40.4)
	45–54	47 (27.5)
	≥55	16 (9.4)
Gender	Male	117 (68.4)
	Female	54 (31.6)
Nationality ^a	Arab	154 (90.1)
	Non-Arab	17 (9.9)
Highest degree of education	No formal education	1 (0.6)
	High school diploma	30 (17.5)
	College or Higher	134 (78.4)
	Vocational training	6 (3.5)
Employment status	Employed	148 (86.5)
	Not employed	23 (13.5)
Marital status	Married	156 (91.2)
	Not married	15 (8.8)
Comorbid disease/s ^b	No	127 (74.3)
	Yes	44 (25.7)
Number of unhealthy dietary changes	None	59 (34.5)
	Yes	
	1	29 (17.0)
	2	28 (16.4)
	3	19 (11.1)
	4	23 (13.5)
Weight gain categories	5	13 (7.6)
	No weight gain	105 (61.4)
	Less than 3 Kg	20 (11.7)
	3–6 Kg	38 (22.2)
	7 Kg or more	8 (4.7)
Perceived glycemic control	Less controlled	36 (21.1)
	The same	97 (56.7)
	More controlled	38 (22.2)
Exercise time difference ^c (Hours/day)		0.00 (1.25)
Sitting/reclining time difference ^c (Hours/day)		1.66 (3.97)
Screen time Difference ^c (Hours/day)		1.83 (3.43)

^a More than 20 different nationalities were reported.

^b The most commonly reported comorbidity was Hypertension.

^c Reported as Mean (SD).

weight. Of all participants, 36 (21.1%), 38 (22.2%), and 97(56.7%) admitted that their blood sugar readings became less controlled, more controlled, and the same respectively. The majority attributed their less controlled readings to less healthy dietary choices, more sedentary behaviours, and more stress during COVID-19 related home confinement measures. As shown in Table 2, participants who reported at least one unhealthy dietary change were 13 times more likely to report greater weight gain than those who did not report such changes (OR 13:10, 95% CI 4.56–37.6, $p < 0.001$). Females were more than 4 times more likely to report greater weight gain compared to males (OR 4.35, 95% CI:1.71–11.05, $p = 0.002$). Similarly, those who were previously active (used to go to the gym before home confinement measures) were more likely to fall in higher weight gain categories (OR: 2.59, 95%CI 1.06–6.36, $p = 0.037$). Upon assessing the predictors of glycemic control, participants who reported five unhealthy dietary behaviours were more than seven times more likely to perceive poorer glycemic control during home confinement measures compared to those who did not report any unhealthy changes (OR: 7.27, 95%CI 1.60–33.5, $p = 0.011$), Table 3.

4. Discussion

This study showed that adults with T2DM experienced some adverse lifestyle changes during the period of home confinement which might result in a rise of diabetes complications and an increase of other non-communicable diseases. The participants reported both favourable and unfavourable dietary changes, reduction in physical activity, increased sedentary behaviours, and weight gain consisting with the results of other studies [12,13]. Persistence of such unhealthy behaviours might lead to obesity and overweight, which in turn can trigger other

Table 2
Determinants and predictors of weight gain during COVID-19 related home confinement among adults with type 2 diabetes mellitus.

Variable		No (%) of participants in each weight gain category			Univariable analysis <i>p</i> -value	Multivariable analysis ^a	
		No increase in weight	Less than 3 kg	3 kg or more		AOR (95%CI)	<i>p</i> -value
Age	18–34	20 (51.3)	4 (10.3)	15 (38.5)	0.046	4.30 (0.61–30.16)	0.143
	35–44	40 (58.0)	8 (11.6)	21 (30.4)		4.51 (0.65–31.34)	0.127
	45–54	31 (66.0)	7 (14.9)	9 (19.1)		5.00 (0.77–32.5)	0.093
	≥55	14 (87.5)	1 (6.3)	1 (6.3)		1 [Reference]	
Gender	Male	84 (71.8)	12 (10.3)	21 (17.9)	<0.001	1 [Reference]	
	Female	21 (38.9)	8 (14.8)	25 (46.3)		4.35 (1.71–11.05)	0.002
Nationality	Arab	6 (35.3)	2 (11.8)	9 (52.9)	0.011	1.81 (0.50–6.54)	0.365
	Non-Arab	99 (64.3)	18 (11.7)	37 (24.0)		1 [Reference]	
Highest degree of education	No formal education	1 (100.0)	0 (0.0)	0 (0.0)	0.259		>0.999
	High school diploma	19 (63.3)	6 (20.0)	5 (16.7)		0.14 (0.15–1.36)	0.091
	College or Higher	83 (61.9)	14 (10.4)	37 (27.6)		0.12 (0.14–1.04)	0.055
	Vocational training	2 (33.3)	0 (0.0)	4 (66.7)		1 [Reference]	
Employment status	Employed	99 (66.9)	14 (9.5)	35 (23.6)	<0.001	1.04 (0.33–3.24)	0.950
	Not employed	6 (26.1)	6 (26.1)	11 (47.8)		1 [Reference]	
Marital status	Married	100 (64.1)	17 (10.9)	39 (25.0)	0.023	0.46 (0.12–1.73)	0.250
	Not married	5 (33.3)	3 (20.0)	7 (46.7)		1 [Reference]	
Comorbid disease/s	No	76 (59.8)	17 (13.4)	34 (26.8)	0.628	0.60 (0.22–1.64)	0.315
	Yes	29 (65.9)	3 (6.8)	12 (27.3)		1 [Reference]	
Being previously active (attending gym regularly)	Yes	13 (40.6)	7 (21.9)	12 (37.5)	0.016	2.59 (1.05–6.36)	0.037
	No	92 (66.2)	13 (9.4)	34 (24.5)		1 [Reference]	
Unhealthy dietary change	No	53 (89.8)	4 (6.8)	2 (3.4)	<0.001	1 [Reference]	
	Yes (at least one unhealthy change)	52 (46.4)	16 (14.3)	44 (39.3)		13.10 (4.56–37.61)	<0.001
Mean time difference for each weight gain category							
Exercise time difference¹ (Hours/day)		0.10	0.30	–0.34	0.074	1.06 (0.78–1.44)	0.719
Sitting/reclining time difference¹ (Hours/day)		1.18	1.55	2.80	0.021	1.13 (1.01–1.26)	0.038
Screen time Difference¹ (Hours/day)		1.55	1.15	2.76	0.063	0.88 (0.77–1.02)	0.087

Abbreviations: AOR adjusted odds ratio; CI, confidence interval.

^a Ordinal logistic regression model was used.

non-communicable diseases. Moreover, obesity increases the risks of infection from SARS-CoV-2 and is associated with more severe infection and higher chances of hospitalization [14]. As shown in Table 2, adopting unhealthy dietary behaviours was found to be a predictor of weight gain in this study supporting the findings of previous studies [15]. Home-based exercise programs can be safe and effective in breaking the cycle of sedentary behaviours adopted during home confinement measures [16]. About one fifth of participants perceived poorer glycaemic control which might render them vulnerable to diabetic complications. Additionally, the poor glycaemic control can lead to immunosuppression and increase the susceptibility to infections including SARS-CoV 2 [17]. In fact, acute or chronic hyperglycaemia in patients with T2DM which frequently accompanies obesity can impair innate and adaptive immunity by disrupting macrophages function, and enhancing viral replication in monocytes, which can result in secondary T cell dysfunction leading to a more severe clinical course of SARS-CoV 2 in a background of pre-existing chronic inflammation that characterizes both T2DM and obesity [14]. The variability in the perceived glycaemic control found in this study as shown in Table 1, reflects the importance of conducting further research with larger sample size that involves actual measurements and follow up of glycaemic parameters such as the glycosylated haemoglobin (HbA1C), and fasting blood glucose to draw a conclusion about the real impact of home confinement on the glycaemic control.

4.1. Strengths and limitations

While this is one of the few studies to assess the perceived impact of COVID-19 related home confinement measures on the lifestyle of diabetics in the Middle East, some limitations must be acknowledged. First, relying on the self-reporting by participants to collect data can

introduce recall bias. Second, our assessment of physical activity was based on changes in exercise time and other sedentary behaviours such as the sitting/reclining and screen times. However, other types of physical activities such as household work might have been affected so the results should be interpreted with caution. Lastly, the small sample size, and the snowballing sampling might have introduced selection bias and compromised the external validity of the study.

5. Conclusion

Further research is needed to explore the persistence of adverse lifestyle changes in the post pandemic era. Diabetics are in need for more intense medical follow ups to restore and optimize their metabolic health after the period of movement restrictions and home confinement.

Funding

None.

CRediT authorship contribution statement

Muna Abed Alah: Conceptualization, Methodology, Data collection and extraction, Interpretation of data, Date analysis, Project administration, Writing – original draft, Final approval of the version to be submitted. **Sami Abdeen:** Methodology, Data collection and extraction, Interpretation of data, Data analysis, Writing – original draft, Final approval of the version to be submitted. **Vahe Kehyayan:** Interpretation of data, Supervision, Writing – review & editing, Final approval of the version to be submitted. **Iheb Bougmiza:** Interpretation of data, Supervision, Writing – review & editing, Final approval of the version to be submitted.

Table 3

Determinants and predictors of the perceived glycemic control during COVID-19 related home confinement among adults with type 2 diabetes mellitus.

Variable		Less Controlled N (%)	Univariable analysis p-value*	Multivariable analysis	
				AOR (95%CI)	p-value
Age	18–34	8 (20.5)	0.229	1.32 (0.11–16.00)	0.826
	35–44	19 (27.5)		3.07 (0.29–32.81)	0.353
	45–54	8 (17.0)		3.22 (0.29–36.36)	0.344
	≥55	1 (6.3)		1 [Reference]	
Gender	Male	20 (17.1)	0.062	0.88 (0.30–2.53)	0.807
	Female	16 (29.6)		1 [Reference]	
Nationality	Arab	30 (19.5)	0.129	0.58 (0.15–2.31)	0.440
	Non-Arab	6 (35.3)		1 [Reference]	
Highest degree of education	No formal education	0 (0.0)	0.834	–	–
	High school diploma	6 (20.0)		–	–
	College or Higher	28 (20.9)		–	–
	Vocational training	2 (33.3)		–	–
Employment status	Employed	27 (18.2)	0.030	0.60 (0.16–2.28)	0.454
	Not employed	9 (39.1)		1 [Reference]	
Marital status	Married	33 (21.2)	>0.999	–	–
	Not married	3 (20.0)		–	–
Comorbid disease/s	No	27 (21.3)	>0.999	–	–
	Yes	9 (20.5)		–	–
Being previously active (attending gym regularly)	Yes	10 (31.3)	0.117	1.72 (0.60–4.97)	0.314
	No	26 (18.7)		1 [Reference]	
Number of unhealthy dietary change	None	7 (11.9)	0.001	1 [Reference]	
	Yes 1	2 (6.9)		0.38 (0.07–2.12)	0.268
	2	7 (25.0)		1.18 (0.30–4.70)	0.815
	3	5 (26.3)		1.09 (0.22–5.41)	0.917
	4	7(30.4)		2.89 (0.70–11.95)	0.143
	5	8 (61.5)		7.27 (1.58–33.53)	0.011
Mean difference for less controlled					
Exercise time difference ¹ (Hours/day)		–0.65	0.001	0.67 (0.43–1.05)	0.080
Sitting/reclining time difference ¹ (Hours/day)		3.06	0.020	0.99 (0.87–1.12)	0.837
Screen time Difference ¹ (Hours/day)		3.75	<0.001	1.16 (0.99–1.34)	0.055

Abbreviations: AOR, adjusted odds ratio; CI, confidence interval.

* Variables with P-value <0.250 were included in the multivariable logistic regression model.

Declaration of competing interest

None.

Acknowledgments

Qatar National Library funded the publication of this article.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.metop.2021.100144>.

Ethical consideration

Ethical approval was obtained from the Institutional Review Board (IRB) of HMC (MRC-01-20-838).

References

- Pranata R, Henrina J, Raffaello WM, Lawrensia S, Huang I. Diabetes and COVID-19: the past, the present, and the future. *Metab Clin Exp* 2021;121. <https://doi.org/10.1016/J.METABOL.2021.154814>.
- Riddle MC, Buse JB, Franks PW, Knowler WC, Ratner RE, Selvin E, et al. COVID-19 in people with diabetes: urgently needed lessons from early reports. *Diabetes Care* 2020;43:1378–81. <https://doi.org/10.2337/DCI20-0024>.
- Jordan RE, Adab P, Cheng KK. Covid-19: risk factors for severe disease and death. *BMJ* 2020;368. <https://doi.org/10.1136/BMJ.M1198>.
- I H, Ma L, R P. Diabetes mellitus is associated with increased mortality and severity of disease in COVID-19 pneumonia - a systematic review, meta-analysis, and meta-regression. *Diabetes Metab Syndr* 2020;14:395–403. <https://doi.org/10.1016/J.DSX.2020.04.018>.
- Eberle C, Stichling S. Impact of COVID-19 lockdown on glycemic control in patients with type 1 and type 2 diabetes mellitus: a systematic review. *Diabetol Metab Syndr* 2021. <https://doi.org/10.1186/S13098-021-00705-9>. 131 2021;13:1–8.
- Tanji Y, Sawada S, Watanabe T, Mita T, Kobayashi Y, Murakami T, et al. Impact of COVID-19 pandemic on glycemic control among outpatients with type 2 diabetes in Japan: a hospital-based survey from a country without lockdown. *Diabetes Res Clin Pract* 2021;176. <https://doi.org/10.1016/J.DIABRES.2021.108840>.
- Karatas S, Yesim T, Beysel S. Impact of lockdown COVID-19 on metabolic control in type 2 diabetes mellitus and healthy people. *Prim Care Diabetes* 2021;15:424–7. <https://doi.org/10.1016/J.PCD.2021.01.003>.
- Alshareef R, Al Zahrani A, Alzahrani A, Ghandoura L. Impact of the COVID-19 lockdown on diabetes patients in Jeddah, Saudi Arabia. *Diabetes Metab Syndr Clin Res Rev* 2020;14:1583–7. <https://doi.org/10.1016/J.DSX.2020.07.051>.
- Means C. Letter to the Editor: mechanisms of increased morbidity and mortality of SARS-CoV-2 infection in individuals with diabetes: what this means for an effective management strategy. *Metab Clin Exp* 2020;108. <https://doi.org/10.1016/J.METABOL.2020.154254>.
- Kumari A, Ranjan P, Vikram NK, Kaur D, Sahu A, Dwivedi SN, et al. A short questionnaire to assess changes in lifestyle-related behaviour during COVID 19 pandemic. *Diabetes Metab Syndr Clin Res Rev* 2020;14:1697–701. <https://doi.org/10.1016/j.dsx.2020.08.020>.
- Chopra S, Ranjan P, Malhotra A, Sahu A, Dwivedi SN, Baitha U, et al. Development and validation of a questionnaire to evaluate the impact of COVID-19 on lifestyle-related behaviours: eating habits, activity and sleep behaviour. *Publ Health Nutr* 2021;24:1275–90. <https://doi.org/10.1017/S1368980020004656>.
- Biamonte E, Pegoraro F, Carrone F, Facchi I, Favacchio G, Lania AG, et al. Weight change and glycemic control in type 2 diabetes patients during COVID-19 pandemic: the lockdown effect. *Endocrine* 2021;72:1. <https://doi.org/10.1007/S12020-021-02739-5>.
- Ruiz-Roso MB, Knott-Torcal C, Matilla-Escalante DC, Garcimartín A, Sampredro-Núñez MA, Dávalos A, et al. COVID-19 lockdown and changes of the dietary pattern and physical activity habits in a cohort of patients with type 2 diabetes mellitus. *Nutrients* 2020;12:1–16. <https://doi.org/10.3390/NU12082327>.
- M D, Gs C, I K, N V, Cm A. Understanding the Co-epidemic of obesity and COVID-19: current evidence, comparison with previous epidemics, mechanisms, and preventive and therapeutic perspectives. *Curr Obes Rep* 2021;10:214–43. <https://doi.org/10.1007/S13679-021-00436-Y>.
- Bennett G, Young E, Butler I, Coe S. The impact of lockdown during the COVID-19 outbreak on dietary habits in various population groups: a scoping review. *Front Nutr* 2021;8:626432. <https://doi.org/10.3389/fnut.2021.626432>.
- Marçal IR, Fernandes B, Viana AA, Ciolac EG. The urgent need for recommending physical activity for the management of diabetes during and beyond COVID-19 outbreak. *Front Endocrinol* 2020;849. <https://doi.org/10.3389/FENDO.2020.584642>.
- Ng V, A E, D K, T S, Gs C, I K, et al. Diabetes mellitus and SARS-CoV-2 infection: pathophysiological mechanisms and implications in management. *Curr Diabetes Rev* 2021;17. <https://doi.org/10.2174/157339981766621010110253>.