



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Sensitivity of wastewater-based epidemiology for detection of SARS-CoV-2 RNA in a low prevalence setting

Joanne Hewitt^{a,*}, Sam Trowsdale^b, Bridget A. Armstrong^c, Joanne R. Chapman^a, Kirsten M. Carter^a, Dawn M. Croucher^a, Cassandra R. Trent^d, Rosemary E. Sim^d, Brent J. Gilpin^c

^a Institute of Environmental Science and Research Ltd, 34 Kenepuru Drive, Porirua, 5240, New Zealand

^b School of Environment, University of Auckland, 23 Symonds Street, Auckland, 1010, New Zealand

^c Institute of Environmental Science and Research Ltd, 27 Creyke Road, Ilam, Christchurch 8041, New Zealand

^d Watcare Services Limited, 52 Aintree Ave, Airport Oaks, Auckland, New Zealand

ARTICLE INFO

Keywords:

Viral shedding
COVID-19
PEG precipitation
RT-qPCR
New Zealand

ABSTRACT

To assist public health responses to COVID-19, wastewater-based epidemiology (WBE) is being utilised internationally to monitor SARS-CoV-2 infections at the community level. However, questions remain regarding the sensitivity of WBE and its use in low prevalence settings. In this study, we estimated the total number of COVID-19 cases required for detection of SARS-CoV-2 RNA in wastewater. To do this, we leveraged a unique situation where, over a 4-month period, all symptomatic and asymptomatic cases, in a population of approximately 120,000, were precisely known and mainly located in a single managed isolation and quarantine facility (MIQF) building. From 9 July to 6 November 2020, 24-hr composite wastewater samples ($n = 113$) were collected daily from the sewer outside the MIQF, and from the municipal wastewater treatment plant (WWTP) located 5 km downstream. New daily COVID-19 cases at the MIQF ranged from 0 to 17, and for most of the study period there were no cases outside the MIQF identified. SARS-CoV-2 RNA was detected in 54.0% (61/113) at the WWTP, compared to 95.6% (108/113) at the MIQF. We used logistic regression to estimate the shedding of SARS-CoV-2 RNA into wastewater based on four infectious shedding models. With a total of 5 and 10 COVID-19 infectious cases per 100,000 population (0.005% and 0.01% prevalence) the predicated probability of SARS-CoV-2 RNA detection at the WWTP was estimated to be 28 and 41%, respectively. When a proportional shedding model was used, this increased to 58% and 87% for 5 and 10 cases, respectively. In other words, when 10 individuals were actively shedding SARS-CoV-2 RNA in a catchment of 100,000 individuals, there was a high likelihood of detecting viral RNA in wastewater. SARS-CoV-2 RNA detections at the WWTP were associated with increasing COVID-19 cases. Our results show that WBE provides a reliable and sensitive platform for detecting infections at the community scale, even when case prevalence is low, and can be of use as an early warning system for community outbreaks.

1. Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has rapidly spread around the world since first detected in late 2019 (World Health Organisation 2020). For many countries, the virus has been extremely difficult to control and continues to circulate in the

community (AJMC Staff 2021). SARS-CoV-2 RNA is shed into wastewater via faeces and respiratory excretions of symptomatic and asymptomatic individuals (Wölfel et al., 2020; Zhang et al., 2020). This means that wastewater-based epidemiology (WBE) can be used to monitor infections at the community scale (e.g. Lodder and Husman 2020, Peccia et al., 2020, Thompson et al., 2020). WBE complements

Abbreviations: WBE, wastewater-based epidemiology; MIQF, managed isolation and quarantine facility; WWTP, wastewater treatment plant; MIF, managed isolation facility; FIPV, feline infectious peritonitis virus; MNV, murine norovirus; CrFK, Crandell-Rees feline kidney; GC, genome copies; PEG, polyethylene glycol; Cq, PCR cycle quantification threshold; LOD, limit of detection; SD, standard deviation; CI, confidence interval; IQR, interquartile range.

* Corresponding author.

E-mail address: joanne.hewitt@esr.cri.nz (J. Hewitt).

<https://doi.org/10.1016/j.watres.2021.118032>

Received 21 August 2021; Received in revised form 24 December 2021; Accepted 30 December 2021

Available online 2 January 2022

0043-1354/© 2021 Elsevier Ltd. All rights reserved.

other COVID-19 surveillance activities (i.e., diagnostic testing of mainly symptomatic individuals) and can operate at a much broader-scale (Bivins et al., 2020; Sims and Kasprzyk-Hordern 2020). WBE has been coupled with modelling to provide advanced warning (4- to 7-days) of an increase in cases (Peccia et al., 2020; Randazzo et al., 2020) and to estimate numbers of asymptomatic and/or undiagnosed cases (Charvarria-Miro et al., 2021; Wu et al., 2020b). WBE has been used at the commercial building or university campus scale to identify unknown cases, and better target public health interventions such as increased screening (Betancourt et al., 2021; Gibas et al., 2021; Harris-Lovett et al., 2021).

However, questions remain regarding the sensitivity of wastewater testing to detect infectious COVID-19 cases in communities where there are no or limited cases (Cao et al., 2020; Foladori et al., 2020; Medema et al., 2020a; Pecson et al., 2021). It is currently unknown how many cases, per unit of population, are required for SARS-CoV-2 RNA to be reliably detected in municipal wastewater. This can best be addressed in a setting with limited and precisely known case numbers. In locations where the virus is widely circulating, undiagnosed and asymptomatic individuals may skew sensitivity estimates towards overestimation. For example, if SARS-CoV-2 RNA was detected in a catchment with five known cases but there were also ten undiagnosed cases, WBE sensitivity might be estimated to be three times higher than reality. Overestimation would potentially be exacerbated when the number of new cases per day, rather than total cases (i.e., incidence rather than prevalence) is used, given that individuals may continue to shed SARS-CoV-2 RNA for weeks following diagnosis (Cevik et al., 2021). While not ideal, daily incidence is the best proxy for total case load in many countries.

In 2020, Aotearoa New Zealand adopted an COVID-19 elimination strategy with the intent of reducing the incidence of cases in the community to zero (e.g., Baker et al., 2020; Jefferies et al., 2020). During the period of our study, all people entering New Zealand were required to enter a government-run hotel-based managed isolation facility (MIF) upon arrival. All people were tested for COVID-19 on at least days 3 and 12, and if COVID-19 symptoms developed (New Zealand Government 2021). Most individuals who tested positive were transferred to a single dedicated managed isolation and quarantine facility (MIQF) in Auckland. In many instances, their close contacts were transferred at the same time. COVID-19 cases detected in the Auckland community, and their close contacts, were transferred from their home to the MIQF. This MIF system for border arrivals, along with a national lockdown and other measures, meant that by the beginning of June 2020, SARS-CoV-2 transmission was effectively eliminated in New Zealand (Baker et al., 2020). As such, most New Zealand COVID-19 cases were contained within one (MIQF) building. The wastewater from this MIQF drained into a wider sewer pipe network with a resident population catchment of about 120,000. There was no sewer input upstream of the MIFQ. By testing paired daily influent wastewater samples from the sewer outside the MIQF and downstream sewer for the presence of SARS-CoV-2 RNA and comparing results to the known number of symptomatic and asymptomatic COVID-19 cases, we were able to calculate the sensitivity of WBE COVID-19 detection. Clinical and wastewater surveillance in the wider Auckland region confirmed that COVID-19 was not in the community during the study period, except for a brief community outbreak described below (Section 3.1). Positive detections in the municipal wastewater were thus solely due to SARS-CoV-2 RNA input from the MIQF, and therefore the data can be used to quantify detection sensitivity. Four shedding models were used to estimate the contribution of COVID-19 cases to the MIQF and downstream WWTP influent wastewater based on modelled infectious status.

2. Materials and methods

2.1. Wastewater sampling

Time-weighted composite samples were collected from two sewer

sites in the same wastewater catchment. At the MIQF, a temporary sewage maintenance access cover and ISCO autosampler (model 3710, Teledyne ISCO, Lincoln, NE) were installed in the sewer on 7 July 2020, and the study ran from 9 July to 6 November 2020. The sampling point included intermittent pumped flow from two separate wastewater sumps on the premises. The sampler was programmed to collect 60 mL wastewater every 10 min per 24-hr period into a large plastic drum without refrigeration. The site was visited daily between 12:30 and 13:00 and, after stirring, a 1 L aliquot of the 24-hr wastewater composite was collected in a plastic bottle. At the inlet to the WWTP, relative to the associated interceptor of the MIQF, an ISCO autosampler (model 4700, Teledyne ISCO) was programmed to collect 200 mL wastewater every 30 min over 24 hr into a large plastic drum with refrigeration. The site was visited daily between 08:00 and 08:30 and, after stirring, a 1 L aliquot of the 24-hr composite was collected in a plastic bottle. All samples were transported overnight on ice to the laboratory, stored at 4 °C until processed, and processed within 18 hr of receipt (Sections 2.2–2.5).

2.2. Quality controls

Feline infectious peritonitis virus (FIPV), an enveloped coronavirus, and murine norovirus (MNV), a non-enveloped virus, were used as external process controls to monitor virus recovery and to assess any RT-qPCR inhibition. Process control recoveries were not used to adjust the SARS-CoV-2 RNA concentration of the samples (as recommended by Kantor et al., 2021).

FIPV (ATCC® VR-990 WSU 79–1146) stocks were prepared, and 50% tissue culture infectious dose (TCID₅₀)/mL determined, in Crandell-Rees Feline Kidney (CrFK) cells (ATCC® CCL-94). MNV stocks were prepared, and plaque forming units (PFU)/mL determined, in RAW 264.7 cells (Hewitt et al., 2009). Working stocks of 10⁶ TCID₅₀/mL FIPV and 10⁶ PFU/mL MNV were prepared from the initial virus preparations by diluting in 2% Eagle's Minimal Essential Media (Gibco, Waltham, MA). Aliquots were stored at –80 °C. Prior to virus concentration (Section 2.3), 2 µL of FIPV and MNV working stocks were added per 100 mL wastewater (e.g., 10 µL each of FIPV and MNV added to 500 mL). For each batch of samples processed, 10 µL each of FIPV and MNV working stocks were also added to 2.5 mL phosphate buffered saline (PBS), pH 7.2 and stored at 4 °C (for up to 18 hr). This was referred to as the FIPV/MNV in PBS process control.

A SARS-CoV-2 positive extraction quality control was prepared from a clinical specimen submitted for diagnostic testing (ESR, Porirua, NZ). The virus suspension was heat inactivated at 56 °C for 30 min, diluted 1/1000 in viral transport media, 200 µL aliquots prepared and stored at –80 °C until required for RNA extraction. The SARS-CoV-2 extraction control was quantified (as described below) and determined as approx. 30,000 genome copies (GC)/mL.

2.3. Viral concentration

Following seeding of FIPV and MNV into wastewater (Section 2.2), viruses from both liquid and solid fractions were recovered, and subsequently concentrated using polyethylene glycol (PEG) precipitation. Wastewater (500 mL) was centrifuged at 10,000 x g for 20 min at 4 °C. The supernatant was transferred to a clean bottle and temporarily stored at 4 °C (maximum 2 hr). The solids were resuspended in a 1:8 (w/v) ratio of glycine buffer (0.05 M glycine, 3% beef extract), and pH adjusted to 9.0. The bottles were sonicated in an ultrasonic water bath for 2 min, and then placed on a horizontal shaker at 200 rpm for 30 min at room temperature. Subsequently, the glycine buffer mixture was centrifuged at 10,000 x g for 20 min at 4 °C and the supernatant combined with the initial, chilled, supernatant. The pH of the entire sample was adjusted to 7.0–7.2. PEG 6000 (Sigma-Aldrich, St. Louis, MO) and sodium chloride (LabServ, Auckland, New Zealand) were added to give final concentrations of 20% and 0.2 M, respectively, and gently mixed at room

temperature until dissolved. The sample was then placed on a horizontal shaker and gently agitated at 70–80 rpm at 4 °C for a minimum of 2 h. The sample was centrifuged at 10,000 x g for 30 min at 4 °C and supernatant discarded. PBS, pH 7.2 was added to resuspend the pellet. The final 'volume before processing' to 'volume after concentration' ratio was approximately 200:1. The final volume was recorded and used in subsequent calculations to determine SARS-CoV-2 RNA GC/L wastewater, and to determine process control recoveries. Wastewater concentrates were stored at 4 °C (maximum 24 hr) until viral RNA was extracted.

2.4 RNA extraction

The High Pure Viral Nucleic Acid Extraction Kit (Roche Molecular Biochemicals Ltd., Mannheim, Germany) was used to extract viral nucleic acid from a 200 µL aliquot of each wastewater concentrate and the FIPV/MNV in PBS process control. One modification was made to the standard protocol; samples were centrifuged at 10,000 x g for 30 s prior to addition to the filter. To monitor for RT-qPCR inhibition and/or poor RNA extraction, a 1:4 dilution of the concentrate and the FIPV/MNV in PBS process control was also prepared prior to RNA extraction. For each batch of RNA extractions, a positive (200 µL pre-aliquoted heat inactivated SARS-CoV-2 suspension) and negative (water) extraction control were used. Viral RNA was stored at 4 °C (maximum 1 hr) or at –80 °C until cDNA synthesis.

2.5. RT-qPCR

Two-step reverse transcription (RT)-qPCR assays were used for the detection of SARS-CoV-2, FIPV and MNV. SARS-CoV-2 RNA was detected using published primers/probe that target the N-gene. Primers and probe sequences are shown in Table S1. (Chinese Center for Disease Control and Prevention, 2020; Dunowska and Ghosh, 2021; Hewitt et al., 2009). For all targets, viral cDNA (5 µL) was first generated from 2.5 µL RNA using Superscript™ III Reverse Transcriptase (Invitrogen) with specific reverse primers (Table S1) according to manufacturer's instructions. RT reactions were carried out at 50 °C for 30 min followed by 95 °C for 4 min. The qPCR consisted of 5 µL cDNA, 10 µL Perfecta ToughMix (Quantabio, Beverly, MA), appropriate primers and probe (Biosearch Technologies, Inc., Teddington, UK) and UltraPure DNase/RNase-free distilled water (DNase/RNase-free water, Invitrogen) to a final volume of 20 µL. Final primer/probe concentrations were 0.5 µM/0.25 µM (N-gene), 0.6 µM/0.15 µM (FIPV) and 0.4 µM/0.2 µM (MNV). The PCR cycling conditions for all three assays were 95 °C for 3 min, followed by 45 cycles of 95 °C for 15 secs and 60 °C for 30 secs. The N-gene qPCR assay was performed using the CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, CA) and the FIPV and MNV assays performed using CFX Connect™ (Bio-Rad). All assays were performed in hard shell, clear well, 96-well PCR plates (Bio-Rad), sealed with Microseal B (Bio-Rad). To increase the sensitivity of detection of SARS-CoV-2 RNA, each viral RNA extract from the wastewater concentrates was tested by RT-qPCR in quadruplicate for both extraction volumes (to give a total of eight reactions per sample). FIPV and MNV RT-qPCR assays were performed in duplicate, for both extraction volumes.

Controls for the N-gene RT-qPCR assay consisted of SARS-CoV-2 (positive) and DNase/RNase-free water (negative) extraction controls, non-template control (RT and qPCR) and reagent blank (qPCR). In addition, 5 µL (total 2×10^2 GC) N-gene synthetic DNA oligonucleotide standard (Dnature Diagnostics & Research Ltd, Gisborne, New Zealand) was included as a positive qPCR control (qPCR only). For FIPV and MNV, a FIPV/MNV in PBS process control (see Section 2.2), RNase/DNase-free water (negative extraction control), non-template water control (RT and qPCR), and reagent blank (qPCR) were used.

PCR cycle quantification threshold (Cq) values were determined in the Bio-Rad CFX Manager™ Software (Bio-Rad) after correcting for

fluorescent drift and manually adjusting to a standardised threshold.

Synthetic SARS-CoV-2 RNA Fragment 1 Standard (National Institute of Standards and Technology, Gaithersburg, MD) was used to generate a RNA standard curve and to determine the limit of detection (LOD). RNA standard dilutions of 10^5 , 10^4 , 10^3 , 10^2 , 50, 25, 10, 5, 2 and 1 GC/reaction were prepared in DNase/RNase free water (Invitrogen) in DNA LoBind tubes (Eppendorf, Hamburg, Germany) to minimise RNA binding. The parameters of the standard curve derived RT-qPCR replicates of 10^5 to 10 GC/reaction were slope, y-intercept, efficiency and R^2 value of –3.580, 37.811, 90.3% and 0.991 respectively.

To determine the 95% LOD (GC/reaction), eight replicates of 10^3 , 10^2 , 50 and 25 GC/reaction, 15 replicates of 10 GC/reaction, and 20 replicates of 5, 2 and 1 GC/reaction were analysed. The RNA standard was detected in all replicates down to 10 GC/reaction (mean PCR Cq value 34.8 ± 0.4), and in 95% of the 5 GC/reaction replicates (mean PCR Cq value 36.1 ± 0.9). One and two GC/reaction were detected in approximately 50% RT-qPCR replicates (mean PCR Cq value 37.2 ± 0.7). The 95% LOD was determined as 5 GC/reaction. This equated to $3.4 \log_{10}$ SARS-CoV-2 GC/L (2500 GC/L) wastewater.

Results were expressed as positive (SARS-CoV-2 RNA detected) if SARS-CoV-2 RNA was detected in at least one RT-qPCR replicate, or negative (SARS-CoV-2 not detected). For the purposes of this study, SARS-CoV-2 RNA quantification was performed for wastewater samples when three or four RT-qPCR replicates were positive and where the mean PCR Cq value ≤ 36.1 (i.e., equivalent to 5 GC/reaction or greater), with a standard deviation (SD) of the PCR Cq values ≤ 1.0 . For these samples, PCR Cq values were converted to GC/reaction using the standard curve, and then expressed as GC/L wastewater using the formula:

$$GC/L = \frac{V_{\text{final}} (\text{mL})}{V_{\text{processed}} (L)} * \frac{V_{\text{eluted}} (\mu\text{L})}{(V_{\text{extracted}} (\mu\text{L}) * V_{\text{PCR}} (\mu\text{L}))} * 1000 * GC / \text{reaction}$$

where V_{final} was the final volume of concentrate (mL), $V_{\text{processed}}$ was the initial volume of wastewater processed (L), V_{eluted} was the volume following RNA extraction (µL), $V_{\text{extracted}}$ was the volume of concentrate extracted (µL), and V_{PCR} was the volume of RNA used in the RT-qPCR reaction (µL). GC/L was then converted to \log_{10} GC/L wastewater.

When three or four replicates were positive, but the calculated mean GC/reaction value was < 5 (i.e., $< \text{LOD}$ per reaction), the SARS-CoV-2 RNA quantification (GC/reaction) was imputed as the LOD (i.e., 5) divided by the square root of 2 (Canales et al., 2018). This equated to $3.2 \log_{10}$ GC/L wastewater for a concentration factor of 200. For samples that did not meet either of the above criteria, results were qualitatively presented as presence/absence detection differentiated by either 1 or 2 positive RT-qPCR replicates.

FIPV and MNV process control recoveries were calculated using the formula,

$$\% \text{ recovery} = \left(\frac{1}{2^{\Delta Cq}} \right) * 100$$

where ΔCq was the difference between the mean PCR Cq of the process control recovered in the sample and mean PCR Cq of the FIPV/MNV in PBS process control. Process control recoveries were considered acceptable for quality purposes if both were greater than 1%. By comparing the mean FIPV and MNV PCR Cq values of undiluted and 1:4 diluted samples, any RT-qPCR inhibition could be evaluated by observing a shift in PCR Cq values between the two samples. The 1:4 dilution would theoretically shift the Cq value by 2 (i.e., a diluted sample gives a higher PCR Cq value when there is no inhibition).

2.6. Epidemiological data

Epidemiological data on notified COVID-19 cases (symptomatic or asymptomatic) were obtained from the EpiSurv Surveillance database (<https://surv.esr.cri.nz/episurv>). These included dates for arrival in the country, transfer to and from the MIQF, onset of illness (if symptomatic),

and collection of clinical samples for diagnostic purposes. Based on these data, we used four models to estimate the shedding of SARS-CoV-2 RNA into wastewater based on modelled infectious status:

- Model 1: Total cases - A count of all COVID-19 cases irrespective of their likely infectious status (conservative overestimate).
- Model 2: Infectious cases - A count of likely infectious COVID-19 cases with a constant shedding pattern beginning three days before reported (symptomatic cases) or imputed (asymptomatic cases, see below) disease onset through to nine days after onset.
- Model 3: Relative infectious cases - A count of likely infectious COVID-19 cases with a proportional shedding pattern based on He et al. (2020) (Fig. 1), with peak shedding on the day of reported (symptomatic cases) or imputed (asymptomatic cases, see below) disease onset, and proportionately less before and afterwards. Due to the calculation, the relative infectious case number was not restricted to whole numbers.
- Model 4: New daily cases - A count of new cases per day (i.e., daily cases), based on reported (symptomatic cases) or imputed (asymptomatic cases, see below) onset of disease.

Estimating the degree of infectiousness for individual cases likely provides more realistic estimates of the number of infected people shedding SARS-CoV-2 RNA to wastewater than the total cases in the catchment (Walsh et al., 2020). We assumed that peak shedding for asymptomatic cases occurred on the date that a positive nasopharyngeal swab sample was collected. We therefore used this sampling date as a proxy “onset date” and that the shedding profile was the same as symptomatic cases.

Often, SARS-CoV-2 RNA contributions made by shedding cases would not have been detected until the following day, due to the residence time of wastewater and the nature of the 24-hr sampling scheme. For example, SARS-CoV-2 RNA shed from a MIQF resident at 2pm on Monday would appear in Tuesday’s 24-hr composite samples collected at the MIQF and WWTP. The date of wastewater samples therefore reflects the contributions from the previous 24 h (i.e., date is the day the




Fig. 1. Model to estimate viral shedding relative to estimated symptom onset date (model 3). Peak shedding at symptom onset date (or estimated onset date), with proportionally less for 3 days prior, and 9 days after, symptom onset date. Model 3 was adapted from He et al. (2020).

24-hr composite sampling ended and collected). Dates relating to occupation and shedding density contribution have been adjusted to reflect the date that wastewater was collected.

2.7. Statistical analysis

Mean (\pm bootstrap 95% confidence intervals (CI)) process control recovery efficiencies were calculated. Logistic regression was used to model the relationship between the number of COVID-19 cases/case-equivalents putatively shedding at the MIQF, as cases per 100,000 population, and detection of SARS-CoV-2 RNA in the paired MIQF and WWTP wastewater samples. The contributions of asymptomatic and symptomatic COVID-19 cases/case-equivalents putatively shedding at the MIQF (per 100,000 population) were regarded in additive models to examine the relative contribution of asymptomatic cases to WWTP wastewater samples. All statistical analyses were performed in R Statistical Computing Software version 4.0.3 (R Core Team 2020) with Tidyverse (Thompson et al., 2020) used to produce graphs.

3. Results

3.1. Cases in managed isolation and quarantine (MIQF)

From 9 July to 6 November 2020, there were a total of 335 COVID-19 cases at the MIQF. Of the 335 total cases, 212 were symptomatic and 123 were asymptomatic. Symptomatic cases had an illness onset which ranged from 30 days before arriving at the MIQF to 3 days after arrival, with an average onset of illness 3.3 days before arrival at the MIQF. Overall, COVID-19 cases stayed at the MIQF from 5 – 52 (mean of 13) days (Fig. 2). During the four-month study, one cluster of 179 community cases, mostly residing in the WWTP catchment being studied, was




Fig. 2. History of each symptomatic (Panel A) and asymptomatic (Panel B) COVID-19 case at the managed isolation and quarantine facility (MIQF). Closed red circles indicate the date reported as symptom onset, open red circles indicate the imputed symptom onset date for asymptomatic cases. Closed black circles indicate the date the case entered the MIQF. Grey lines indicate dates each case was present in MIQF. Black crosses indicate the date the case exited the MIQF.

reported. All cases were rapidly identified (onset of illness 31 July – 11 September 2020), with 155 cases being transferred to the MIQF between 11 August and 11 September (Fig. 2). On 29 August 2020, the total number of cases at the MIQF peaked at 107. This was equivalent to 40 infectious cases (model 2) or 16 relative infectious cases (model 3), and 17 new daily cases (model 4). There were no other known COVID-19 cases outside of the MIQF during this study period. At any one time, there were 5 – 107 total cases (model 1) (equivalent to 4.1 – 89.2 cases per 100,000 population), 2.5 – 50.3 infectious cases (model 2) or 0.2 – 15.1 relative infectious cases (model 3). New daily cases at MIQF (model 4) ranged from a total of 0 – 17 (or 0 – 14.1 per 100,000 population), with an average of two new cases per day. There were between 41 and 121 (average 93) staff, security, or other non-guests onsite at the MIQF each day (none of them cases), giving a total population at the MIQF each day of about 300–500 individuals who potentially contributed to the wastewater.

3.2. Process control recovery and quality controls

Two samples from the MIQF, collected on 19 August and 25 September 2020, had unusual low pH/visual properties with low FIPV recovery (0.3% and 0.5% respectively). This may have been a result of excess acidic commercial cleaning products being used at the MIQF. Data from the four samples (two MIFQ and two WWTP wastewater) collected on these two dates were excluded from further analyses. Data from the remaining 113 paired samples ($n = 226$) were used for subsequent analyses.

Process control recoveries (%) (mean \pm 95% CI) from MIQF wastewater samples were 25.4% (22.5, 28.5) for FIPV and 20.8% (18.1, 23.8) for MNV (Fig. S1a). Recoveries from WWTP wastewater samples were 20.7% (16.7, 25.1) for FIPV and 33.9% (30.2, 38.2) for MNV (Fig. S1a, Table S2, Table S3). In samples diluted 1:4 prior to RNA extraction, FIPV and MNV recovery data showed that, overall, there was no significant RT-qPCR inhibition detected in the MIQF or WWTP wastewater samples (Fig. S1b, Table S2, Table S3). The negative extraction, non-template controls and reagent blanks were negative for all RT-qPCR assays.

3.3. Detection and quantification of SARS-CoV-2 RNA in wastewater

SARS-CoV-2 RNA was detected in 95.6% (108/113) of the MIQF wastewater samples. Mean (\pm SD) PCR Cq values for samples ranged from 26.2 (\pm 0.1) to 37.9 (\pm 0.4) (Fig. 3a). For the majority (84.3%, 91/108) of samples where SARS-CoV-2 RNA was detected, all four RT-qPCR replicates were positive. SARS-CoV-2 RNA concentrations in MIQF

wastewater ranged from 3.2 to 6.0 log₁₀ GC/L. Non-detection of SARS-CoV-2 RNA in MIQF wastewater occurred on 1 July, 25–27 July, and 3 August 2020.

SARS-CoV-2 RNA was detected in 54.0% (61/113) of the WWTP samples. Of the 61 samples where SARS-CoV-2 RNA was detected, 16.4% (10/61) had four positive RT-qPCR replicates. For these ten samples, mean (\pm SD) PCR Cq values ranged from 33.9 (\pm 0.7) to 38.1 (\pm 1.2) (Fig. 3b), with SARS-CoV-2 RNA concentrations that ranged from 3.3 to 3.8 log₁₀ GC/L wastewater from the WWTP influent. For SARS-CoV-2 RNA detections with 1 to 3 positive RT-qPCR replicates, the mean (\pm SD) PCR Cq value was 37.3 \pm 0.8 (range 35.6 to 40.4). Notably, over half of the positives (55.7%, 34/61) had only 1 of 4 positive RT-qPCR replicates (Fig. 3b).

3.4. Relationship between COVID-19 cases and wastewater analysis

As may be expected, quantifiable SARS-CoV-2 RNA concentrations at the WWTP occurred most often when there were the highest number of individuals potentially shedding SARS-CoV-2 RNA (Fig. 4a and 4b). In addition to simply counting COVID-19 total cases, shedding was modelled on the more restricted shedding assumptions described above.

When SARS-CoV-2 RNA was detected in the MIQF wastewater ($n = 108$), there were 6 – 107 total cases (i.e., 5 – 89 per 100,000 population) (model 1). This is equivalent to 3 – 60 (i.e., 2.5 – 50 per 100,000 population) infectious cases (model 2) and 0.2 – 9.1 (i.e., 0.2 – 7.6 per 100,000 population) relative infectious cases (model 3). Non-detections of SARS-CoV-2 RNA in the MIQF wastewater were associated with 5 – 11 total MIQF cases (i.e., 4 – 9 per 100,000 population) compared to at least six total cases when SARS-CoV-2 RNA was detected. The MIQF wastewater samples with SARS-CoV-2 RNA concentrations more than 4.0 log₁₀ GC/L were collected on days when there were more cases (i.e., 7 – 107 cases (i.e., 5.8 – 89.2 per 100,000 population), compared to 0 – 60 cases (i.e., 0 – 50 per 100,000 population) depending on the shedding assumptions (Table 1).

It is clear from the data that detecting SARS-CoV-2 RNA at the WWTP was more likely when there were at least eight total cases (i.e., using model 1) (Table 2). This threshold reduced to four cases using model 2 and one case using model 3. SARS-CoV-2 RNA detections at the WWTP, particularly those with three or four RT-qPCR replicates, were associated with an increase in the number of total (model 1), infectious (model 2), relative infectious (model 3), or new daily (model 4) cases in the catchment (Table 2).

Logistic regression models were fitted to data excluding and including the period of the community COVID-19 cluster for each




Fig. 3. Number of RT-qPCR replicates positive for SARS-CoV-2 RNA in the MIQF (Panel A) and WWTP wastewater (Panel B) samples. Mean RT-qPCR Cq values are shown with an inverted y-axis (as lower Cq values represent higher viral concentrations).




Fig. 4. Detection of SARS-CoV-2 RNA in wastewater collected at the MIQF (Panel A) and at WWTP (Panel B). Red triangles indicate MIQF positive samples, yellow triangles indicate MIQF negative samples, red circles indicate WWTP positive samples, and yellow circles indicate WWTP negative samples. On the y-axis, \log_{10} genome copies/L of 1 or 2 have been imputed to represent the 1 or 2 positive RT-qPCR replicates. Blue shading indicates COVID-19 cases at the MIQF on given day (model 1), purple shading estimated infectious cases (model 2), beige shading estimated relative infectious cases (model 3), and green shading new daily cases (model 4). The period when community cases were identified (28 July to 11 September 2020) is indicated by orange bar on x-axis.

Table 1

Estimated COVID-19 case equivalents at the MIQF when SARS-CoV-2 RNA was either not detected, detected at $\leq 4.0 \log_{10}$ GC/L or detected at $> 4.0 \log_{10}$ GC/L wastewater collected at the MIQF ($n = 113$).

SARS-CoV-2 RNA in MIQF wastewater	No of samples	Average number of COVID-19 cases at the MIQF (Minimum - Maximum)			
		Total cases Model 1	Infectious cases Model 2	Relative infectious cases Model 3	New daily cases Model 4
Not detected	5	8.2 (5 - 11)	5.2 (3 - 7)	1.7 (0.8 - 3.0)	0.8 (0 - 2)
Detected ($\leq 4.0 \log_{10}$ GC/L)	56	21.1 (6 - 83)	12.8 (3 - 43)	4.2 (0.2 - 12.2)	1.8 (0 - 9)
Detected ($> 4.0 \log_{10}$ GC/L)	52	49.5 (7 - 107)	27.4 (4 - 60)	8.0 (2.0 - 17.7)	3.9 (0 - 17)
Total	113	33.6 (5 - 107)	19.2 (3 - 60)	5.8 (0.2 - 17.7)	2.7 (0 - 17)

shedding model (Fig. 5). When the data during the community cluster were excluded, increases in relative infectious cases and new daily cases, was not significantly associated with SARS-CoV-2 RNA detection at the WWTP ($df = 67$, model 3: $p = 0.113$; model 4: $p = 0.372$). Case load and

Table 2

Estimated COVID-19 case equivalents at the MIQF when SARS-CoV-2 RNA was either not detected, detected in 1 or 2 RT-qPCR replicates, or detected in 3 or 4 RT-qPCR replicates at the WWTP ($n = 113$).

SARS-CoV-2 RNA in WWTP wastewater	No of samples	Average number of COVID-19 cases at the MIQF (Minimum - Maximum)			
		Total cases Model 1	Infectious cases Model 2	Relative infectious cases Model 3	New daily cases Model 4
Not detected	52	20.1 (5 - 71)	12.6 (3 - 28)	4.0 (0.2 - 15.0)	1.8 (0 - 11)
Detected: 1 or 2 replicates	42	32.1 (8 - 83)	19.6 (4 - 57)	6.0 (1.0 - 15.9)	2.5 (0 - 9)
Detected: 3 or 4 replicates	19	75.0 (18 - 107)	37.2 (10 - 60)	10.3 (2.0 - 17.7)	5.7 (0 - 17)

infectious cases were significantly associated with an increase in SARS-CoV-2 RNA detection at the WWTP ($df = 67$, model 1: $p = 0.012$, model 2: $p = 0.033$) (Table S2).

However, when the community cluster data was included, increasing cases for all models were associated with an increased probability of detection at the WWTP ($df = 112$, models 1, 2 & 3: $p < 0.001$, model 4: p




Fig. 5. Logistic regression of the probability of detecting SARS-CoV-2 RNA at the WWTP, estimated using A) MIQF infectious cases (model 2) for the whole study period; B) MIQF relative infectious cases (model 3) for the whole study period; C) New daily cases (model 4) for the whole study period; D) MIQF infectious cases (model 2) excluding the community cluster data; E) MIQF relative infectious cases (model 3) excluding the community cluster data; F) New daily cases (model 4) excluding the community cluster data.

= 0.003) (Table S2). The proportion of symptomatic to asymptomatic cases at the MIQF varied with SARS-CoV-2 RNA detections at the WWTP (Fig. 6). The interquartile range (IQR) of ratios was higher in the 19 WWTP samples with three or four positive RT-qPCR replicates (IQR = 2.5 – 5.8 symptomatic cases to every asymptomatic case) than for the 42 WWTP samples with only one or two positive RT-qPCR replicates (IQR = 0.78 – 2.4 symptomatic cases to every asymptomatic case) and for the 52 WWTP samples where SARS-CoV-2 RNA was not detected (IQR =

0.71 – 1.8 symptomatic cases to every asymptomatic case).

Logistic regression models with terms for symptomatic and asymptomatic case contributions for infectious cases (model 2) and relative infectious cases (model 3), and new daily cases (model 4) were significantly associated with an increase in probability of detection of SARS-CoV-2 RNA at the WWTP for symptomatic cases (df = 112, model 2 & 3: $p = 0.002$, model 4: $p = 0.008$) (Supp. Table S2). However, asymptomatic cases were not significantly associated with an increase in probability of detection of SARS-CoV-2 RNA at the WWTP for any shedding model (df = 112, model 2: $p = 0.488$, model 3: $p = 0.661$, model 4: $p = 0.461$) (Supp. Table S2). This indicates that asymptomatic cases are likely to be shedding less detectable SARS-CoV-2 RNA into wastewater.

4. Discussion

In this study, by comparing actual case counts with models estimating number of infectious individuals, we have shown that SARS-CoV-2 RNA can be detected in wastewater when relatively few infectious individuals are contributing to the sample. For example, detection probability was 87% when the infectious case prevalence (model 3) was 0.01% (i.e., 10 infectious cases per 100,000 population). While SARS-CoV-2 RNA can be detected when there is only one infectious case in 100,000 population, the predicted probability was lower at 29%. Hence, in communities with very low (i.e., single/few) cases of COVID-19, false negative results may occur. This false negative result may occur at the sample scale, because the amount of viral RNA within the sample means that it cannot be detected by the RT-qPCR. This may be addressed in part




Fig. 6. The proportion of symptomatic cases to asymptomatic cases (model 1) at the MIQF by the number of RT-qPCR replicates positive for SARS-CoV-2 RNA at the WWTP.

by increasing the concentration factor, increasing the amount of template tested and/or increasing the number of technical replicates tested. Alternatively, the sample could be a true negative (i.e., no SARS-CoV-2 RNA is present in the sample) but be a false negative at the community scale because the sample collected did not capture the intermittent and low levels of SARS-CoV-2 RNA in the wastewater system. Even if sampling could somehow test all the wastewater in a community, shedding from infectious individuals whose residence is served by a septic tank or who are otherwise not connected to the wastewater system will also not be detected. These factors must be considered when interpreting and communicating results to public health authorities.

WBE sensitivity estimates

Previous estimates of the sensitivity of SARS-CoV-2 RNA detection in wastewater have varied widely. Differences in sensitivity between studies may be caused by variation in the number of asymptomatic and pre-symptomatic individuals, and COVID-19 testing rates, in different localities. As the number of undetected COVID-19 cases contributing to wastewater increases, precision of sensitivity estimates reduces. Most previous studies were based on reported cases, which are assumed to underrepresent actual infections in communities with widespread community transmission. Further, the various viral concentration and detection methods used across studies will affect recovery efficiency and limits of detection. Additionally, few studies have considered the effect that variable shedding patterns over the course of an infection will have on estimates (Crank et al., 2022). Hart et al. (2020) suggest that, depending on local conditions, a single COVID-19 case would be theoretically detectable using WBE from a catchment of 100 (1% infection rate, least sensitive) to 2000,000 (0.00005% infection rate, most sensitive) people, based on excretion rates, wastewater flow rates and detection limit of 10 GC/mL (i.e., 10^4 GC/L) wastewater. Our data tested these estimates and found that detection with 0.00005% prevalence would be highly unlikely, whereas detection with a 1% prevalence would be highly likely. Indeed, detection with prevalence of 0.005% was possible, but was more reliable (90–100%) at a 0.03% or higher prevalence (i.e., 30 total cases in 100,000 population). The sensitivity reported in our study is similar to that reported by Hata et al. (2021) who concluded that, while there was a possibility that SARS-CoV-2 RNA could be detected in wastewater when the number of cases was less than 1 per 100,000 (0.001%), detections were more likely when COVID-19 cases exceeded 10 per 100,000 population (0.01%). Medema et al. (2020b) reported somewhat higher sensitivity, with positive detections in wastewater occurring when the prevalence of COVID-19 cases was higher than 5 cases per 100,000 population (0.005%) (Table 3). Other

studies (e.g., Baldovin et al., 2021, Hong et al., 2021) have reported lower sensitivity.

Sensitivity estimates will also be influenced by factors affecting RNA decay, such as distance travelled and time spent in the sewer network, composition of the wastewater, collection and transport methods, and laboratory handling and extraction methods (Kantor et al., 2021). To date, studies seeking to determine the rate of SARS-CoV-2 decay under real-world scenarios are largely lacking (although see (Bivins et al., 2020), Carducci et al., 2020, Wurtzer et al., 2021).

Selecting an appropriate method for recovering viral genetic material is crucial to the success of WBE studies. Much research effort has recently focused on finding optimal methods for SARS-CoV-2 RNA recovery. The method used for this study, whereby viral particles were concentrated from the both the solid and liquid phases prior to PEG (20%) precipitation, compares well or better (mean recovery efficiencies ranging from ~20–35%, Figure S1) than other methods, including those based on PEG-precipitation, most often with no phase separation (Torlii et al. 2022; Philo et al., 2021). Other studies (e.g. Barril et al., 2021, Flood et al., 2021, Salvo et al., 2021) have demonstrated that the PEG precipitation-based methods can outperform other common methods such as filtration and flocculation, but that specifics of the methodology can have a large influence on recovery efficiencies. The methods described here provide an alternative approach to those currently deployed in WBE studies of SARS-CoV-2, particularly when sensitivity is paramount, such as in low case prevalence settings.

Use of shedding models

The amount of shedding by individuals is known to vary person to person, and over time (Hoffmann and Alsing 2021). In this study, some individuals stayed at the MIQF for over 30 days (average 13 days), during which time the amount of SARS-CoV-2 RNA shed via faeces and respiratory excretions likely varied from high to undetectable levels. Using total COVID-19 cases as an indicator of SARS-CoV-2 RNA shedding per person provides a conservative estimate of the sensitivity of detecting SARS-CoV-2 RNA in wastewater. This is because individuals towards the end of their infectious period, and potentially shedding at low levels, are counted equally with people at the start of their infectious period. Both models 2 and 3 assumed that viral shedding was minimal nine days after illness onset (Cevik et al., 2021). While SARS-CoV-2 RNA continues to be shed for prolonged periods in some individuals (Gupta et al., 2020), which could have resulted in a sensitivity overestimate in our study, the impact of this is likely to be minimal, as almost all cases left the MIQF (and most presumably from the larger WWTP catchment) within 12 days of reported illness onset. Models 3 and 4 provided similar estimates of detection probability. This similarity likely reflects that detection (and movement into MIQF) for most cases occurred in the early stages of their infection. Unsurprisingly, the probability of SARS-CoV-2 RNA detection at the WWTP increased as case numbers rose. This was evident during the period of community cases within the study catchment, when SARS-CoV-2 RNA was detected with quantifiable concentrations at the WWTP. Over the course of this study, the number of new daily cases (model 4) ranged from 0 to 17, with an average of two new cases per day. Using new daily cases to estimate a positive detection of SARS-CoV-2 RNA therefore had little support beyond 10 new daily cases due to the low case prevalence during the study period (Fig. 5c).

Shedding by asymptomatic cases

Our data suggest that symptomatic individuals may shed a higher amount of SARS-CoV-2 RNA into wastewater than asymptomatic individuals (Fig. 6). Wastewater testing is potentially one of the best ways to detect asymptomatic cases in a community, as they are otherwise unlikely to be tested unless they are close contacts of symptomatic cases. During our study, asymptomatic case load was not significantly associated with increased SARS-CoV-2 RNA detection in wastewater for any

Table 3

Predicted probabilities of SARS-CoV-2 RNA detection at the WWTP by selected equivalent case counts (per 100,000 population) from logistic regression models fitted with community cluster data.

COVID-19 case count per 100,000 (prevalence)	Expected probability of positive tests in WWTP wastewater (95% Confidence interval)			
	Total cases (model 1)	Infectious cases (model 2)	Relative infectious cases (model 3)	New daily cases (model 4)
1 (0.001%)	24% (13 – 38)	19% (10 – 34)	29% (18 – 43)	46% (36 – 56)
5 (0.005%)	28% (17 – 42)	28% (17 – 41)	58% (47 – 68)	74% (57 – 86)
10 (0.01%)	34% (23 – 46)	41% (30 – 52)	87% (70 – 95)	93% (70 – 99)
20 (0.02%)	47% (37 – 58)	69% (56 – 80)	99% (92 – 100) *	–
40 (0.04%)	74% (59 – 85)	96% (83 – 99)	100% (99 – 100) *	–

* Extrapolation beyond available data.

shedding models. It is possible that asymptomatic cases in our study may have been tested longer after infection than symptomatic cases, many of whom were tested immediately upon developing symptoms. Studies specifically investigating differences in shedding between symptomatic and asymptomatic individuals, particularly in faeces, would be valuable. We are aware of only a few studies where SARS-CoV-2 RNA has been detected in the faeces of asymptomatic individuals (e.g., Li et al., 2020, Yuan et al., 2021).

Near-source tracking

As part of this study, we have showed that at a building level, wastewater analysis has a very high probability of detecting SARS-CoV-2 RNA shed from COVID-19 cases, even when total infectious COVID-19 cases are few. This wastewater 'near-source tracking' has been used to target surveillance where it is most needed (Medema et al., 2020a) such as in university campuses and schools (Crowe et al., 2021), elderly care facilities and hospitals (Davó et al., 2021; Spurbek et al., 2021), and households (Döhla et al., 2020). Similar to the concept of the MIQF used in our study, Hong et al. (2021) used a semi-controlled community, namely a hospital with COVID-19 cases in Jeddah, Saudi Arabia, to examine the detection sensitivity of WBE. They calculated sensitivity of ≥ 253 positive cases per 10,000 population (2.5% prevalence) for reliable detection in wastewater collected via a grab-sample. Baldovin et al. (2021) sampled localised urban wastewater lines in Padua, Italy during a time of high COVID-19 case numbers. The authors calculated a detection sensitivity of 1 case per 531 hospitalised inhabitants (0.19% prevalence) for reliable detection.

Sampling frequency

Daily sampling, as undertaken in this study, provides the best understanding of potential viral RNA shedding in the community and the earliest response opportunity. Based on our study, detecting 0.8 to 4.2 cases in 100,000 is possible, although shedding dynamics, and wastewater factors such as dilution and distance of cases from the sampling site, sewer layout, flow rates and other variables means that the likelihood of detection in a single wastewater sample might only be 20–50%. As daily testing may not be practical, investigations around the optimal frequency of wastewater sampling in a low prevalence setting would be a fruitful area of research.

Conclusions

By quantifying the detection limit of SARS-CoV-2 RNA in wastewater where the number of contributing symptomatic and asymptomatic COVID-19 cases was known, and through modelling viral RNA shedding, this study demonstrates that even with low numbers of infectious individuals (10 – 31) in a WWTP catchment there is a high probability (>90% depending on shedding assumption model) of detecting SARS-CoV-2 RNA in wastewater. As such, wastewater can be used as both an early warning system of community outbreaks and for longer term surveillance. However, it is important to carefully consider how cases are quantified/modelled, and the relative contributions of symptomatic and asymptomatic individuals when estimating sensitivity and interpreting results of WBE studies of SARS-CoV-2.

Funding information

This work was supported by the Ministry of Business and Innovation COVID-19 Innovation Acceleration Fund, grant number CIAF-0840. The funders had no role in study design; collection, analysis, and interpretation of data; manuscript writing or the submission process.

Research data

The raw data underpinning this paper is presented as Supplementary Tables S3 and S5.

Declaration of Competing Interest

None.

Acknowledgements

The authors would like to thank residents and staff at the MIQF and WWTP catchment for their wastewater inputs, and staff who collected samples and helped with logistics. Carmen Vettori, Donna Marie Warren and Mary Brunsdon (ESR) assisted with sample accessioning and laboratory support services, Alex Eustace, Amy Bradshaw and Tammy Waters (ESR) helped with sample processing and/or analysis, Beverly Horn and Jenny Ralston (ESR) provided assistance with statistical analysis and graphics respectively. Magda Dunowska (Massey University, New Zealand) provided the feline infectious peritonitis virus (FIPV), ATCC VR-990 strain WSU 79–1146 and designed the primers. Pradip Gyawali (ESR) designed the FIPV probe. Murine norovirus was kindly provided by Prof Herbert Virgin (Washington University School of Medicine, MO, US). Watershed Engineering Ltd conducted mapping of the sewershed and developed estimates of resident population. Liza Lopez (ESR) extracted case data from EpiSurv - a secure web-based application based on the Surveillance Information New Zealand (SurvINZ) architecture. The authors thank the team that maintain EpiSurv and those who have uploaded case data. We also appreciate comments on an earlier version from Erasmus Smit, Gerard Sonder and Phil Carter (ESR).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.watres.2021.118032.

References

- AJMC Staff. (2021) A Timeline of COVID-19 Developments in 2020. <https://www.ajmc.com/view/a-timeline-of-covid19-developments-in-2020>. Accessed on 30 March 2021.
- Baker, M.G., Wilson, N., Anglemeyer, A., 2020. Successful elimination of Covid-19 transmission in New Zealand. *N. Engl. J. Med.* 383 (8), e56 <https://doi.org/10.1056/NEJMc2025203>.
- Baldovin, T., Amoruso, I., Fonzo, M., Buja, A., Baldo, V., Cocchio, S., Bertonecello, C., 2021. SARS-CoV-2 RNA detection and persistence in wastewater samples: an experimental network for COVID-19 environmental surveillance in Padua, Veneto Region (NE Italy). *Sci. Total Environ.* 760, 143329 <https://doi.org/10.1016/j.scitotenv.2020.143329>.
- Barril, P.A., Pianciola, L.A., Mazzeo, M., Ousset, M.J., Jaureguiberry, M.Y., Alessandrello, M., Sánchez, G., Oteiza, J.M., 2021. Evaluation of viral concentration methods for SARS-CoV-2 recovery from wastewaters. *Sci. Total Environ.* 756, 144105 <https://doi.org/10.1016/j.scitotenv.2020.144105>.
- Betancourt, W.Q., Schmitz, B.W., Innes, G.K., Prasek, S.M., Pogreba Brown, K.M., Stark, E.R., Foster, A.R., Sprissler, R.S., Harris, D.T., Sherchan, S.P., Gerba, C.P., Pepper, I.L., 2021. COVID-19 containment on a college campus via wastewater-based epidemiology, targeted clinical testing and an intervention. *Sci. Total Environ.* 779, 146408 <https://doi.org/10.1016/j.scitotenv.2021.146408>.
- Bivins, A., Greaves, J., Fischer, R., Yinda, K.C., Ahmed, W., Kitajima, M., Munster, V.J., Bibby, K., 2020a. Persistence of SARS-CoV-2 in water and wastewater. *Environ. Sci. Technol. Lett.* 7 (12), 937–942 <https://doi.org/10.1021/acs.estlett.0c00730>.
- Bivins, A., North, D., Ahmad, A., Ahmed, W., Alm, E., Been, F., Bhattacharya, P., Bijlsma, L., Boehm, A.B., Brown, J., Buttiglieri, G., Calabro, V., Carducci, A., Castiglioni, S., Gurol, Z.C., Chakraborty, S., Costa, F., Curcio, S., de los Reyes, F.L., Vela, J.D., Farkas, K., Fernandez-Casi, X., Gerba, C., Gerrity, D., Girones, R., Gonzalez, R., Haramoto, E., Harris, A., Holden, P.A., Islam, M.T., Jones, D.L., Kasprzyk-Hordern, B., Kitajima, M., Kotlarz, N., Kumar, M., Kuroda, K., La Rosa, G., Malpei, F., Mauts, M., McLellan, S.L., Medema, G., Meschke, J.S., Mueller, J., Newton, R.J., Nilsson, D., Noble, R.T., van Nuijs, A., Peccia, J., Perkins, T.A., Pickering, A.J., Rose, J., Sanchez, G., Smith, A., Stadler, L., Stauber, C., Thomas, K., van der Voorn, T., Wigginton, K., Zhu, K., Bibby, K., 2020b. Wastewater-based epidemiology: global collaborative to maximize contributions in the fight against COVID-19. *Environ. Sci. Technol.* 54 (13), 7754–7757 <https://doi.org/10.1021/acs.est.0c02388>.

- Canales, R.A., Wilson, A.M., Pearce-Walker, J.I., Verhoughstraete, M.P., Reynolds, K.A., 2018. Methods for handling left-censored data in quantitative microbial risk assessment. *Appl. Environ. Microbiol.* 84 (20) <https://doi.org/10.1128/AEM.01203-18>.
- Cao, B., Gu, A.Z., Hong, P.-Y., Ivanek, R., Li, B., Wang, A., Wu, J., 2020. Editorial perspective: viruses in wastewater: wading into the knowns and unknowns. *Environ. Res.* 196, 110255 [doi:10.1016/j.envres.2020.110255](https://doi.org/10.1016/j.envres.2020.110255).
- Carducci, A., Federigi, I., Liu, D., Thompson, J.R., Verani, M., 2020. Making Waves: coronavirus detection, presence and persistence in the water environment: state of the art and knowledge needs for public health. *Water Res.* 179, 115907 <https://doi.org/10.1016/j.watres.2020.115907>.
- Cevik, M., Tate, M., Lloyd, O., Maraolo, A.E., Schafers, J., Ho, A., 2021. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *The Lancet Microbe* 2 (1), e13–e22 [https://doi.org/10.1016/S2666-5247\(20\)30172-5](https://doi.org/10.1016/S2666-5247(20)30172-5).
- Chavarría-Miro, G., Anfruns-Estrada, E., Martínez-Velazquez, A., Vazquez-Portero, M., Guix, S., Paraira, M., Galofre, B., Sanchez, G., Pinto, R.M., Bosch, A., 2021. Time evolution of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2) in wastewater during the first pandemic wave of COVID-19 in the metropolitan area of Barcelona, Spain. *Appl. Environ. Microbiol.* 87 (7) <https://doi.org/10.1128/AEM.02750-20>.
- Chinese Center for Disease Control and Prevention, 2020. Technical guidelines for COVID-19 laboratory testing. *China CDC Weekly* 2 (19), 332–336 <https://doi.org/10.46234/ccdcw2020.085>.
- Crank, K., Chen, W., Bivins, S., Lowry, A., Bibby, K., 2022. Contribution of SARS-CoV-2 RNA shedding routes to RNA loads in wastewater. *Sci. Total Environ.* 806 (2), 150376 <https://doi.org/10.1016/j.scitotenv.2021.150376>.
- Crowe, J., Schnaubelt, A.T., Schmidt-Bonne, S., Angell, K., Bai, J., Eske, T., Nicklin, M., Pratt, C., White, B., Crotts-Hannibal, B., Staffend, N., Herrera, V., Cobb, J., Conner, J., Carstens, J., Tempero, J., Bouda, L., Ray, M., Lawler, J.V., Campbell, W. S., Lowe, J.-M., Santarpia, J., Bartelt-Hunt, S., Wiley, M., Brett-Major, D., Logan, C., Broadhurst, M.J., 2021. Pilot program for test-based SARS-CoV-2 screening and environmental monitoring in an urban public school district. *medRxiv*, 2021.2004.2014.21255036. <https://doi.org/10.1101/2021.04.14.21255036>.
- Davó, L., Seguí, R., Botija, P., Beltrán, M.J., Albert, E., Torres, I., López-Fernández, P.A., Ortí, R., Maestre, J.F., Sánchez, G., Navarro, D., 2021. Early detection of SARS-CoV-2 infection cases or outbreaks at nursing homes by targeted wastewater tracking. *Clin. Microbiol. Infect.* 27 (7), 1016–1063 <https://doi.org/10.1016/j.cmi.2021.02.003>.
- Döhla, M., Wilbring, G., Schulte, B., Kümmerer, B.M., Diegmann, C., Sib, E., Richter, E., Haag, A., Engelhart, S., Eis-Hübing, A.M., Exner, M., Streeck, H., Schmuthausen, R. M., 2020. SARS-CoV-2 in environmental samples of quarantined households. *medRxiv*, 2020.2005.2028.20114041. <https://doi.org/10.1101/2020.05.28.20114041>.
- Dunowska, M., Ghosh, S., 2021. In vitro effects of doxycycline on replication of feline coronavirus. *Pathogens* 10 (3), 312 <https://doi.org/10.3390/pathogens10030312>.
- Flood, M.T., D'Souza, N., Rose, J.B., Aw, T.G., 2021. Methods evaluation for rapid concentration and quantification of SARS-CoV-2 in raw wastewater using droplet digital and quantitative RT-PCR. *Food Environ. Virol.* 13, 303–315 <https://doi.org/10.1007/s12560-021-09488-8>.
- Foladori, P., Cutrupi, F., Segata, N., Manara, S., Pinto, F., Malpei, F., Bruni, L., La Rosa, G., 2020. SARS-CoV-2 from faeces to wastewater treatment: what do we know? A review. *Sci. Total Environ.* 743, 140444 <https://doi.org/10.1016/j.scitotenv.2020.140444>.
- Gibas, C., Lambirth, K., Mittal, N., Juel, M.A.I., Barua, V.B., Brazell, L.R., Hinton, K., Lontai, J., Stark, N., Young, I., Quach, C., Russ, M., Kauer, J., Nicolosi, B., Akella, S., Tang, W., Chen, D., Schlueter, J., Munir, M., 2021. Implementing building-level SARS-CoV-2 wastewater surveillance on a university campus. *Sci. Total Environ.* 782, 146749 <https://doi.org/10.1016/j.scitotenv.2021.146749>.
- Gupta, S., Parker, J., Smits, S., Underwood, J., Dolwani, S., 2020. Persistent viral shedding of SARS-CoV-2 in faeces - a rapid review. *Colorectal Dis.* 22 (6), 611–620 <https://doi.org/10.1111/codi.15138>.
- Harris-Lovett, S., Nelson, K.L., Beamer, P., Bischel, H.N., Bivins, A., Bruder, A., Butler, C., Camenisch, T.D., Long, S.K.D., Karthikeyan, S., Larsen, D.A., Meierdiercks, K., Mouser, P.J., Pagsuyoin, S., Prasek, S.M., Radniecki, T.S., Ram, J.L., Roper, D.K., Safford, H., Sherchan, S.P., Shuster, W., Stalder, T., Wheeler, R.T., Korfmacher, K.S., 2021. Wastewater surveillance for SARS-CoV-2 on college campuses: initial efforts, lessons learned and research needs. *Int. J. Environ. Res. Public Health* 18 (9), 4455 <https://doi.org/10.3390/ijerph18094455>.
- Hart, O.E., Halden, R.U., 2020. Computational analysis of SARS-CoV-2/COVID-19 surveillance by wastewater-based epidemiology locally and globally: feasibility, economy, opportunities and challenges. *Sci. Total Environ.* 730, 138875 <https://doi.org/10.1016/j.scitotenv.2020.138875>.
- Hata, A., Hara-Yamamura, H., Meuchi, Y., Imai, S., Honda, R., 2021. Detection of SARS-CoV-2 in wastewater in Japan during a COVID-19 outbreak. *Sci. Total Environ.* 758, 143578 <https://doi.org/10.1016/j.scitotenv.2020.143578>.
- He, X., Lau, E.H.Y., Wu, P., Deng, X.L., Wang, J., Hao, X.X., Lau, Y.C., Wong, J.S.Y., Guan, Y.J., Tan, X.H., Mo, X.N., Chen, Y.Q., Liao, B.L., Chen, W.L., Hu, F.Y., Zhang, Q., Zhong, M.Q., Wu, Y.R., Zhao, L.Z., Zhang, F.C., Cowling, B.J., Li, F., Leung, G.M., 2020. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.* 26 (5) <https://doi.org/10.1038/s41591-020-0869-5>.
- Hewitt, J., Rivera-Aban, M., Greening, G.E., 2009. Evaluation of murine norovirus as a surrogate for human norovirus and hepatitis A virus in heat inactivation studies. *J. Appl. Microbiol.* 107 (1), 65–71 <https://doi.org/10.1111/j.1365-2672.2009.04179.x>.
- Hoffmann, T., Alsing, J., 2021. Faecal shedding models for SARS-CoV-2 RNA amongst hospitalised patients and implications for wastewater-based epidemiology. *medRxiv*, 2021.2003.2016.21253603. <https://doi.org/10.1101/2021.03.16.21253603>.
- Hong, P.-Y., Rachmadi, A.T., Mantilla-Calderson, D., Alkahtani, M., Bashawri, Y.M., Al Qarni, H., O'Reilly, K.M., Zhou, J., 2021. Estimating the minimum number of SARS-CoV-2 infected cases needed to detect viral RNA in wastewater: to what extent of the outbreak can surveillance of wastewater tell us? *Environ. Res.* 195, 110748 <https://doi.org/10.1016/j.envres.2021.110748>.
- Jefferies, S., French, N., Gilkison, C., Graham, G., Hope, V., Marshall, J., McElroy, C., McNeill, A., Muellner, P., Paine, S., Prasad, N., Scott, J., Sherwood, J., Yang, L., Priest, P., 2020. COVID-19 in New Zealand and the impact of the national response: a descriptive epidemiological study. *Lancet Public Health* 5 (11), E612–E623 [https://doi.org/10.1016/S2468-2667\(20\)30225-5](https://doi.org/10.1016/S2468-2667(20)30225-5).
- Kantor, R.S., Nelson, K.L., Greenwald, H.D., Kennedy, L.C., 2021. Challenges in measuring the recovery of SARS-CoV-2 from wastewater. *Environ. Sci. Technol.* 55 (6), 3514–3519 <https://doi.org/10.1021/acs.est.0c08210>.
- Li, W., Su, Y.Y., Zhi, S.B., Huang, J., Zhuang, C.L., Bai, W.Z., Wan, Y., Meng, X.R., Zhang, L., Zhou, Y.B., Luo, Y.Y., Ge, S.X., Chen, Y.K., Ma, Y., 2020. Virus shedding dynamics in asymptomatic and mildly symptomatic patients infected with SARS-CoV-2. *Clin. Microbiol. Infect.* 26 (11) <https://doi.org/10.1016/j.cmi.2020.07.008>.
- Lodder, W., Husman, A.M.D., 2020. SARS-CoV-2 in wastewater: potential health risk, but also data source. *Lancet Gastroenterol. Hepatol.* 5 (6), 533–534 [https://doi.org/10.1016/S2468-1253\(20\)30087-X](https://doi.org/10.1016/S2468-1253(20)30087-X).
- Medema, G., Been, F., Heijnen, L., Petterson, S., 2020a. Implementation of environmental surveillance for SARS-CoV-2 virus to support public health decisions: opportunities and challenges. *Current Opinion in Environmental Science and Health* 17, 49–71 <https://doi.org/10.1016/j.coesh.2020.09.006>.
- Medema, G., Heijnen, L., Elsinga, G., Italiaander, R., Brouwer, A., 2020b. Presence of SARS-Coronavirus-2 RNA in sewage and correlation with reported COVID-19 prevalence in the early stage of the epidemic in The Netherlands. *Environ. Sci. Technol. Lett.* 7 (7), 511–516 <https://doi.org/10.1021/acs.estlett.0c00357>.
- New Zealand Government. (2021) Welcome pack: managed isolation facility for returnees to New Zealand. <https://www.miq.govt.nz/being-in-managed-isolation/entering-isolation/welcome-pack/welcome-pack/>. Accessed on 11 January 2021.
- Peccia, J., Zulli, A., Brackney, D.E., Grubaugh, N.D., Kaplan, E.H., Casanovas-Massana, A., Ko, A.I., Malik, A.A., Wang, D., Wang, M.K., Warren, J.L., Weinberger, D.M., Arnold, W., Omer, S.B., 2020. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat. Biotechnol.* 38 (10), 1164–1167 <https://doi.org/10.1038/s41587-020-0684-z>.
- Pecson, B.M., Darby, E., Haas, C.N., Amha, Y., Bartolo, M., Danielson, R., Dearborn, Y., Di Giovanni, G., Ferguson, C., Fevig, S., Gaddis, E., Gray, D., Lukaski, G., Mull, B., Olivas, L., Olivieri, A., Qu, Y., Consortium, S.-C.-I., 2021. Reproducibility and sensitivity of 36 methods to quantify the SARS-CoV-2 genetic signal in raw wastewater: findings from an interlaboratory methods evaluation in the U.S. *Environ. Sci.: Water Res. Technol.* 7, 504–520 <https://doi.org/10.1039/D0EW00946F>.
- Philo, S.E., Keim, E.K., Swanson, R., Ong, A.Q.W., Burnor, E.A., Kossik, A.L., Harrison, J.C., Demeke, B.A., Zhou, N.A., Beck, N.K., Shirai, J.H., Meschke, J.S., 2021. A comparison of SARS-CoV-2 wastewater concentration methods for environmental surveillance. *Sci. Total Environ.* 760 (15), 144215 <https://doi.org/10.1016/j.scitotenv.2020.144215>.
- R Core Team, 2020. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Randazzo, W., Truchado, P., Cuevas-Ferrando, E., Simón, P., Allende, A., Sánchez, G., 2020. SARS-CoV-2 RNA in wastewater anticipated COVID-19 occurrence in a low prevalence area. *Water Res.* 181, 115942 <https://doi.org/10.1016/j.watres.2020.115942>.
- Salvo, M., Moller, A., Alvareda, E., Gamazo, P., Colina, R., Victoria, M., 2021. Evaluation of low-cost viral concentration methods in wastewaters: implications for SARS-CoV-2 pandemic surveillances. *J. Virol. Methods* 297, 114249 <https://doi.org/10.1016/j.jviromet.2021.114249>.
- Sims, N., Kasprzyk-Hordern, B., 2020. Future perspectives of wastewater-based epidemiology: monitoring infectious disease spread and resistance to the community level. *Environ. Int.* 139, 105689 <https://doi.org/10.1016/j.envint.2020.105689>.
- Spurbeck, R.R., Minard-Smith, A.T., Catlin, L.A., 2021. Applicability of neighborhood and building scale wastewater-based genomic epidemiology to track the SARS-CoV-2 pandemic and other pathogens. *medRxiv* <https://doi.org/10.1101/2021.02.18.21251939>.
- Thompson, J.R., Nanchaiah, Y.V., Gu, X.Q., Lee, W.L., Rajal, V.B., Haines, M.B., Girones, R., Ng, L.C., Alm, E.J., Wuertz, S., 2020. Making waves: wastewater surveillance of SARS-CoV-2 for population-based health management. *Water Res.* 184, 116181 <https://doi.org/10.1016/j.watres.2020.116181>.
- Torii, S., Oishi, W., Zhu, Y., Thakali, O., Malla, B., Yu, Z., Zhao, B., Arakawa, C., Kitajima, M., Hata, A., Ihara, M., Kyuwa, S., Sano, D., Haramoto, E., Katayama, H., 2022. Comparison of five polyethylene glycol precipitation procedures for the RT-qPCR based recovery of murine hepatitis virus, bacteriophage phi6, and pepper mild mottle virus as a surrogate for SARS-CoV-2 from wastewater. *Sci. Total Environ.* 807 (2), 150722 <https://doi.org/10.1016/j.scitotenv.2021.150722>.
- Walsh, K.A., Jordan, K., Clyne, B., Rohde, D., Drummond, L., Byrne, P., Ahern, S., Carty, P.G., O'Brien, K.K., O'Murchu, E., O'Neill, M., Smith, S.M., Ryan, M., Harrington, P., 2020. SARS-CoV-2 detection, viral load and infectivity over the course of an infection. *J. Infect.* 81 (3), 357–371 <https://doi.org/10.1016/j.jinf.2020.06.067>.
- Wölfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Muller, M.A., Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., Hoelscher, M., Bleicker, T., Brünink, S., Schneider, J., Ehmann, R., Zwirgmaier, K., Drosten, C., Wendtner, C.,

2020. Virological assessment of hospitalized patients with COVID-2019. *Nature* 581 (7809), 465–469 <https://doi.org/10.1038/s41586-020-2196-x>.
- World Health Organisation. (2020) WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19—11-march-2020>. Accessed on 11 January 2021.
- Wu, F.Q., Zhang, J.B., Xiao, A., Gu, X.Q., Lee, W.L., Armas, F., Kauffman, K., Hanage, W., Matus, M., Ghaeli, N., Endo, N., Duvallet, C., Poyet, M., Moniz, K., Washburne, A.D., Erickson, T., Chai, P., Thompson, J., Alm, E., 2020. SARS-CoV-2 titers in wastewater are higher than expected from clinically confirmed cases. *MSystems* 5 (4) <https://doi.org/10.1128/mSystems.00614-20>.
- Wurtzer, S., Waldman, P., Ferrier-Rembert, A., Frenois-Veyrat, G., Mouchel, J.M., Boni, M., Maday, Y., OBEPINE consortium, Marechal, V., Moulin, L., 2021. Several forms of SARS-CoV-2 RNA can be detected in wastewaters: implications for wastewater-based epidemiology and risk assessment. *Water Res.* 198, 117183 <https://doi.org/10.1016/j.watres.2021.117183>.
- Yuan, C., Wang, H., Li, K., Tang, A., Dai, Y., Wu, B., Zhang, H., Chen, J., Liu, J., Wu, W., Gu, S., Wang, H., Xu, H., Wu, M., Yu, M., Wang, Y., Yu, X., He, J., Liu, S., Zhang, Y., Tong, Z., Yan, J., 2021. SARS-CoV-2 viral shedding characteristics and potential evidence for the priority for faecal specimen testing in diagnosis. *PLoS ONE* 16 (2), e0247367 <https://doi.org/10.1371/journal.pone.0247367>.
- Zhang, W., Du, R.H., Li, B., Zheng, X.S., Yang, X.L., Hu, B., Wang, Y.Y., Xiao, G.F., Yan, B., Shi, Z.L., Zhou, P., 2020. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. *Emerg. Microbes Infect.* 9 (1), 386–389 <https://doi.org/10.1080/22221751.2020.1729071>.