



RESEARCH ARTICLE

Open Access

# Membrane TNF-alpha-activated programmed necrosis is mediated by Ceramide-induced reactive oxygen species

Shidrokh Ardestani<sup>1</sup>, Desirae L Deskins<sup>1</sup> and Pampee P Young<sup>1,2,3\*</sup>

## Abstract

**Background:** Programmed necrosis is a form of caspase-independent cell death whose molecular regulation is poorly understood. While tumor necrosis factor-alpha (TNF- $\alpha$ ) has been identified as an activator of programmed necrosis, the specific context under which this can happen is unclear. Recently we reported that TNF- $\alpha$  can be expressed by human tumor cells as both a membrane tethered (mTNF- $\alpha$ ) and a soluble (sTNF- $\alpha$ ) form. Whereas low level, tumor-derived sTNF- $\alpha$  acts as a tumor promoter, tumor cell expression of mTNF- $\alpha$  significantly delays tumor growth in mice, in large part by induction of programmed necrosis of tumor associated myeloid cells. In this study we sought to determine the molecular mechanism involved in mTNF- $\alpha$  oxidative stress-induced cell death by evaluating the known pathways involved in TNF receptor-induced programmed necrosis.

**Methods:** The source of Reactive Oxygen Species (ROS) in mTNF- $\alpha$  treated cells was determined by coculturing RAW 264.7 monocytic and L929 fibroblasts cells with fixed B16F10 control or mTNF- $\alpha$  expressing-melanoma cells in the presence of inhibitors of NADPH and mitochondria ROS. To identify the down-stream effector of TNF- $\alpha$  receptors (TNFR), level of phospho-RIP-1 and ceramide activity were evaluated.

To determine whether mTNF-mediated cell death was dependent on a specific TNFR, cell death was measured in primary CD11b myeloid cells isolated from wild-type or TNFR-1, TNFR-2, TNFR-1 and TNFR-2 double knockout mice, cocultured with various TNF- $\alpha$  isoform.

**Results:** Tumor derived-mTNF- $\alpha$  increased ROS-mediated cytotoxicity, independent of caspase-3 activity. Although TNFR on target cells were required for this effect, we observed that mTNF-induced cell death could be mediated through both TNFR-1 and the death domain-lacking TNFR-2. ROS generation and cytotoxicity were inhibited by a mitochondrial respiratory chain inhibitor but not by an inhibitor of NADPH oxidase. mTNF- $\alpha$  mediated cytotoxicity was independent of RIP-1, a serine/threonine kinase that serves as a main adaptor protein of sTNF- $\alpha$  induced programmed necrosis. Instead, mTNF- $\alpha$ -induced ROS and cell death was prohibited by the ceramide-activated protein kinase (CAPK) inhibitor.

**Conclusion:** These findings demonstrate that the mTNF- $\alpha$  isoform is an effective inducer of programmed necrosis through a caspase independent, ceramide-related pathway. Interestingly, unlike sTNF $\alpha$ , mTNF-induced programmed necrosis is not dependent on the presence of TNFR1.

**Keywords:** Membrane-TNF-alpha (mTNF- $\alpha$ ), Reactive oxygen species (ROS), Ceramide, Programmed necrosis, Mitochondria

\* Correspondence: pampee.young@vanderbilt.edu

<sup>1</sup>Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center, 1161 21st Avenue South, C2217A MCN, Nashville, TN 37232, USA

<sup>2</sup>Department of Internal Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA

Full list of author information is available at the end of the article

## Background

Tumor necrosis factor-alpha (TNF- $\alpha$ ) is an inflammatory cytokine, that activates cell inflammation, proliferation, survival and cell death depending on autocrine/paracrine signals, and on the cellular context [1,2]. The soluble homotrimeric form of TNF- $\alpha$  (sTNF- $\alpha$ ) that is released from the cell surface activates multiple signal transduction pathways including NFkB survival pathway. In addition to activation of cell survival pathways, sTNF- $\alpha$  can induce cell death [3,4]. Activation of caspases and initiation of apoptosis has been described as the classic form of TNF-mediated cell death. Recent evidence suggests that sTNF- $\alpha$  can also trigger an alternative form of cell death that is distinct from apoptosis. This form of cell death is referred to as “programmed necrosis” and is dependent on the generation of reactive oxygen species [5,6].

So far, in the majority of studies TNF-mediated programmed necrosis have been attributed to the biological and mechanistic function of sTNF- $\alpha$  and its interaction with TNFR-1 in the presence of pharmacological or genetic inhibition of apoptosis [7,8]. TNF- $\alpha$  can also exist as a membrane-anchored protein [9,10]. Like sTNF $\alpha$ , (mTNF-a) is biologically active and binds either of the two TNF-receptors [11]. A recent study from our laboratory indicated that human lung NSCLC express both soluble and membrane isoforms. Using a murine lung cancer model we showed that unlike sTNF- $\alpha$ , mTNF- $\alpha$  exhibits inhibitory effects on tumor growth and myeloid content. We demonstrated that mTNF- $\alpha$  efficiently induced myeloid cell death through induction of ROS-mediated necrosis in the absence of any apoptosis inhibitors [12]. Currently nothing has been reported on how mTNF- $\alpha$  mediates programmed necrosis.

Soluble TNF-induced programmed necrosis typically occurs where apoptosis is inhibited, and is mediated through a few defined pathways. In all cases, the serine/threonine kinase receptor interacting protein-1 (RIP-1) has been shown to play a central role in initiation of programmed necrosis [13,14], mainly through nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (NOX) or mitochondria [15-18]. Recent studies have also described a role for ceramide-mediated programmed necrosis. An increase in the level of intracellular ceramide has been linked to increased redox reaction within the cell [19,20], suggesting the potential for crosstalk between ceramide, ROS, and TNF- $\alpha$  pathways in this process. In spite of these observations, the specific mechanism by which ceramide signaling leads to increased redox reactions is not fully understood.

In this study we sought to determine the molecular pathway involved in mTNF- $\alpha$ -mediated oxidative stress-induced cell death. Using inhibitors targeting mitochondrial electron transport chain and NADPH oxidase we concluded that mitochondrial-dependent oxidative stress was the major source of mTNF-induced intracellular ROS

generation, regulated by ceramide-activated protein kinase (CAPK) activity. To our knowledge, this is the first report identifying mTNF- $\alpha$  isoform as a potent activator of programmed necrosis, even in the absence of inhibitors of apoptosis, via ceramide-dependent mitochondrial ROS generation.

## Results and discussion


### mTNF- $\alpha$ is an inducer of cell death

To investigate the ability of the membrane versus soluble TNF- $\alpha$  isoforms to induce cell death, RAW 264.7 cells, a line derived from murine leukemic monocytes/macrophage cells (target), were mixed with 1% paraformaldehyde-fixed B16F10 melanoma cells (effector) expressing empty vector,  $\pm 100$  U/ml rTNF- $\alpha$  (FxB16<sub>cont</sub> or FxB16<sub>cont</sub> + rTNF), or fixed mTNF-expressing B16F10 cells (FxB16<sub>mTNF</sub>) at a target: effector ratio of 1:10. Paraformaldehyde-fixed tumor cells were used to eliminate the endogenous sTNF- $\alpha$  (Figure 1A). As measured by the MTT dye reduction assay, incubation with FxB16<sub>mTNF</sub> resulted in more than  $70 \pm 12\%$  cytotoxicity of RAW 264.7 myeloid cells after 48 hours of incubation, as compared to RAW 264.7 cells incubated with FxB16<sub>cont</sub> (Figure 1B,  $P < 0.05$ ). In contrast FxB16<sub>cont</sub> + rTNF increased RAW 264.7 cell survival compared to control.

To determine the molecular pathway leading to mTNF-induced cell death, we utilized the highly TNF-sensitive L929 fibrosarcoma cell line. As shown in Figure 1C, mTNF- $\alpha$  isoform resulted in more than 50% cell death compared to L929 cocultured with FxB16<sub>cont</sub> as determined by MTT reduction assay. Cellular toxicity causes membrane damage and results in the release of lactate dehydrogenase (LDH) from the cytoplasm and thus LDH in the media can be used to measure cell death. To confirm the results obtained with MTT assay, we measured LDH release in L929 cell in the presence of control or mTNF-expressing tumor cells. The mTNF- $\alpha$  isoform increased the level of LDH leakage by 3.7 fold over control (Figure 1D).

### Membrane TNF- $\alpha$ -induced cell death can be mediated through both TNFR-1 and TNFR-2

mTNF- $\alpha$  signal transduction has been linked to a cooperative signaling between TNFR-1 and TNFR-2 [7,21]. Next we sought to determine whether mTNF-mediated cell death was dependent on a specific TNF receptor. Primary CD11b myeloid cells were isolated from wild-type (WT), TNFR-1 knockout (TNFR-1KO), TNFR-2KO or TNFR-1 and TNFR-2 double knockout (TNFR-DKO) and cocultured with fixed control tumor cells with or without rTNF or fixed mTNF-expressing tumor cells for 48 hours. Cell cytotoxicity was determined by MTT assay. As shown in Figure 2, presence of either TNFR-1 or TNFR-2 resulted in increased levels of mTNF-induced cytotoxicity similar



**Figure 1 Membrane TNF- $\alpha$  isoform effectively induces cell death.** **A**, Schematic diagram of fixed B16F10 cells cocultured with L929/RAW264.7. **B**, RAW cells were cocultured with paraformaldehyde-fixed control (FxB16<sub>cont</sub>), control + rTNF $\alpha$  (FxB16<sub>cont</sub> + TNF $\alpha$ ), or mTNF $\alpha$  (FxB16<sub>mTNF</sub>) for 24 hours. Cell death was measured by MTT assay. **C**, MTT assay showing the cytotoxic effects of mTNF- $\alpha$  isoform on L929. **D**, LDH assay measuring L929 cells percent LDH leakage, in the presence of control or mTNF-expressing fixed B16F10 tumor cells. Data show the percentage of LDH leakage into media to total LDH (media + cells). Each sample was assayed in triplicate, with each experiment repeated at least 3 times independently. Data are expressed as average  $\pm$  S.E. \* $P$  < 0.05 and \*\* $P$  < 0.005, 1-way ANOVA of log (10) transformed data with Student's t-test as post-test (**B**), students's t-test (**C** and **D**).


to WT-CD11b (~17% cell death in WT-CD11b, ~20% in TNFR-1KO-CD11b and TNFR-2KO-CD11b). In contrast, control cells treated with rTNF improved cell survival in WT- and TNFR-1KO-CD11b and resulted only in ~4% cell death in TNFR-2KO-CD11b and ~5% cell death in TNFR-DKO-CD11b. Interestingly, mTNF-mediated cell cytotoxicity was reversed in TNFR-DKO-CD11b cells. These findings suggested that mTNF-induced cell death can be mediated through both TNFR-1 and TNFR-2.

#### Membrane TNF- $\alpha$ exerted cell cytotoxicity by increasing intracellular ROS production

Induction of cell death by sTNF- $\alpha$  occurs mainly through activation of caspases leading to apoptosis. To test whether the mTNF- $\alpha$  isoform exerts its cell toxicity in

part by activating the caspase pathway, we determined the level of cleavage/activation of caspase-3 proteins in L929 cells treated with fixed control or mTNF-expressing tumor cells. As presented in Figure 3A, treatment of L929 cells or RAW cells with FxB16<sub>mTNF</sub> did not result in an increased level of active-caspase-3.

Previously we have shown that mTNF- $\alpha$ -treated myeloid cells exhibit increased intracellular ROS and decreased cell survival [12]. To demonstrate an association between intracellular ROS level and cell death in L929 fibrosarcomas, cells incubated with different TNF- $\alpha$  isoforms were measured using CM-H2DCFDA (which quantitatively reacts with oxygen species to produce a highly fluorescent dye). L929 cells incubated with FxB16<sub>mTNF</sub> resulted in a 60% increase in CM-H2DCFDA fluorescence, indicating an



**Figure 2 Membrane TNF-induced cell death can be mediated through both TNFR-1 and TNFR-2.** Freshly isolated CD11b myeloid cells from wild type mice (WT-CD11b), mice deficient in TNF receptor 1 (TNFR-1KO-CD11b), TNFR-2KO-CD11b or both receptors (TNFR-DKO-CD11b). Cells were cocultured with paraformaldehyde-fixed control (FxB16<sub>cont</sub>), control in the presence of 100 U/ml recombinant TNF $\alpha$  (FxB16<sub>cont</sub> + rTNF), or mTNF $\alpha$  (FxB16<sub>mTNF</sub>) for 24 hours. Percentage of cell death was measured by MTT assay. Data present mean percentage (bars, mean  $\pm$  S.E.) of three replicates from 3 independent experiments.

increase in the level of ROS ( $P > 0.05$ ; Figure 3B). Furthermore, incubation of L929 cells with ROS scavenger N-acetyl-cysteine (NAC) reduced mTNF-mediated ROS level (130% of control in FxB16<sub>mTNF</sub>; 94% of control in FxB16<sub>mTNF</sub> + NAC; Figure 3C,  $P < 0.5$ ). This was followed by 4-fold decrease in LDH release in mTNF-treated L929 cells supplied with NAC ( $P < 0.05$ ; Figure 3D).

#### Inhibition of mitochondrial respiratory chain decreased mTNF-mediated ROS generation

Next we sought to determine the source of ROS in response to mTNF- $\alpha$  in L929 cells. There is report of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase-1 (NOX1) and mitochondria as the two major source of TNF- $\alpha$  induced ROS production [22,23]. To do so, the NADPH-dependent oxidase and the mitochondrial respiratory chain complex II were blocked using DPI (2  $\mu$ M) and TTFA (0.5  $\mu$ M) respectively. The NADPH-dependent oxidase inhibitor DPI did not inhibit the ROS production induced by mTNF- $\alpha$  ( $156.7 \pm 8.3\%$  of control in FxB16<sub>mTNF</sub> and  $155.4 \pm 14.4\%$  in FxB16<sub>mTNF</sub> + DPI;  $P < 0.05$ ; Figure 4A) and further had no effect on the LDH leakage (270% of control in FxB16<sub>mTNF</sub> and 280% in FxB16<sub>mTNF</sub> + DPI;  $P < 0.05$ ; Figure 4B). However, addition of TTFA into L929 cells cocultured with mTNF-expressing tumor cells, reduced CM-H2DCFDA oxidation (108% of control) and LDH release (10% of control, Figure 4B). These data suggested that mitochondria are the source of mTNF-induced ROS generation and cell death.

#### Membrane TNF-mediated ROS production involves ceramide pathway


TNFR-mediated mitochondrial ROS generation can be induced through RIP-1 kinase activity or through a

ceramide-dependent signaling pathway initiated by SMase activity (Figure 5A). To determine the specific pathway responsible for activation of mitochondrial ROS production we analyzed level of active RIP1 by evaluating its phosphorylation in L929 or RAW 264.7 in the presence of FxB16<sub>cont</sub> or FxB16<sub>mTNF</sub> by immunoblot analysis. As shown in Figure 5B, treatment of both cell lines with fixed membrane expressing tumor cells, FxB16<sub>mTNF</sub> did not increase the level of RIP-1 phosphorylation. To further confirm that the RIP-1 kinase signaling pathway is not involved in the generation of mitochondrial ROS we assessed cell death by MTT assay with L929 cells incubated with fixed tumor cells, with or without the RIP-1 inhibitor necrostatin (Figure 5C). The inhibitor did not cause any change in L929 cell death when incubated with FxB16<sub>mTNF</sub> compared to control.

There is evidence to support the role of ceramide as a second messenger of TNF- $\alpha$  activated cells involved in activation of programmed necrosis [24]. Next we evaluated the role of ceramide signaling in TNF-induced ROS production and survival. Addition of myriocin, a ceramide inhibitor, reduced the cell death seen in L929 cells incubated with FxB16<sub>mTNF</sub> to a level similar to that seen with cells incubated with control tumor cells (Figure 5D). Furthermore addition of DMAP (1 mM), a CAPK inhibitor, reduced mTNF-induced ROS by 60% ( $138 \pm 15.6\%$  of control in FxB16<sub>mTNF</sub>;  $80 \pm 2.9\%$  in FxB16<sub>mTNF</sub> + DMAP; Figure 5E). Percentage of LDH leakage was also reduced from 276% in mTNF-treated cells to 163% in mTNF-treated cells supplied with DMAP ( $P < 0.005$ , Figure 5F). These findings suggested that mTNF-induced mitochondrial ROS generation requires protein kinase activity associated with ceramide. This was further confirmed in RAW 264.7 cell lines. Similar to L929 cells, inhibition of CAPK in mTNF- $\alpha$  treated Raw 264.7 cells reduced LDH leakage (Figure 5G).

#### Conclusions

Previously necrotic cell death has been defined as a sudden, unregulated form of cell death which leads to inflammation and tissue damage. However, in recent years accumulating evidence suggests that not all form of necrotic cell death is accidental but can instead be a programmed event [5,6]. There have been several reports of TNF-induced programmed necrosis, mainly in the context of the soluble form of TNF- $\alpha$ . Importantly, induction of programmed necrosis by sTNF- $\alpha$  typically requires the presence of inhibitors of caspases [25,26]. Here, we present for the first time that the lesser-known membrane form of TNF- $\alpha$  has the ability to induce programmed necrosis through ROS generation, independent of caspase inhibitors. In this study we explored the mechanism of mTNF-mediated ROS generation and programmed necrosis.




In our study treatment of mTNF-induced L929 cells with mitochondrial inhibitor complex II increased ROS reduction and improved survival, suggesting a role for mitochondrial complex II in mTNF-mediated programmed necrosis. The plasma membrane-associated NADPH oxidases (NOX) have been proposed as an alternate source of ROS production [15,17,27]. In contrast to what we observed with mitochondrial inhibitor, inhibition of NOX failed to inhibit ROS generation and to increase cell viability.

Although, RIP1/RIP3 kinases have been shown to orchestrate the programmed necrosis pathway activity of sTNF- $\alpha$ , we did not detect any phosphorylated RIP1; instead we found that ceramide pathway was involved in mTNF- $\alpha$ -induced cell death. Ceramide pathway has been identified as an alternative mechanism for induction of

programmed necrosis [13,28]. An enhanced level of ceramide has been shown to contribute to depletion of mitochondrial reduced glutathione [29] and increasing mitochondria susceptibility to GD3, a ceramide-derived ganglioside. GD3 traffics to the mitochondria and directly induces ROS production [30,31]. In our study mTNF-induced ROS and cell death seems to be regulated through activity of ceramide since the inhibitor of CAPK, blocked mTNF-mediated ROS and cell death.

The molecular mechanism of the two different TNF- $\alpha$  isoforms remains elusive. It is interesting that although sTNF- $\alpha$  and mTNF- $\alpha$  have similar structures and are able to interact with both TNF- $\alpha$  receptors, they exert opposing effects on tumor growth and cell survival. This raises questions about the plausible molecular mechanisms to account for this difference.



**Figure 4** Inhibition of mitochondrial respiratory chain decreases mTNF-mediated ROS generation. **A** and **B**, L929 cell cocultured with mTNF-expressing tumor cells in the absence or presence of NOX inhibitor-DPI (2  $\mu$ M) and mitochondrial complex II inhibitor-TTFA (0.5  $\mu$ M) for 24 hours. TTFA reduced ROS level, shown by reduction of CM-H2DCFDA intensity (**A**) and LDH leakage into media (**B**). Addition of NOX inhibitor-DPI had no effects on both ROS generation of LDH level. Data are represented as % of CM-H2DCFDA intensity (**A**) or LDH in media/total LDH (**B**) to L929 cells cocultured with control expressing tumor cells with or without DPI or TTFA. Data are expressed as average  $\pm$  S.E. \* $P < 0.05$ .

The variation in biological response of soluble versus the membrane form of cytokines has been observed in other members of the TNF receptor superfamily. Membrane form of Fas ligand, for example, has been shown to have opposing role in modulating cell death when compared to its soluble form [32]. Previous work by Gregory and colleague has shown that in glaucoma, full-length FasL accelerates retinal ganglion cells death. By contrast, FasL-deficiency, or administration of soluble FasL, has a protective effect [33]. TNF-related apoptosis-inducing ligand (TRAIL), which has important functions in inducing apoptosis, has shown to have differential activation capacity toward TRAIL-R1 and R2 as a soluble form versus membrane form [34].

Unlike soluble TNF- $\alpha$  which its signal transduction is mediated mainly through TNFR-1 [5,35], it is evident from our study that mTNF- $\alpha$  is efficient in activation of both TNFR-1 and TNFR-2. Members of the tumor necrosis factor receptors superfamily (TNFR-1, Fas, TRAIL) that are capable of inducing cell death usually contain a conserved DD in the intracellular region that is required for activation cell death [1,36]. It is interesting that in our study TNFR-2 which lacks a DD-domain, also seems to be sufficient for induction of mTNF- $\alpha$ -mediated cell death. This phenomenon has been reported in other studies. For instance, TNF-R2 shown to trigger cell death in the rhabdomyosarcoma cell line KYM-1 [37] and the stimulation of CD30 induces cell death in T cell hybridomas [38]. It is not yet clear how TNF receptor superfamily members lacking a death domain (i.e. TNF-R2) execute their death inducing capability. This effect could be FADD-dependent [39] or could be mediated through cooperative activity with other receptors such as Fas/FasL [40].


It has been proposed that membrane anchored TNF- $\alpha$  creates a more stable contact with TNFR1 or TNFR2 with longer half-life. This will result in the formation of higher-order receptor complexes (receptor clustering) [41]. These clusters of receptors are formed in lipid rafts abundant in sphingomyelin which are the precursor of ceramide [42]. Formation of the raft and receptor clustering recruit sphingomylinases to the membrane and result in increased levels of ceramide. Ceramide will decrease mitochondrial membrane integrity and ROS leakage [43]. It is possible that mTNF- $\alpha$ -mediated ceramide production is regulated through multiple TNF receptor superfamily members clustering. This requires further investigation.

In conclusion we have demonstrated that mTNF- $\alpha$  can induce cell death independent of caspase inhibitors by increasing ROS. This occurs through RIP-1-independent, ceramide-dependent activation of mitochondrial ROS. Molecular mechanism, leading to mTNF-induced ceramide formation and mitochondrial-ROS generation, remains to be investigated.

## Methods

### Mice, cell lines and materials

Wild-type C57Bl/6 (WT) mice were purchased from Jackson Laboratory. Homozygous mutants for TNFR-1, TNFR-2, and TNFR-1/2 double knockout (TNFR-DKO) on a C57Bl/6 background were a generous gift from Dr. D. Polk. B16F10 melanoma, RAW264.7 and L929 cells were purchased from American Type Culture Collection (ATCC) and were maintained in DMEM with 10% (v/v) fetal bovine serum, heat inactive fetal bovine serum and MEM with 10% (v/v) horse serum, respectively. N-Acetyl-



L-cysteine (NAC), 2-Thenoyltrifluoroacetone (TTFA), myxothiazol and 4-(Dimethylamino)pyridine (DMAP) were purchased from Sigma-Aldrich and prepared fresh on the day of the experiment.

### Constructs

Membrane TNF-expressing cells were generated by cloning the mTNF $\Delta$ 1-9, K11E sequence encoding a mutant transmembrane TNF- $\alpha$  protein with a deletion at the cleavage site between pre-sequence and mature membrane TNF- $\alpha$  (BCCM/LMBP plasmid collection, Ghent University) into the BamH1-EcoR1 site of LZRS-IRES-Neo retroviral vector, conferring neomycin resistance. This mutation prevents cleavage of the 26-kDa membrane TNF- $\alpha$  into secretory TNF- $\alpha$  isoform. An empty LZRS vector was used as a control vector [12].

### Surface expression of TNF- $\alpha$

Trypsinized cells were incubated with anti-TNF- $\alpha$  antibody (1  $\mu$ l/2.5  $\times$  10 $^4$  cells, Southern Biotech) for 30 minutes on ice. PE conjugated secondary antibody (0.125  $\mu$ g/10 $^6$  cells/100  $\mu$ l) was added for 30 minutes on ice. Surface expression of TNF- $\alpha$  was measured using flow cytometry.

### TNF- $\alpha$ cytotoxicity assay

Overnight cultured RAW 264.7, L929 or freshly isolated bone marrow CD11b cells (EasySep, StemCell Technologies) (2.5  $\times$  10 $^4$ /100  $\mu$ L/well, target) were cocultured with Paraformaldehyde-fixed [44] control, control + 100 U/ml of recombinant TNF- $\alpha$  or mTNF- $\alpha$  (effector) at target:effector ratio of 1:10 and incubated for 48 hours. Cells were labeled with 100  $\mu$ l of PBS containing 0.5 mg/mL of 3-[4,5-dimethylthiazol-2-yl] 2,5-diphenyltetrazolium bromide (MTT) (Sigma) for 2 hours at 37°C then lysed with 0.1 ml DMSO. Photometric measurement was carried out at 540 nm. Percentage of cell death was calculated by using the following formula: Cell death (%) = (1 – OD<sub>sample</sub>/OD<sub>control</sub>)  $\times$  100.

### Lactate dehydrogenase (LDH) assay

Early cell damage was determined using the LDH cytotoxicity detection kit (Promega, Madison, WI), which quantifies the LDH release from the cells into the culture medium. Cells were seeded in a 96-well plates at a density of 2.5  $\times$  10 $^4$  cells/well overnight to promote adherence. Cells were cocultured with fixed B16F10 control or mTNF- $\alpha$  in the absence or presence of the indicated treatments for 24 hours. Supernatants from the cultures were collected and used in the LDH assay as instructed by the manufacturer. LDH activity was detected separately in the supernatant and cell lysate. The percentage of LDH leakage was calculated as 100  $\times$  (LDH<sub>supernatant</sub> / (LDH<sub>supernatant</sub> + lysate)).

### Measurement of intracellular ROS

The oxidant-sensing probe CM-H2DCFDA (Invitrogen) was used to detect intracellular reactive oxygen species (ROS). An overnight culture of cells (2.5  $\times$  10 $^4$ ) were loaded with 10  $\mu$ M CM-H2DCFDA for 30 minutes and cocultured with 2.5  $\times$  10 $^5$  fixed B16F10 control or mTNF-expressing cell in the absence or presence of the indicated treatments for 6 hours. Fluorescence was determined using a luminescence spectrophotometer (Spectra max, Molecular Devices) with an excitation wavelength of 495 nm and emission wavelength of 525 nm.

### Immunoblot analysis

Target cells were plated at a density of 10 $^6$  cells overnight (37°C, 5% CO<sub>2</sub>). Cells then were treated with fixed control or mTNF-expressing B16F10 at target cell/fixed tumor cell ratio of 1:10. After 30 minutes incubation at 37°C, target cells were lysed with RIPA buffer [3 M NaCl, 1 M Tris, 0.5 M EDTA, 10% SDS, 1% NP40 substitute and 1 $\times$  Complete Protease Inhibitor Cocktail (Roche)]. Whole cell lysate was evaluated for caspase-3 cleavage (Cell Signaling Technology) and RIP-1 (BD biosciences) by Western blot analysis.

### Statistical analysis

The statistical significance between experimental and control groups was determined by performing log (10) transformation of the data such that we can apply the parametric analysis, ANOVA followed by Student's t-test as a post-test using Prism software (Graphpad, San Diego, CA). A P-value of <0.05 was considered statistically significant.

### Abbreviations

LDH: Lactate dehydrogenase; TTFA: Thenoyltrifluoroacetone; CAPK: Ceramide-activated protein kinase; DPI: Diphenyleneiodonium; DMAP: 4-Dimethylaminopyridine.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

SA contributed in design, development of methodology, acquisition of data, and writing of the manuscript. DD contributed in design and acquisition, analysis and interpretation of data. PPY contributed in design, development of methodology, analysis and interpretation of data and writing of the manuscript and supervised the study. All authors read and approved the final manuscript.

### Acknowledgements

This work was supported by funding from the NIH (R01-HL088424) to PPY, Veterans Affairs Merit Award to PPY, Vanderbilt University Clinical and Translational Science Award (UL1 RR024975-01 from NCRR/NIH) to PPY, Immunobiology of Blood and Vascular Systems training grant (5 T32 HL069765) to SA.

### Author details

<sup>1</sup>Department of Pathology, Microbiology and Immunology, Vanderbilt University Medical Center, 1161 21st Avenue South, C2217A MCN, Nashville, TN 37232, USA. <sup>2</sup>Department of Internal Medicine, Vanderbilt University

Medical Center, Nashville, TN 37232, USA. <sup>3</sup>Department of Veterans Affairs Medical Center, Nashville, TN 37212, USA.

Received: 13 June 2013 Accepted: 23 October 2013

Published: 1 November 2013

## References

1. Rangamani P, Sirovich L: Survival and apoptotic pathways initiated by TNF-alpha: modeling and predictions. *Biotechnol Bioeng* 2007, **97**:1216–1229.
2. Wallach D, Varfolomeev EE, Malinin NL, Goltsev YV, Kovalenko AV, Boldin MP: Tumor necrosis factor receptor and Fas signaling mechanisms. *Annu Rev Immunol* 1999, **17**:331–367.
3. Baud V, Karin M: Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol* 2001, **11**:372–376.
4. Balkwill F: Tumour necrosis factor and cancer. *Nat Rev Cancer* 2009, **9**:361–371.
5. Morgan MJ, Kim YS, Liu ZG: TNFalpha and reactive oxygen species in necrotic cell death. *Cell Res* 2008, **18**:343–349.
6. Wu W, Liu P, Li J: Necroptosis: an emerging form of programmed cell death. *Crit Rev Oncol Hematol* 2012, **82**:249–258.
7. Chan FK, Shisler J, Bixby JG, Felices M, Zheng L, Appel M, Orenstein J, Moss B, Lenardo MJ: A role for tumor necrosis factor receptor-2 and receptor-interacting protein in programmed necrosis and antiviral responses. *J Biol Chem* 2003, **278**:51613–51621.
8. Arnott CH, Scott KA, Moore RJ, Robinson SC, Thompson RG, Balkwill FR: Expression of both TNF-alpha receptor subtypes is essential for optimal skin tumour development. *Oncogene* 2004, **23**:1902–1910.
9. Perez C, Albert I, DeFay K, Zachariaes N, Gooding L, Kriegler M: A nonsecretable cell surface mutant of tumor necrosis factor (TNF) kills by cell-to-cell contact. *Cell* 1990, **63**:251–258.
10. Horiuchi T, Mitoma H, Harashima S, Tsukamoto H, Shimoda T: Transmembrane TNF-alpha: structure, function and interaction with anti-TNF agents. *Rheumatology (Oxford)* 2010, **49**:1215–1228.
11. Horiuchi T, Morita C, Tsukamoto H, Mitoma H, Sawabe T, Harashima S, Kashiwagi Y, Okamura S: Increased expression of membrane TNF-alpha on activated peripheral CD8+ T cells in systemic lupus erythematosus. *Int J Mol Med* 2006, **17**:875–879.
12. Ardestani S, Li B, Deskins DL, Wu H, Massion PP, Young PP: Membrane versus soluble isoforms of TNF-a exert opposing effects on tumor growth and survival of tumor-associated myeloid cells. *Cancer Res* 2013, **73**:3938–3950.
13. Li J, McQuade T, Siemer AB, Napetschnig J, Moriwaki K, Hsiao YS, Damko E, Moquin D, Walz T, McDermott A, et al: The RIP1/RIP3 necrosome forms a functional amyloid signaling complex required for programmed necrosis. *Cell* 2012, **150**:339–350.
14. Moquin D, Chan FK: The molecular regulation of programmed necrotic cell injury. *Trends Biochem Sci* 2010, **35**:434–441.
15. Fan J, Frey RS, Rahman A, Malik AB: Role of neutrophil NADPH oxidase in the mechanism of tumor necrosis factor-alpha-induced NF-kappa B activation and intercellular adhesion molecule-1 expression in endothelial cells. *J Biol Chem* 2002, **277**:3404–3411.
16. Frey RS, Rahman A, Kefer JC, Minshall RD, Malik AB: PKCzeta regulates TNF-alpha-induced activation of NADPH oxidase in endothelial cells. *Circ Res* 2002, **90**:1012–1019.
17. Hordijk PL: Regulation of NADPH oxidases: the role of Rac proteins. *Circ Res* 2006, **98**:453–462.
18. Kim YS, Morgan MJ, Choksi S, Liu ZG: TNF-induced activation of the Nox1 NADPH oxidase and its role in the induction of necrotic cell death. *Mol Cell* 2007, **26**:675–687.
19. Garcia-Ruiz C, Colell A, Mari M, Morales A, Fernandez-Checa JC: Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. Role of mitochondrial glutathione. *J Biol Chem* 1997, **272**:11369–11377.
20. Huwiler A, Pfeilschifter J, van den Bosch H: Nitric oxide donors induce stress signaling via ceramide formation in rat renal mesangial cells. *J Biol Chem* 1999, **274**:7190–7195.
21. Lazdins JK, Grell M, Walker MR, Woods-Cook K, Scheurich P, Pfizenmaier K: Membrane tumor necrosis factor (TNF) induced cooperative signaling of TNFR60 and TNFR80 favors induction of cell death rather than virus production in HIV-infected T cells. *J Exp Med* 1997, **185**:81–90.
22. Bedard K, Krause KH: The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 2007, **87**:245–313.
23. Liu Y, Fiskum G, Schubert D: Generation of reactive oxygen species by the mitochondrial electron transport chain. *J Neurochem* 2002, **80**:780–787.
24. Corda S, Laplace C, Vicaut E, Duranteau J: Rapid reactive oxygen species production by mitochondria in endothelial cells exposed to tumor necrosis factor-alpha is mediated by ceramide. *Am J Respir Cell Mol Biol* 2001, **24**:762–768.
25. Zhang DW, Shao J, Lin J, Zhang N, Lu BJ, Lin SC, Dong MQ, Han J: RIP3, an energy metabolism regulator that switches TNF-induced cell death from apoptosis to necrosis. *Science* 2009, **325**:332–336.
26. Robaye B, Mosselmans R, Fiers W, Dumont JE, Galand P: Tumor necrosis factor induces apoptosis (programmed cell death) in normal endothelial cells in vitro. *Am J Pathol* 1991, **138**:447–453.
27. Quinlan CL, Orr AL, Perevoshchikova IV, Treberg JR, Ackrell BA, Brand MD: Mitochondrial complex II can generate reactive oxygen species at high rates in both the forward and reverse reactions. *J Biol Chem* 2012, **287**:27255–27264.
28. Vandenabeele P, Declercq W, van Herreweghe F, Vanden Berghe T: The role of the kinases RIP1 and RIP3 in TNF-induced necrosis. *Sci Signal* 2010, **3**:re4.
29. Mari M, Colell A, Morales A, Paneda C, Varela-Nieto I, Garcia-Ruiz C, Fernandez-Checa JC: Acidic sphingomyelinase downregulates the liver-specific methionine adenosyltransferase 1A, contributing to tumor necrosis factor-induced lethal hepatitis. *J Clin Invest* 2004, **113**:895–904.
30. Garcia-Ruiz C, Colell A, Morales A, Calvo M, Enrich C, Fernandez-Checa JC: Trafficking of ganglioside GD3 to mitochondria by tumor necrosis factor-alpha. *J Biol Chem* 2002, **277**:36443–36448.
31. Garcia-Ruiz C, Colell A, Paris R, Fernandez-Checa JC: Direct interaction of GD3 ganglioside with mitochondria generates reactive oxygen species followed by mitochondrial permeability transition, cytochrome c release, and caspase activation. *FASEB J* 2000, **14**:847–858.
32. Date T, Mochizuki S, Belanger AJ, Yamakawa M, Luo Z, Vincent KA, Cheng SH, Gregory RJ, Jiang C: Differential effects of membrane and soluble Fas ligand on cardiomyocytes: role in ischemia/reperfusion injury. *J Mol Cell Cardiol* 2003, **35**:811–821.
33. Hohlbaum AM, Moe S, Marshak-Rothstein A: Opposing effects of transmembrane and soluble Fas ligand expression on inflammation and tumor cell survival. *J Exp Med* 2000, **191**:1209–1220.
34. Wajant H, Moosmayer D, Wuest T, Bartke T, Gerlach E, Schonherr U, Peters N, Scheurich P, Pfizenmaier K: Differential activation of TRAIL-R1 and -2 by soluble and membrane TRAIL allows selective surface antigen-directed activation of TRAIL-R2 by a soluble TRAIL derivative. *Oncogene* 2001, **20**:4101–4106.
35. Rodriguez-Bermejo G, Fraile B, Paniagua R, Aller P, Royuela M: Expression of NF-kappaB-related proteins and their modulation during TNF-alpha-provoked apoptosis in prostate cancer cells. *Prostate* 2012, **72**:40–50.
36. Gupta S: A decision between life and death during TNF-alpha-induced signaling. *J Clin Immunol* 2002, **22**:185–194.
37. Grell M, Scheurich P, Meager A, Pfizenmaier K: TR60 and TR80 tumor necrosis factor (TNF)-receptors can independently mediate cytosis. *Lymphokine Cytokine Res* 1993, **12**:143–148.
38. Lee SY, Park CG, Choi Y: T cell receptor-dependent cell death of T cell hybridomas mediated by the CD30 cytoplasmic domain in association with tumor necrosis factor receptor-associated factors. *J Exp Med* 1996, **183**:669–674.
39. Depuydt B, van Loo G, Vandenabeele P, Declercq W: Induction of apoptosis by TNF receptor 2 in a T-cell hybridoma is FADD dependent and blocked by caspase-8 inhibitors. *J Cell Sci* 2005, **118**:497–504.
40. Teh HS, Seebaran A, Teh SJ: TNF receptor 2-deficient CD8 T cells are resistant to Fas/Fas ligand-induced cell death. *J Immunol* 2000, **165**:4814–4821.
41. Grell M, Douni E, Wajant H, Lohden M, Clauss M, Maxeiner B, Georgopoulos S, Lesslauer W, Kollrias G, Pfizenmaier K, Scheurich P: The transmembrane form of tumor necrosis factor is the prime activating ligand of the 80 kDa tumor necrosis factor receptor. *Cell* 1995, **83**:793–802.
42. Grassme H, Jendrossek V, Bock J, Riehle A, Gulbins E: Ceramide-rich membrane rafts mediate CD40 clustering. *J Immunol* 2002, **168**:298–307.

43. Gudz TI, Tserng KY, Hoppel CL: Direct inhibition of mitochondrial respiratory chain complex III by cell-permeable ceramide. *J Biol Chem* 1997, **272**:24154–24158.
44. Zhang H, Yan D, Shi X, Liang H, Pang Y, Qin N, Chen H, Wang J, Yin B, Jiang X, et al: Transmembrane TNF-alpha mediates “forward” and “reverse” signaling, inducing cell death or survival via the NF-kappaB pathway in Raji Burkitt lymphoma cells. *J Leukoc Biol* 2008, **84**:789–797.

doi:10.1186/1750-2187-8-12

**Cite this article as:** Ardestani et al.: Membrane TNF-alpha-activated programmed necrosis is mediated by Ceramide-induced reactive oxygen species. *Journal of Molecular Signaling* 2013 **8**:12.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

