

Complete genome sequence of *Enterococcus faecium* strain AK_C_05 with potential characteristics applicable in livestock industry

Hyunok Doo^{1#}, Jin Ho Cho^{2#}, Minho Song^{3#}, Eun Sol Kim¹, Sheena Kim¹, Gi Beom Keum¹, Jinok Kwak¹, Srinivas Pandey¹, Sumin Ryu¹, Yejin Choi¹, Juyoun Kang¹, Hyeun Bum Kim^{1*} and Ju-Hoon Lee^{4,5,6*}

¹Department of Animal Biotechnology, Dankook University, Cheonan 31116, Korea

²Division of Food and Animal Science, Chungbuk National University, Cheongju 28644, Korea

³Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea

⁴Department of Food Animal Biotechnology, Seoul National University, Seoul 08826, Korea

⁵Department of Agricultural Biotechnology, Seoul National University, Seoul 08826, Korea

⁶Center for Food and Bioconvergence, Seoul National University, Seoul 08826, Korea



Received: Aug 2, 2023
 Revised: Sep 5, 2023
 Accepted: Sep 25, 2023

#These authors contributed equally to this work.

*Corresponding author

Hyeun Bum Kim
 Department of Animal Biotechnology,
 Dankook University, Cheonan 31116,
 Korea.
 Tel: +82-41-550-3653
 E-mail: hbkim@dankook.ac.kr

Ju-Hoon Lee
 Department of Agricultural
 Biotechnology, Seoul National
 University, Seoul 08826, Korea.
 Tel: +82-2-880-4854
 E-mail: juhlee@snu.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Hyunok Doo
<https://orcid.org/0000-0003-4329-4128>

Abstract

The *Enterococcus faecium* (*E. faecium*) strain AK_C_05 was isolated from *cheonggukjang*, the Korean traditional food, collected from a local market in South Korea. In this report, we presented the complete genome sequence of *E. faecium* strain AK_C_05. The genome of *E. faecium* strain AK_C_05 genome consisted of one circular chromosome (2,691,319 bp) with a guanine + cytosine (GC) content of 38.3% and one circular plasmid (177,732 bp) with a GC content of 35.48%. The Annotation results revealed 2,827 protein-coding sequences (CDSs), 18 rRNAs, and 68 tRNA genes. It possesses genes, which encodes enzymes such as alpha-galactosidase (EC 3.2.1.22), beta-glucosidase (EC 3.2.1.21) and alpha-L-arabinofuranosidase (EC 3.2.1.55) enabling efficient utilization of carbohydrates. Based on Clusters of Orthologous Groups analysis, *E. faecium* strain AK_C_05 showed specialization in carbohydrate transport and metabolism indicating the ability to generate energy using a variety of carbohydrates.

Keywords: *Enterococcus faecium*, Livestock, Carbohydrates

The Enterococci bacteria belong to lactic acid bacteria (LAB) group, which can be found in fermented foods [1]. Especially, *Enterococcus faecium* is also utilized as probiotics, which could enhance the microbial balance in animals [2]. Despite of safety concerns regarding its use as probiotics, recent research has explored the use of *Enterococcus faecium* as a feed additive for livestock to enhance growth performance [1,3].

In the present study, the *E. faecium* strain AK_C_05 was isolated from homemade *cheonggukjang*, the Korean traditional food, collected from a local market in Cheonan (36.802917° N, 127.149796° E), Chungcheongnam-do, South Korea. Then, the whole genome sequencing was performed to understand the genomic characteristics of *E. faecium* strain AK_C_05 as a potential probiotic in

Jin Ho Cho
<https://orcid.org/0000-0001-7151-0778>
 Minh Song
<https://orcid.org/0000-0002-4515-5212>
 Eun Sol Kim
<https://orcid.org/0000-0001-8801-421X>
 Sheena Kim
<https://orcid.org/0000-0002-5410-1347>
 Gi Beom Keum
<https://orcid.org/0000-0001-6006-9577>
 Jinok Kwak
<https://orcid.org/0000-0003-1217-3569>
 Srinivas Pandey
<https://orcid.org/0000-0002-6947-3469>
 Sumin Ryu
<https://orcid.org/0000-0002-1569-3394>
 Yejin Choi
<https://orcid.org/0000-0002-7434-299X>
 Juyoun Kang
<https://orcid.org/0000-0002-3974-2832>
 Hyeun Bum Kim
<https://orcid.org/0000-0003-1366-6090>
 Ju-Hoon Lee
<https://orcid.org/0000-0003-0405-7621>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This research was supported by a grant (22193MFDS538) from Ministry of Food and Drug Safety in 2022.

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Doo H, Kim HB, Lee JH.
 Data curation: Keum GB, Choi Y, Kang J.
 Formal analysis: Kim ES, Kim S, Keum GB, Ryu S.
 Methodology: Cho JH, Song M.
 Validation: Kim S, Kwak J, Pandey S.
 Writing - original draft: Doo H, Cho JH, Song M.
 Writing - review & editing: Doo H, Cho JH, Song M, Kim ES, Kim S, Keum GB, Kwak J, Pandey S, Ryu S, Choi Y, Kang J, Kim HB, Lee JH.

Ethics approval and consent to participate

This article does not require IRB/ACUC approval because there are no human and animal participants.

the livestock industry. The *E. faecium* strain AK_C_05 was cultivated in Enterococcosel broth (MBcell, Seoul, South Korea) at 37°C for 24 hours. Genomic DNA was extracted from the cultured *E. faecium* pellet using CTAB DNA extraction method. The complete genome of the *E. faecium* AK_C_05 was sequenced using the Oxford Nanopore Technologies MinION platform at eGnome (Seoul, South Korea). Briefly, library preparation was performed using Native barcoding Sequencing Kit (SQK_NBD114.24, V14) following the manufacturer's instructions (Oxford Nanopore Technologies, Oxford, UK). The prepared library was loaded into the MinION MK1b sequencing device (Oxford Nanopore) equipped with a MinION flow cell (MIN114, R10.4.1, Oxford Nanopore). The Oxford Nanopore sequencing produced 79,247 of long reads, resulting in a total of 572,297,864 base pairs. De novo assemble was performed using a Flye assembler v2.9.2, followed by polishing using the Homopolish polisher v0.4.1. The quality of genome assembly was assessed using Quality Assessment Tool for Genome Assemblies (QUAST) v5.2.0 [4]. The quantitative assessment of the genome completeness was conducted using the Benchmarking Universal Single-Copy Orthologs (BUSCO) v5.4.6 [5]. To annotate and predict the protein coding genes, rRNA, and tRNA genes of *E. faecium* strain AK_C_05, the Rapid Annotation using Subsystem Technology (RAST) v2.0 tool was utilized [6]. The functional categorization of all predicted protein coding genes was performed using the Clusters of Orthologous Groups (COGs)-based EggNOG-mapper v2.0 [7]. Furthermore, the presence of virulence factors and antibiotic resistance in *E. faecium* strain AK_C_05 was predicted using the BLASTn method, with reference to the Virulence Factor Database (VFDB) and the Comprehensive Antibiotic Resistance Database (CARD) [8,9].

The complete genome of the *E. faecium* strain AK_C_05 contain one circular chromosome (2,691,319 bp) with a guanine + cytosine (GC) content of 38.3% and one circular plasmid (177,732 bp) with a GC content of 35.48%. A total of 2,827 predicted protein-coding sequence, 18 rRNA genes, and 68 tRNA genes were identified in *E. faecium* strain AK_C_05. The most abundant COGs category, excluding Function unknown [S], was Carbohydrate transport and metabolism [G], which accounted for 235 genes, representing 10.4% of the total genes identified. The genome feature and map of *E. faecium* strain AK_C_05 were presented in Table 1, Figs. 1A and 1B.

Based on its specific focus on carbohydrate transport and metabolism, *E. faecium* strain AK_C_05 possesses genes and enzymes, such as alpha-galactosidase (EC 3.2.1.22), beta-glucosidase (EC 3.2.1.21) and alpha-L-arabinofuranosidase (EC 3.2.1.55), that enable efficient utilization of carbohydrates and the capacity to derive energy from diverse carbohydrate substrates. This characteristic makes *E. faecium* strain AK_C_05 a potential candidate for application in the livestock industry. The complete genome of *E. faecium* strain AK_C_05 has indicated the presence of the antibiotic resistance gene *aac* (6')-II in the chromosome and not in the plasmid, confirming

Table 1. Genome features of *Enterococcus faecium* strain AK_C_05

Property	Term	
	Chromosome	Plasmid
Contig length (bp)	2,691,319 bp	177,732 bp
No. of contig	1 (chromosome)	1 (plasmid)
Guanine + cytosine (G + C)	38.3	35.48
Protein-coding genes	2,629	198
rRNA genes	18	-
tRNA genes	68	-
Genbank Accession No.	CP128995	CP128994




Fig. 1. Genome map of *Enterococcus faecium* strain AK_C_05 and the functional categorization of predicted protein coding genes. The outer ring represents the positions of all annotated gene coding regions (ORFs), while the inner ring in red indicates the guanine + cytosine (GC) content. Peaks in pink and green indicate GC skew. The orange and sky-blue arrows represent rRNA and tRNA operons, respectively. (A) The annotated ORFs are color-coded based on their Clusters of Orthologous Groups (COG) assignments. (B) The COG functional categories of the predicted protein coding genes are represented.

that there is no potential for transmission of the resistance gene to other microorganisms. In the plasmid of *E. faecium* strain AK_C_05, the *filA* gene was detected, while no other virulence factors were identified. Interestingly, the *filA* gene's ability to facilitate adhesion to the cell wall is regarded as a beneficial trait for probiotics [10]. Overall, our results indicate that *E. faecium* AK_C_05 could be a promising functional probiotic for improving growth performance in the livestock industry.

NUCLEOTIDE SEQUENCE ACCESSION NUMBER(S)

The complete genome sequences of *Enterococcus faecium* strain AK_C_05 were deposited in GeneBank under the accession numbers CP128994.1 and CP128995.1. The BioSample accession number is SAMN35654454, and BioProject accession number is PRJNA980926.

REFERENCES

1. Franz CMAP, Stiles ME, Schleifer KH, Holzapfel WH. Enterococci in foods—a conundrum for food safety. *Int J Food Microbiol.* 2003;88:105-22. [https://doi.org/10.1016/S0168-1605\(03\)00174-0](https://doi.org/10.1016/S0168-1605(03)00174-0)
2. Bhardwaj A, Gupta H, Kapila S, Kaur G, Vij S, Malik RK. Safety assessment and evaluation of probiotic potential of bacteriocinogenic *Enterococcus faecium* KH 24 strain under in vitro and in vivo conditions. *Int J Food Microbiol.* 2010;141:156-64. <https://doi.org/10.1016/j.ijfoodmicro.2010.05.001>
3. Sato Y, Kuroki Y, Oka K, Takahashi M, Rao S, Sukegawa S, et al. Effects of dietary supplementation with *Enterococcus faecium* and *Clostridium butyricum*, either alone or in combination, on growth and fecal microbiota composition of post-weaning pigs at a commercial farm. *Front Vet Sci.* 2019;6:26. <https://doi.org/10.3389/fvets.2019.00026>
4. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome assemblies. *Bioinformatics.* 2013;29:1072-5. <https://doi.org/10.1093/bioinformatics/btt086>
5. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics.* 2015;31:3210-2. <https://doi.org/10.1093/bioinformatics/btv351>
6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST server: rapid annotations using subsystems technology. *BMC Genom.* 2008;9:75. <https://doi.org/10.1186/1471-2164-9-75>
7. Cantalapiedra CP, Hernández-Plaza A, Letunic I, Bork P, Huerta-Cepas J. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol Biol Evol.* 2021;38:5825-9. <https://doi.org/10.1093/molbev/msab293>
8. Chen L, Yang J, Yu J, Yao Z, Sun L, Shen Y, et al. VFDB: a reference database for bacterial virulence factors. *Nucleic Acids Res.* 2005;33:D325-8. <https://doi.org/10.1093/nar/gki008>
9. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, et al. The comprehensive antibiotic resistance database. *Antimicrob Agents Chemother.* 2013;57:3348-57. <https://doi.org/10.1128/aac.00419-13>
10. Li B, Zhan M, Evivie SE, Jin D, Zhao L, Chowdhury S, et al. Evaluating the safety of potential probiotic *Enterococcus durans* KLDS6.0930 using whole genome sequencing and oral toxicity study. *Front Microbiol.* 2018;9:1943. <https://doi.org/10.3389/fmicb.2018.01943>