

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Isopropyl-4-methoxy-5-methylphenyl acetate

 Radouane Oubabi,<sup>a</sup> Aziz Auhmani,<sup>a</sup> My Youssef Ait Itto,<sup>a</sup> Mohamed Driss<sup>b</sup> and El Hassane Soumhi<sup>c\*</sup>
<sup>a</sup>Laboratoire de Synthèse Organique et Physico-Chimie Moléculaire, Faculté des Sciences-Semlalia, Université Cadi Ayyad, BP 2390, 40001, Marrakech, Morocco,

<sup>b</sup>Laboratoire de Matériaux et Cristallographie, Faculté des Sciences de Tunis, Université de Tunis ElManar, 2092 ElManar II, Tunis, Tunisia, and <sup>c</sup>Equipe de Chimie des Matériaux et de l'Environnement, FSTG-Marrakech, Université Cadi Ayyad, Bd Abdelkrim Khattabi, BP 549, Marrakech, Morocco

Correspondence e-mail: eh\_soumhi@yahoo.fr


Received 9 October 2013; accepted 23 October 2013

 Key indicators: single-crystal X-ray study;  $T = 300$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.046;  $wR$  factor = 0.132; data-to-parameter ratio = 18.4.

In the title compound,  $\text{C}_{13}\text{H}_{18}\text{O}_3$ , the benzene ring is almost perpendicular to the acetoxy plane, making a dihedral angle of  $89.33$  ( $11$ )°. In the crystal, molecules are linked by weak  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, forming a zigzag chain along the  $c$ -axis direction.

### Related literature

For background to natural monoterpenic phenols and their derivatives, see: Yuan-Lang & Erdtman (1962); Ündeğer *et al.* (2009); Osorio *et al.* (2006). For a related structure, see: Rajouani *et al.* (2008).



### Experimental

#### Crystal data

$\text{C}_{13}\text{H}_{18}\text{O}_3$   
 $M_r = 222.27$   
 Monoclinic,  $P2_1/c$   
 $a = 10.829$  (2) Å  
 $b = 9.600$  (2) Å  
 $c = 12.530$  (3) Å  
 $\beta = 100.34$  (2)°  
 $V = 1281.4$  (5) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.08$  mm<sup>-1</sup>  
 $T = 300$  K  
 $0.3 \times 0.15 \times 0.1$  mm

#### Data collection

Enraf–Nonius CAD-4 diffractometer  
 Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\text{min}} = 0.521$ ,  $T_{\text{max}} = 0.992$   
 3366 measured reflections  
 2780 independent reflections  
 1594 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.018$   
 2 standard reflections every 60 min  
 intensity decay: 1%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.132$   
 $S = 1.02$   
 2780 reflections  
 151 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.16$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.13$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C3}-\text{H1}\cdots\text{O2}^i$	0.93	2.60	3.523 (3)	171

 Symmetry code: (i)  $x, -y - \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS*; data reduction: *MolEN* (Fair, 1990); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5314).

### References

- Enraf–Nonius (1989). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.  
 Fair, C. K. (1990). *MolEN*. Enraf–Nonius, Delft, The Netherlands.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
 Osorio, E., Arango, G., Robledo, S., Muñoz, D., Jaramillo, L. & Vélez, I. (2006). *Acta Farm. Bonaer.* **25**, 405–413.  
 Rajouani, N., Ait Itto, My. Y., Benharref, A., Auhmani, A. & Daran, J.-C. (2008). *Acta Cryst.* **E64**, o762.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Ündeğer, Ü., Başaran, A., Degen, G. H. & Başaran, N. (2009). *Food Chem. Toxicol.* **47**, 2037–2043.  
 Yuan-Lang, C. & Erdtman, H. (1962). *Acta Chem. Scand.* **16**, 1291–1295.

## supplementary materials

*Acta Cryst.* (2013). E69, o1715 [doi:10.1107/S160053681302922X]

## 2-Isopropyl-4-methoxy-5-methylphenyl acetate

Radouane Oubabi, Aziz Auhmani, My Youssef Ait Itto, Mohamed Driss and El Hassane Soumhi

### 1. Comment

The *p*-methoxythymol is a natural phenol extracted from heartwood of *tetraclinis articulate* (Yuan-Lang *et al.*, 1962). In several recent studies, natural monoterpenic phenols and their derivatives have revealed extensive biological activities, ranging from antimicrobial (Ündeğer *et al.*, 2009) to Antileishmanial and Cytotoxic activities (Osorio *et al.*, 2006). In the aim of preparing monoterpenic phenol derivatives, we report here, the hemisynthesis of 2-isopropyl-4-methoxy-5-methylphenyl acetate from naturally occurred *p*-methoxythymol. Thus, treatment of *p*-methoxythymol with acetic anhydride in pyridine, and provides the title compound as colorless crystals in 72.4% yield. Its structure was fully characterized by its mass and NMR spectroscopic data. Furthermore, the crystallographic study made it possible to determine the stereochemistry. The main geometric features of this group are in good agreement with those observed in a similar compound (Rajouani *et al.*, 2008).

### 2. Experimental


A solution of *p*-methoxythymol (111 mg, 0.617 mmol) in acetic anhydride (20 ml) and pyridine (20 ml) was heated under reflux for 24 h. After cooling, the mixture was acidified with 1 N HCl solution then extracted with ether (3 × 20 ml). The organic layer was washed with water, dried on anhydrous Na<sub>2</sub>SO<sub>4</sub> and then evaporated under reduced pressure. The obtained residue was chromatographed on silica gel column using hexane and ethyl acetate (97/3) as eluent, to give 2-isopropyl-4-methoxy-5-methylphenyl acetate (100 mg) in 72.4% yield.

### 3. Refinement

All H-atoms were located in a difference map and refined using a riding model with C—H = 0.93–0.98 Å, and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Computing details

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); cell refinement: *CAD-4 EXPRESS* (Enraf–Nonius, 1989); data reduction: *MolEN* (Fair, 1990); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *WinGX* (Farrugia, 2012).


**Figure 1**

The molecule structure of the title compound with 50% probability ellipsoids.

## 2-Isopropyl-4-methoxy-5-methylphenyl acetate

### Crystal data

$C_{13}H_{18}O_3$

$M_r = 222.27$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2ybc$

$a = 10.829\ (2)\ \text{\AA}$

$b = 9.600\ (2)\ \text{\AA}$

$c = 12.530\ (3)\ \text{\AA}$

$\beta = 100.34\ (2)^\circ$

$V = 1281.4\ (5)\ \text{\AA}^3$

$Z = 4$

$F(000) = 480$

$D_x = 1.152\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 10\text{--}15^\circ$

$\mu = 0.08\ \text{mm}^{-1}$

$T = 300\ \text{K}$

Prism, colourless

$0.3 \times 0.15 \times 0.1\ \text{mm}$

### Data collection

Enraf-Nonius CAD-4

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega/2\theta$  scans

Absorption correction:  $\psi$  scan

(North *et al.*, 1968)

$T_{\min} = 0.521$ ,  $T_{\max} = 0.992$

3366 measured reflections

2780 independent reflections

1594 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

$\theta_{\max} = 27.0^\circ$ ,  $\theta_{\min} = 2.7^\circ$

$h = -13 \rightarrow 1$

$k = -1 \rightarrow 12$

$l = -15 \rightarrow 15$

2 standard reflections every 60 min

intensity decay: 1%

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.046$   
 $wR(F^2) = 0.132$   
 $S = 1.02$   
 2780 reflections  
 151 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: difference Fourier map  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0525P)^2 + 0.2201P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.13 \text{ e } \text{\AA}^{-3}$   
 Extinction correction: *SHELXL97* (Sheldrick,  
 2008),  $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.016 (3)

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.76460 (12)	0.08305 (13)	0.33870 (9)	0.0587 (4)
O2	0.76646 (18)	-0.07552 (19)	0.46746 (13)	0.1062 (7)
O3	0.43208 (12)	-0.20942 (15)	0.03401 (10)	0.0673 (4)
C1	0.68418 (16)	-0.00154 (18)	0.26312 (13)	0.0496 (4)
C2	0.73387 (16)	-0.09160 (17)	0.19660 (13)	0.0476 (4)
C3	0.64806 (16)	-0.16266 (18)	0.11839 (14)	0.0522 (4)
H1	0.6777	-0.2242	0.0715	0.063*
C4	0.52031 (16)	-0.14316 (18)	0.10945 (13)	0.0514 (4)
C5	0.47183 (17)	-0.0533 (2)	0.17877 (14)	0.0562 (5)
C6	0.55666 (17)	0.0175 (2)	0.25510 (14)	0.0570 (5)
H2	0.5272	0.0792	0.3020	0.068*
C7	0.79534 (17)	0.0364 (2)	0.44091 (15)	0.0612 (5)
C8	0.8695 (2)	0.1412 (3)	0.51249 (17)	0.0803 (7)
H3	0.8841	0.1081	0.5860	0.096*
H4	0.8239	0.2274	0.5082	0.096*
H5	0.9485	0.1559	0.4896	0.096*
C9	0.4748 (2)	-0.2910 (2)	-0.04666 (16)	0.0742 (6)
H6	0.5283	-0.2355	-0.0830	0.089*
H7	0.4041	-0.3229	-0.0983	0.089*
H8	0.5210	-0.3697	-0.0132	0.089*
C10	0.33271 (18)	-0.0334 (3)	0.16960 (19)	0.0812 (7)
H9	0.3168	0.0323	0.2233	0.097*
H10	0.2943	-0.1210	0.1810	0.097*
H11	0.2982	0.0011	0.0986	0.097*

C11	0.87402 (16)	-0.11360 (19)	0.20594 (15)	0.0546 (5)
H12	0.9162	-0.0529	0.2641	0.066*
C12	0.9114 (2)	-0.2629 (2)	0.23722 (18)	0.0747 (6)
H13	0.8718	-0.3250	0.1813	0.090*
H14	0.8850	-0.2857	0.3043	0.090*
H15	1.0009	-0.2723	0.2458	0.090*
C13	0.9195 (2)	-0.0725 (3)	0.10231 (18)	0.0785 (6)
H16	1.0093	-0.0797	0.1133	0.094*
H17	0.8949	0.0218	0.0837	0.094*
H19	0.8830	-0.1335	0.0445	0.094*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0685 (8)	0.0536 (7)	0.0512 (7)	-0.0086 (6)	0.0036 (6)	-0.0056 (6)
O2	0.1266 (15)	0.1007 (13)	0.0758 (10)	-0.0443 (11)	-0.0236 (10)	0.0304 (10)
O3	0.0564 (7)	0.0792 (9)	0.0621 (8)	-0.0073 (7)	-0.0007 (6)	-0.0123 (7)
C1	0.0553 (10)	0.0460 (10)	0.0448 (9)	-0.0017 (8)	0.0022 (8)	0.0014 (8)
C2	0.0509 (9)	0.0444 (9)	0.0464 (9)	-0.0011 (8)	0.0054 (7)	0.0033 (8)
C3	0.0574 (11)	0.0510 (10)	0.0475 (9)	-0.0004 (8)	0.0079 (8)	-0.0034 (8)
C4	0.0508 (10)	0.0521 (10)	0.0481 (9)	-0.0037 (8)	0.0006 (8)	0.0019 (8)
C5	0.0516 (10)	0.0583 (11)	0.0562 (10)	0.0076 (9)	0.0030 (8)	0.0052 (9)
C6	0.0624 (11)	0.0543 (11)	0.0543 (11)	0.0077 (9)	0.0108 (9)	-0.0040 (9)
C7	0.0533 (11)	0.0727 (13)	0.0545 (11)	-0.0043 (10)	0.0013 (9)	0.0009 (10)
C8	0.0706 (13)	0.0957 (17)	0.0675 (13)	-0.0095 (12)	-0.0071 (11)	-0.0155 (12)
C9	0.0796 (14)	0.0819 (15)	0.0586 (11)	-0.0200 (12)	0.0054 (10)	-0.0148 (11)
C10	0.0567 (12)	0.0940 (17)	0.0900 (16)	0.0120 (12)	0.0052 (11)	-0.0100 (13)
C11	0.0504 (10)	0.0560 (11)	0.0559 (10)	-0.0048 (9)	0.0053 (8)	-0.0048 (9)
C12	0.0615 (12)	0.0719 (14)	0.0860 (15)	0.0071 (11)	0.0006 (11)	0.0076 (12)
C13	0.0652 (13)	0.0903 (16)	0.0839 (15)	-0.0030 (12)	0.0236 (11)	0.0106 (13)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

O1—C7	1.341 (2)	C11—C12	1.522 (3)
O1—C1	1.421 (2)	C8—H3	0.9600
O2—C7	1.183 (2)	C8—H4	0.9600
O3—C4	1.374 (2)	C8—H5	0.9600
O3—C9	1.420 (2)	C9—H6	0.9600
C1—C2	1.376 (2)	C9—H7	0.9600
C1—C6	1.379 (2)	C9—H8	0.9600
C2—C3	1.401 (2)	C10—H9	0.9600
C2—C11	1.516 (2)	C10—H10	0.9600
C3—C4	1.380 (2)	C10—H11	0.9600
C3—H1	0.9300	C11—H12	0.9800
C4—C5	1.392 (3)	C12—H13	0.9600
C5—C6	1.380 (2)	C12—H14	0.9600
C5—C10	1.502 (3)	C12—H15	0.9600
C6—H2	0.9300	C13—H16	0.9600
C7—C8	1.484 (3)	C13—H17	0.9600
C11—C13	1.521 (3)	C13—H19	0.9600

C7—O1—C1	117.57 (14)	O3—C9—H7	109.5
C4—O3—C9	118.02 (15)	H6—C9—H7	109.5
C2—C1—C6	122.33 (16)	O3—C9—H8	109.5
C2—C1—O1	120.23 (15)	H6—C9—H8	109.5
C6—C1—O1	117.31 (16)	H7—C9—H8	109.5
C1—C2—C3	116.58 (16)	C5—C10—H9	109.5
C1—C2—C11	122.40 (15)	C5—C10—H10	109.5
C3—C2—C11	121.02 (16)	H9—C10—H10	109.5
C4—C3—C2	121.33 (17)	C5—C10—H11	109.5
C4—C3—H1	119.3	H9—C10—H11	109.5
C2—C3—H1	119.3	H10—C10—H11	109.5
O3—C4—C3	123.79 (16)	C2—C11—C13	111.77 (15)
O3—C4—C5	115.01 (16)	C2—C11—C12	111.54 (15)
C3—C4—C5	121.20 (16)	C13—C11—C12	110.65 (17)
C6—C5—C4	117.32 (16)	C2—C11—H12	107.5
C6—C5—C10	121.61 (18)	C13—C11—H12	107.5
C4—C5—C10	121.07 (17)	C12—C11—H12	107.5
C1—C6—C5	121.23 (17)	C11—C12—H13	109.5
C1—C6—H2	119.4	C11—C12—H14	109.5
C5—C6—H2	119.4	H13—C12—H14	109.5
O2—C7—O1	122.63 (18)	C11—C12—H15	109.5
O2—C7—C8	125.96 (19)	H13—C12—H15	109.5
O1—C7—C8	111.41 (18)	H14—C12—H15	109.5
C7—C8—H3	109.5	C11—C13—H16	109.5
C7—C8—H4	109.5	C11—C13—H17	109.5
H3—C8—H4	109.5	H16—C13—H17	109.5
C7—C8—H5	109.5	C11—C13—H19	109.5
H3—C8—H5	109.5	H16—C13—H19	109.5
H4—C8—H5	109.5	H17—C13—H19	109.5
O3—C9—H6	109.5		
C7—O1—C1—C2	-95.9 (2)	C3—C4—C5—C6	1.3 (3)
C7—O1—C1—C6	88.2 (2)	O3—C4—C5—C10	0.3 (3)
C6—C1—C2—C3	0.8 (3)	C3—C4—C5—C10	-179.34 (18)
O1—C1—C2—C3	-174.89 (14)	C2—C1—C6—C5	-0.3 (3)
C6—C1—C2—C11	-179.47 (17)	O1—C1—C6—C5	175.57 (16)
O1—C1—C2—C11	4.8 (2)	C4—C5—C6—C1	-0.8 (3)
C1—C2—C3—C4	-0.3 (2)	C10—C5—C6—C1	179.85 (19)
C11—C2—C3—C4	179.98 (16)	C1—O1—C7—O2	6.1 (3)
C9—O3—C4—C3	-7.0 (3)	C1—O1—C7—C8	-174.65 (16)
C9—O3—C4—C5	173.38 (16)	C1—C2—C11—C13	-118.73 (19)
C2—C3—C4—O3	179.64 (15)	C3—C2—C11—C13	61.0 (2)
C2—C3—C4—C5	-0.8 (3)	C1—C2—C11—C12	116.80 (19)
O3—C4—C5—C6	-179.06 (15)	C3—C2—C11—C12	-63.5 (2)

Hydrogen-bond geometry (Å, °)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C3-H1\cdots O2^i$	0.93	2.60	3.523 (3)	171

Symmetry code: (i)  $x, -y-1/2, z-1/2$ .