

The Pol-e-Pill Finally Arrives

James A. Levine¹ and Ronald M. Davis²

The global obesity epidemic affects billions of people (1), and, in its wake, a worldwide epidemic of type 2 diabetes has followed (2,3). Obesity is not only associated with diabetes, of course, but also with the other partners in the X-gang: hypertension, hyperlipidemia, and cardiovascular disease (4). As exemplified in an original article in this issue of *Diabetes* (5), the link between diabetes and obesity continues to be scrutinized in detail. In the wake of this metabolic catastrophe, the ill effects of joint pain, degenerative arthritis, and sleep apnea ensue (6). A ripple of despair similarly washes over the oncology community, who now realize that the global tragedy of obesity claims cancer lives too (7–10). “But how about us?” the immunologists (11) and vascular (4) specialists cry; obesity dulls the body’s response to infection, impairs wound healing, and, consequently, diminishes reproductive capacity (12).

All of this is important, but none of it matters to Keith Mitchell, who expresses his frustration to his impotent physician, “Doc, I’ve tried every diet on earth. Why can’t you help me?” (13). Keith returns home bereft of hope to later apologize to his children, with whom he is too tired to play (14), and to his wife, whom he feels ill adept to love.

Obesity touches every aspect of society, from the economic viability of nations (15) to the look of defeat reflected in the eyes of Keith’s daughter as she gazes upon her broken father (16). The obesity epidemic accounted for 27% of the increase in health care spending in the U.S. between 1987 and 2001 (17). Is it any surprise that we in the medical community feel despair at our incapacity to provide scalable real-world solutions to the 300 trillion excess calories stored in the human fuel reserve and its associated torrent of disease? (An average person with a BMI of 22 kg/m² has 13 kg of fat mass, and an average person with a BMI of 33 kg/m² has 36 kg of fat mass. The difference is 20,000 g or 180,000 kcal. With 1.5 billion [1,500,000,000] people in the world with obesity, the excess stored calories approximate 270,000,000,000,000 kcal.)

The notion of a polypill to prevent X-gang disease has been touted for a decade (18–24). The most popular variant of this concept is a tablet that people take as early

as childhood to prevent the onset of metabolic diseases and even obesity itself. If such a tablet were available, clinically effective, safe, and affordable, we would run out of ink (or toner cartridges) prescribing it. Prescribing such a polypill would become the mandate of thousands of health care professionals worldwide. We would all sleep better at night, knowing that Keith was.

What would be the mode of action of such a polypill? A polypill might function via several pathways such as through molecular targets (e.g., the insulin/insulin-receptor pathways) (25), on specific tissues such as muscle and adipose (26), and/or through neuronal (26) and humeral effectors (27). Because the discovery path for a single pill with such broad-ranging targets is prolonged and exhaustive (28), a polypill that contains multiple, preexisting chemicals is a more logical approach for preventing metabolic disease and obesity.

The problem with a broadly functioning, chemically complex polypill is that side effects and multiple interactions with other medications are likely to result once global deployment has been achieved. However, like crossing the road, we must consider the benefit-to-risk ratio. Understanding that more than 5% of the U.S. population has diabetes and, for the first time in history, at least one or two of Keith’s daughter’s friends will develop type 2 diabetes (2,3), what safety profile should we tolerate? A few deaths sacrificed for the salvation of the many is the burden long born by federal agencies such as the Food and Drug Administration (29).

With the magnitude of obesity-associated health problems and the impact of obesity on health care costs, cogent arguments for urgent implementation of an effective polypill are easy proffered. It is therefore with delight, at the dawn of a new year, that we are able to announce the release to market of an effective Pol-e-pill. The Pol-e-pill, discovered two centuries ago (30), has been independently tested at multiple centers across the world and has been the focus of numerous publications and government reports (31–37). The Pol-e-pill’s mode of action affects multiple molecular targets and it is efficacious at squelching the gang of four (diabetes, hypertension, hyperlipidemia, and cardiovascular disease) (38–41), positively affecting multiple other obesity-associated comorbidities and, when prescribed at a high enough dose, decreasing body weight (35,42–45).

The Pol-e-pill has an excellent dose-response curve (46) and a superb safety profile. At low doses, the Pol-e-pill is essentially free of side effects (although in the unwell, the low-dose Pol-e-pill needs to be carefully monitored). However, at supra-optimal doses, the Pol-e-pill does have an increased incidence of adverse effects, which can be ameliorated by decreasing the dose (47). Better still, the Pol-e-pill is not constrained through any active patents, and so its cost is low. The Pol-e-pill can be potentially prescribed across the world today.

So, it is finally here—Keith’s dream—an effective “prescribe-able” solution for preventing and treating diabetes,

From the ¹Department of Internal Medicine, Mayo Clinic, Rochester, Minnesota; and the ²Center for Health Promotion and Disease Prevention, Henry Ford Health System, Detroit, Michigan.

Corresponding author: James A. Levine, levine.james@mayo.edu.

Received 24 January 2008 and accepted 27 January 2008.

R.M.D. is president of the American Medical Association.

DOI: 10.2337/db08-0108

© 2008 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

See accompanying article, p. 1834.

obesity, and most obesity-assisted complications. The dose-response effects, safety profile, and low cost render the Pol-e-pill amenable to worldwide prescription. Check the cartridge in your Pharma-labeled pen and the toner level in your printer—Keith is on his way.

The Pol-e-pill is exercise. Prescribe it. The package insert, with details on indications and contraindications, benefits and risks, dosage and administration, supporting evidence, and additional resources for clinicians and patients, is available at <http://exercisemedicine.org/>. We urge physicians to prescribe exercise just as they would a more costly tablet.

REFERENCES

- James PT, Leach R, Kalamara E, Shayeghi M: The worldwide obesity epidemic. *Obes Res* 9:228S–233S, 2001
- Meetoo D, McGovern P, Safadi R: An epidemiological overview of diabetes across the world. *Br J Nurs* 16:1002–1007, 2007
- Gesta S, Tseng YH, Kahn CR: Developmental origin of fat: tracking obesity to its source. *Cell* 131:242–256, 2007
- Vongpatanasin W: Cardiovascular morbidity and mortality in high-risk populations: epidemiology and opportunities for risk reduction. *J Clin Hypertens (Greenwich)* 9:11–15, 2007
- Jocken JWE, Roepstorff C, Goossens GH, van der Baan P, van Baak M, Saris WHM, Kiens B, Blaak EE: Hormone-sensitive lipase serine phosphorylation and glycerol exchange across skeletal muscle in lean and obese subjects: effect of β -adrenergic stimulation. *Diabetes* 57:1834–1841
- Christensen R, Bartels EM, Astrup A, Bliddal H: Effect of weight reduction in obese patients diagnosed with knee osteoarthritis: a systematic review and meta-analysis. *Ann Rheum Dis* 66:433–439, 2007
- Hursting SD, Lashinger LM, Colbert LH, Rogers CJ, Wheatley KW, Nunez NP, Mahabir S, Barrett JC, Forman MR, Perkin SN: Energy balance and carcinogenesis: underlying pathways and targets for intervention. *Curr Cancer Drug Targets* 7:484–491, 2007
- Larsson SC, Wolk A: Overweight, obesity and risk of liver cancer: a meta-analysis of cohort studies. *Br J Cancer* 97:1005–1008, 2007
- Jacobs ET, Thompson PA, Martinez ME: Diet, gender, and colorectal neoplasia. *J Clin Gastroenterol* 41:731–746, 2007
- Dai Z, Xu YC, Niu L: Obesity and colorectal cancer risk: a meta-analysis of cohort studies. *World J Gastroenterol* 13:4199–4206, 2007
- Ferrante AW, Jr.: Obesity-induced inflammation: a metabolic dialogue in the language of inflammation. *J Intern Med* 262:408–414, 2007
- Jackson G: The importance of risk factor reduction in erectile dysfunction. *Curr Urol Rep* 8:463–466, 2007
- Werder O: Battle of the bulge: an analysis of the obesity prevention campaigns in the United States and Germany. *Obes Rev* 8:451–457, 2007
- Kawaji K, Fujita Y, Yajima Y, Shirataka M, Kubo H: Usefulness of anaerobic threshold in estimating intensity of exercise for diabetics. *Diabetes Res Clin Pract* 6:303–309, 1989
- Runge CF: Economic consequences of the obese. *Diabetes* 56:2668–2672, 2007
- Neumark-Sztainer D: Preventing the broad spectrum of weight-related problems: working with parents to help teens achieve a healthy weight and a positive body image. *J Nutr Educ Behav* 37:S133–S140, 2005
- Thorpe KE, Florence CS, Howard DH, Joski P: The impact of obesity on rising medical spending. *Health Aff (Millwood)* W4 (Suppl.):480–486, 2004
- Combination pharmacotherapy for cardiovascular disease. *Ann Intern Med* 143:593–599, 2005
- Franco OH, Karnik K, Bonneux L: The future of metabolic syndrome and cardiovascular disease prevention: polytype or polyhope? Tales from the polyera. *Horm Metab Res* 39:627–631, 2007
- Athyros VG, Tziomalos K, Mikhailidis DP, Pagourelas ED, Kakafika AI, Skaperdas A, Hatzitolios A, Karagiannis A: Do we need a statin-nicotinic acid-aspirin mini-polypill to treat combined hyperlipidaemia? *Expert Opin Pharmacother* 8:2267–2277, 2007
- Jamieson MJ, Naghavi M: Multi-constituent cardiovascular pills (MCCP)—challenges and promises of population-based prophylactic drug therapy for prevention of heart attack. *Curr Pharm Des* 13:1069–1076, 2007
- Rosenthal T, Gavras I: Fixed-drug combinations as first-line treatment for hypertension. *Prog Cardiovasc Dis* 48:416–425, 2006
- Sleight P, Pouleur H, Zannad F: Benefits, challenges, and registerability of the polypill. *Eur Heart J* 27:1651–1656, 2006
- Daviglus ML, Lloyd-Jones DM, Pirzada A: Preventing cardiovascular disease in the 21st century: therapeutic and preventive implications of current evidence. *Am J Cardiovasc Drugs* 6:87–101, 2006
- Karlsson HK, Zierath JR: Insulin signaling and glucose transport in insulin resistant human skeletal muscle. *Cell Biochem Biophys* 48:103–113, 2007
- Chalidakov GN, Stankulov IS, Hristova M, Ghenev PI: Adipobiology of disease: adipokines and adipokine-targeted pharmacology. *Curr Pharm Des* 9:1023–1031, 2003
- Abate N, Garg A: Heterogeneity in adipose tissue metabolism: causes, implications and management of regional adiposity. *Prog Lipid Res* 34:53–70, 1995
- Terstappen GC, Schlupen C, Raggiaschi R, Gavraghi G: Target deconvolution strategies in drug discovery. *Nat Rev Drug Discov* 6:891–903, 2007
- Miller HI, Henderson DR: Governmental influences on drug development: striking a better balance. *Nat Rev Drug Discov* 6:532–539, 2007
- Horsley W: Diabetes. *Experto Crede. Lancet* 5:174–175, 1827
- Dugdill L, Graham RC, McNair F: Exercise referral: the public health panacea for physical activity promotion? A critical perspective of exercise referral schemes; their development and evaluation. *Ergonomics* 48:1390–1410, 2005
- Karacabey K: Effect of regular exercise on health and disease. *Neuro Endocrinol Lett* 26:617–623, 2005
- Hesselink MK, van Baak MA: Physical activity and health, novel concepts and new targets: report from the 12th Conference of the International Research Group on the Biochemistry of Exercise. *Proc Nutr Soc* 63:189–197, 2004
- Stewart KJ: Exercise training and the cardiovascular consequences of type 2 diabetes and hypertension: plausible mechanisms for improving cardiovascular health. *JAMA* 288:1622–1631, 2002
- Blair SN, LaMonte MJ, Nichaman MZ: The evolution of physical activity recommendations: how much is enough? *Am J Clin Nutr* 79:913S–920S, 2004
- Blair SN, Kohl HW 3rd, Barlow CE, Paffenbarger RS Jr, Gibbons LW, Macera CA: Changes in physical fitness and all-cause mortality: a prospective study of healthy and unhealthy men. *JAMA* 273:1093–1098, 1995
- National Center for Chronic Disease Prevention and Health Promotion: *Physical Activity and Health: A Report of the Surgeon General*. Atlanta, Georgia, U.S. Department of Health and Human Services, 1996
- Eriksson J, Taimela S, Koivisto VA: Exercise and the metabolic syndrome. *Diabetologia* 40:125–135, 1997
- Duncan GE: Exercise, fitness, and cardiovascular disease risk in type 2 diabetes and the metabolic syndrome. *Curr Diab Rep* 6:29–35, 2006
- Lavie CJ, Milani RV: Cardiac rehabilitation and exercise training programs in metabolic syndrome and diabetes. *J Cardiopulm Rehabil* 25:59–66, 2005
- Carroll S, Duffield M: What is the relationship between exercise and metabolic abnormalities? A review of the metabolic syndrome. *Sports Med* 34:371–418, 2004
- Miller WC, Koceja DM, Hamilton EJ: A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. *Int J Obes Relat Metab Disord* 21:941–947, 1997
- Donnelly JE, Smith B, Jacobsen DJ, Kirk E, Dubose K, Hyder M, Bailey B, Washburn R: The role of exercise for weight loss and maintenance. *Best Pract Res Clin Gastroenterol* 18:1009–1029, 2004
- Miller WC: How effective are traditional dietary and exercise interventions for weight loss? *Med Sci Sports Exerc* 31:1129–1134, 1999
- Schoeller DA: But how much physical activity? *Am J Clin Nutr* 78:669–670, 2003
- Bamman MM: The Exercise Dose-Response: Key Lessons from the Past (Essay on an APS Classic Paper). *Am J Physiol Endocrinol Metab* 2007
- Cosca DD, Navazio F: Common problems in endurance athletes. *Am Fam Physician* 76:237–244, 2007