



MEETING ABSTRACT

Open Access

Opposite effect of Valproate on Tax and HBZ expression in T-lymphocytes from HTLV-1 asymptomatic carriers and HAM/TSP patients

Gildas Belrose¹, Antoine Gross², Stéphane Olindo³, Agnès Lézin¹, Maryvonne Dueymes¹, Didier Smadja³, Yuetsu Tanaka⁴, Luc Willems⁵, Jean-Michel Mesnard², Jean-Marie Peloponese², Raymond Césaire^{1*}

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses
Leuven and Gembloux, Belgium. 5-8 June 2011

A determinant of HTLV-1 associated myelopathy/tropical spastic paraparesis (HAM/TSP) development is HTLV-1-infected cell burden. Viral proteins Tax and HBZ, encoded by the positive and negative strands of the pX region respectively, play a key role in HTLV-1 persistence. Tax drives CD4⁺ T-cell clonal expansion but is the immunodominant antigen. Valproate (VPA), an histone deacetylase inhibitor, has been proposed to trigger Tax expression and expose latent HTLV-1 reservoir to immune destruction. We evaluated the impact of VPA treatment on Tax, Gag and HBZ expression in cultured lymphocytes from asymptomatic HTLV-1 carriers and HAM/TSP patients. Around one-fifth of provirus-positive CD4⁺ T cells spontaneously became Tax-positive. The estimation rose up to two-thirds of Tax-positive infected cells when VPA was added. VPA enhanced Gag p19 protein release. Tax and Gag mRNA levels spontaneously peaked, before a decline concomitant to HBZ mRNA increase. VPA treatment enhanced and prolonged Tax mRNA expression, while it blocked HBZ expression. This is the first *ex vivo* study on the balance between Tax and HBZ expression (i.e. sense and antisense transcription). Our data suggest that besides modulation of the expression of Tax, another mechanism involving repression of HBZ may determine the outcome of VPA treatment on HTLV-1-infected cell proliferation and survival.

Author details

¹Laboratoire de Virologie-Immunologie, EA 4537, Centre Hospitalier Universitaire de Fort-de-France, Fort-de-France, Martinique. ²CEAPB, CNRS UMR 5236, Université de Montpellier, Montpellier, France. ³Service de Neurologie, EA 4537, Centre Hospitalier Universitaire de Fort-de-France, Fort-de-France, Martinique. ⁴Department of Immunology, Graduate School and Faculty of Medicine, University of the Ryukyus, Okinawa, Japan. ⁵Cellular and Molecular Biology, Agro-Bio Tech, Gembloux, Belgium.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A198

Cite this article as: Belrose *et al.*: Opposite effect of Valproate on Tax and HBZ expression in T-lymphocytes from HTLV-1 asymptomatic carriers and HAM/TSP patients. *Retrovirology* 2011 8(Suppl 1):A198.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: raymond.cesaire@chu-fortdefrance.fr

¹Laboratoire de Virologie-Immunologie, EA 4537, Centre Hospitalier Universitaire de Fort-de-France, Fort-de-France, Martinique
Full list of author information is available at the end of the article