Pharmacognosy Magazine's new and improved Impact Factor – 1.525

Nandan S.

Assoc-Editor, Pharmacogn. Mag., Splendid Plaza, No. 26, Wheeler Road, Cox Town, Bangalore 560 005, India

Submitted: 28-08-2013 Published: 07-09-2013

Thank you for your continuous support for the *Pharmacognosy Magazine*. As a valued reader, we would like to share important news with you. The 2012 Journal Citation Reports[®] from Thomson Reuters is now released and we are delighted to announce that the *Pharmacognosy Magazine* has its new Impact Factor.

We are pleased to announce that our Impact Factor has increased to **1.525**, retaining our position as one of the top pharmaceutical journals in India (2012 Journal Citation Reports[®], ©Thomson Reuters, 2013).^[1]

Few of the achievements this year

- Country-based ranking (India) based on Impact Factor

 6.
- Subject-based ranking (Medicinal Chemistry) based on Impact Factor – 44.
- Immediacy Index 0.135.
- Impact Factor increased from 1.159^[2] (2011 Journal Citation Reports[®], ©Thomson Reuters, 2012) to 1.525 (2012 Journal Citation Reports[®], ©Thomson Reuters, 2013).

Some of the top cited articles in past 2 years

- Total phenolic distribution of juice, peel, and seed extracts of four pomegranate cultivars.^[3]
- Total polyphenolic (flavonoids) content and antioxidant capacity of different Ziziphora clinopodioides Lam. Extracts.^[4]

Access this article online Quick Response Code: Website: www.phcog.com DOI: 10.4103/0973-1296.117807

Address for correspondence:

Nandan S., Assoc-Editor, Pharmacogn. Mag., Splendid Plaza, No. 26, Wheeler Road, Cox Town, Bangalore 560 005, India. E-mail: nandan.s@phcog.net

- Studies on the active components and antioxidant activities of the extracts of Mimosa pudica Linn. from southern China.^[5]
- Title: Chemical composition and antimicrobial activities of the essential oils from three ecotypes of Zataria multiflora.^[6]
- Effects of *Hypericum perforatum* extract on rat irritable bowel syndrome.^[7]
- Isolation and identification of phenolic compounds from *Gynura divaricata* leaves.^[8]
- A HPLC-DAD method for the simultaneous determination of five marker components in the traditional herbal medicine Bangpungtongsung-san.
- Phytochemical constituents and antioxidant activities of the whole leaf extract of Aloe ferox Mill.^[10]
- Screening and analysis of the multiple absorbed bioactive components and metabolites of Baihe Zhimu Tang by the metabolic fingerprinting technique and liquid chromatography/diode array detectionelectrospray ionization-mass spectrometry.^[11]
- Determination of some phenolic compounds in *Crocus* sativus L. corms and its antioxidant activities study. [12]

We are extremely thankful to the authors, reviewers, readers, and Publisher (Medknow – Wolters Kluwer) for their support.

REFERENCES

- 2012 Journal Citation Reports® Science Edition. Thomson Reuters: 2013.
- Mueen Ahmed KK. 2012 Another successful new year for Pharmacogn Mag. Pharmacogn Mag 2012;8:1-3.
- Gözlekçi S, Saraçoglu O, Onursal E, Özgen M. Total phenolic distribution of juice, peel, and seed extracts of four pomegranate cultivars. Pharmacogn Mag 2011;7:161-4.
- Tian S, Shi Y, Zhou X, Ge L, Upur H. Total polyphenolic (flavonoids) content and antioxidant capacity of different Ziziphora clinopodioides Lam. extracts. Pharmacogn Mag 2011:7:65-8.
- Zhang J, Yuan K, Zhou WI, Zhou J, Yang P. Studies on the active components and antioxidant activities of the extracts of Mimosa pudica Linn. from southern China. Pharmacogn Mag 2011;7:35-9.

- Zomorodian K, Saharkhiz MJ, Rahimi MJ, Bandegi A, Shekarkhar G, Bandegani A, et al. Chemical composition and antimicrobial activities of the essential oils from three ecotypes of Zataria multiflora. Pharmacogn Mag 2011;7:53-9.
- Mozaffari S, Esmaily H, Rahimi R, Baeeri M, Sanei Y, Asadi-Shahmirzadi A, et al. Effects of Hypericum perforatum extract on rat irritable bowel syndrome. Pharmacogn Mag 2011;7:213-23.
- Wan C, Yu Y, Zhou S, Tian S, Cao S. Isolation and identification of phenolic compounds from *Gynura divaricata* leaves. Pharmacogn Mag 2011;7:101-8.
- Weon JB, Yang HJ, Ma JY, Ma CJ. A HPLC-DAD method for the simultaneous determination of five marker components in the traditional herbal medicine Bangpungtongsung-san. Pharmacogn Mag 2011;7:60-4.
- 10. Wintola OA, Afolayan AJ. Phytochemical constituents and

- antioxidant activities of the whole leaf extract of *Aloe ferox* Mill. Pharmacogn Mag 2011;7:325-33.
- Qin K, Cai H, Liu X, Lu T, Fang Q, Yao Z, et al. Screening and analysis of the multiple absorbed bioactive components and metabolites of Baihe Zhimu Tang by the metabolic fingerprinting technique and liquid chromatography/diode array detectionelectrospray ionization-mass spectrometry. Pharmacogn Mag 2011;7:177-85.
- Hosseinzadeh H, Shamsaie F, Mehri S. Antioxidant activity of aqueous and ethanolic extracts of *Crocus sativus* L. stigma and its bioactive constituents, crocin and safranal. Pharmacogn Mag 2009;5:419-24.

Cite this article as: Nandan S. Pharmacognosy Magazine's new and improved Impact Factor - 1.525. Phcog Mag 2013;9:283-4.

Source of Support: Nil. Conflict of Interest: None declared.