



DATA NOTE

The complete genome sequence of *Toxicodendron radicans*, Eastern Poison Ivy [version 1; peer review: 2 approved, 1 approved with reservations]

Toby Pirro, Stacy Pirro

Biodiversity, IRIDIAN GENOMES, Bethesda, MD, 20817, USA

V1 First published: 20 Aug 2020, 9:1015
<https://doi.org/10.12688/f1000research.25556.1>
 Latest published: 20 Aug 2020, 9:1015
<https://doi.org/10.12688/f1000research.25556.1>

Abstract

Eastern Poison Ivy (*Toxicodendron radicans*, Anacardiaceae) is well known in Eastern North America for causing contact dermatitis, an itchy and painful rash in most people who come in contact with it. We present the whole genome sequence and annotation of this species. A total of 96,255,779 paired-ends reads consisting of 28.9 G bases were obtained by sequencing one leaf from a wild-collected plant. The reads were assembled by a *de novo* method followed by alignment to related species. Annotation was performed via GenMark-ES. The raw and assembled data is publicly available via GenBank: Sequence Read Archive ([SRR10325927](https://www.ncbi.nlm.nih.gov/sra/SRR10325927)) and Assembly ([GCA_009867345](https://www.ncbi.nlm.nih.gov/assembly/GCA_009867345)).

Keywords

Toxicodendron radicans, Eastern Poison Ivy, genome, assembly, annotation



This article is included in the **Genome Sequencing gateway**.

Open Peer Review

Reviewer Status

	Invited Reviewers		
	1	2	3
version 1 20 Aug 2020	 report	 report	 report
1. Rongxi Sun , Jiangxi Agricultural University, Nanchang, China			
2. Xuming Zhou , Chinese Academy of Sciences, Beijing, China			
3. Hyungtaek Jung , The University of Queensland, St Lucia, Australia			

Any reports and responses or comments on the article can be found at the end of the article.

Corresponding author: Stacy Pirro (info@iridiangenomes.org)

Author roles: **Pirro T:** Resources; **Pirro S:** Data Curation, Funding Acquisition, Investigation, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was supported by IRIDIAN GENOMES (IRGEN-43853).

Copyright: © 2020 Pirro T and Pirro S. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Pirro T and Pirro S. **The complete genome sequence of *Toxicodendron radicans*, Eastern Poison Ivy [version 1; peer review: 2 approved, 1 approved with reservations]** F1000Research 2020, 9:1015
<https://doi.org/10.12688/f1000research.25556.1>

First published: 20 Aug 2020, 9:1015 <https://doi.org/10.12688/f1000research.25556.1>

Introduction

Eastern Poison Ivy (*Toxicodendron radicans*, Anacardiaceae) is well known in the Eastern United States, Canada, Mexico and parts of China for causing contact dermatitis, an itchy and painful rash in most of the human population (Barceloux, 2008). The rash, along with accompanying blisters, is caused by urushiol, an oil compound in the plant's sap. Urushiol-induced allergic rashes are a Type IV hypersensitivity reaction (Kalish & Johnson, 1990). This type of reaction is a cell-mediated response and can take hours to days to produce symptoms (Williams *et al.*, 1999). Approximately 15% of people have no allergic reaction to urushiol, but most people experience a reaction from between 5–12 days after exposure. Allergic reactions can increase in duration and severity after each incident (Bonnekoh *et al.*, 2019).

Exposure to the urushiol in *Toxicodendron radicans* can cause hypersensitivity in related plants, such as mango, *Mangifera indica* (Yoo & Carius, 2019), and the Chinese Lacquer Tree, *Toxicodendron vernicifluum*. Consumer products made from the Chinese Lacquer Tree are manufactured by curing the urushiol-containing sap to a clear, hard, waterproof substance. Improperly cured products can cause contact dermatitis in urushiol-hypersensitive people several years after manufacture, and present an ongoing health challenge to lacquerware workers (Ma *et al.*, 2012).

A complete genome sequence for this species will allow the insight into the evolution of the urushiol biosynthetic pathway.

Methods

A leaf from a single wild-collected *Toxicodendron radicans* plant was used as the source of genomic DNA. Extraction was performed on tissue from a single leaf using the Qiagen DNAeasy genomic extraction kit for plants, using the standard

process. A paired-end sequencing library was constructed using the Illumina TruSeq kit, according to the manufacturer's instructions. The library was sequenced on an Illumina Hi-Seq platform in paired-end, 2 × 150bp format.

The resulting fastq files were trimmed of adapter/primer sequence and low-quality regions with Trimmomatic (v0.33) (Bolger *et al.*, 2014). The trimmed sequence was assembled by SPAdes (v2.5) (Bankevich *et al.*, 2012) followed by a finishing step using RagTag (v1.0.0) (Alonge, 2020) to make additional contig joins based on conserved regions in related plant species: *Mangifera indica* (mango, GCA_011075055) and *Pistacia vera* (pistachio, GCA_008641045). Default procedures were used for all assembly steps.

Annotation was performed using GeneMark-ES (v2.0) (Lomsadze *et al.*, 2005). Annotation was performed fully *de novo* without a curated training set and default parameters.

Results

The genome assembly yielded a total sequence length of 454,874,194 bp over 270,263 scaffolds with an N50 of 1,945,245. The GeneMark-ES annotation resulted in 42,021 genes.

Data availability

Underlying data

Raw and assembled data is publicly available via GenBank:

Raw genome of *Toxicodendron radican*, Accession number SRR10325927: <https://www.ncbi.nlm.nih.gov/sra/?term=SRR10325927>

Assembly of *Toxicodendron radican*, Accession number GCA_009867345: https://www.ncbi.nlm.nih.gov/assembly/GCA_009867345.1/

References

- Alonge M: Ragtag: **Reference-guided genome assembly correction and scaffolding**. GitHub archive. 2020.
[Publisher Full Text](#)
- Bankevich A, Nurk S, Antipov D, *et al.*: SPAdes: **A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing**. *J Comput Biol*. 2012; **19**(5): 455–477.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Barceloux DG: **Medical Toxicology of Natural Substances: Foods, Fungi, Medicinal Herbs, Plants, and Venomous Animals**. John Wiley and Sons. 2008.
[Publisher Full Text](#)
- Bolger AM, Lohse M, Usadel B: **Trimmomatic: A flexible trimmer for Illumina Sequence Data**. *Bioinformatics*. 2014; **30**(15): 2114–20.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bonnekoh H, Poch G, Metz M: **Severe contact dermatitis caused by urushiol in Japanese lacquer**. *Contact Dermatitis*. 2019; **80**(1): 55–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kalish RS, Johnson KL: **Enrichment and function of urushiol (poison ivy)-specific T lymphocytes in lesions of allergic contact dermatitis to urushiol**. *J Immunol*. 1990; **145**(11): 3706–3713.
[PubMed Abstract](#)
- Lomsadze A, Ter-Hovhannisyan V, Chernoff YO, *et al.*: **Gene identification in novel eukaryotic genomes by self-training algorithm**. *Nucleic Acids Res*. 2005; **33**(20): 6494–6506.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ma XM, Lu R, Miyakoshi T: **Recent advances in research on lacquer allergy**. *Allergol Int*. 2012; **61**(1): 45–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Williams JV, Light J, Marks Jr JG: **Individual variations in allergic contact dermatitis from urushiol**. *Arch Dermatol*. 1999; **135**(8): 1002–1003.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Yoo MJ, Carius BM: **Mango Dermatitis After Urushiol Sensitization**. *Clin Pract Cases Emerg Med*. 2019; **3**(4): 361–363.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:   

Version 1

Reviewer Report 07 July 2021

<https://doi.org/10.5256/f1000research.28204.r87603>

© 2021 Jung H. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Hyungtaek Jung

School of Biological Sciences, The University of Queensland, St Lucia, Qld, Australia

The authors have assembled the genome by employing Illumina sequencing technology using the leaf tissue of *T. radicans*. There is no doubt that the data generated in the current study will make a valuable contribution to the scientific community, however, I have several concerns about the analyses and/or the interpretation of the data as it currently stands.

Methods:

The fact is Illumina-based genome assemblies contain many misassemblies. Given the circumstances, why did authors choose Illumina reads over PacBio and Oxford Nanopore? What's the overall sequencing and assembly summary? Any chloroplast DNA contamination? If yes, how did you treat them? Where are the executed scripts? Annotation with GeneMark-ES without any RNA-Seqs?

Results:

How did you evaluate the assembly accuracy (e.g. BUSCO)? Where are the structural (RepeatMasker) and functional annotations? It needs more explanation.

Is the rationale for creating the dataset(s) clearly described?

Partly

Are the protocols appropriate and is the work technically sound?

Partly

Are sufficient details of methods and materials provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics and Bioinformatics

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 24 June 2021

<https://doi.org/10.5256/f1000research.28204.r87604>

© 2021 Zhou X. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Xuming Zhou

Key Laboratory of Animal Ecology and Conservation Biology, Chinese Academy of Sciences, Beijing, China

This study reported the complete genome sequence of *Toxicodendron radicans*, which could help future research of this species. The manuscript is well written and should be considered for indexing prior to two questions:

1. The author should provide more details about the annotation, for example, how *de novo* and homologous approaches were performed.
2. The author should compare their genome quality to their relative species.

Is the rationale for creating the dataset(s) clearly described?

Yes

Are the protocols appropriate and is the work technically sound?

Yes

Are sufficient details of methods and materials provided to allow replication by others?

No

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genomics, Phylogenetics, Aging

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 13 November 2020

<https://doi.org/10.5256/f1000research.28204.r73876>

© 2020 Sun R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Rongxi Sun

Jiangxi Agricultural University, Nanchang, China

I have now thoroughly reviewed the manuscript "The complete genome sequence of *Toxicodendron radicans*, Eastern Poison Ivy". Given urushiol in this plant' sap causing dermatitis, an itchy and painful rash in most of the human population, the new assembly of complete genome sequence of this species helps to understand the evolution of the urushiol biosynthetic pathway. However, there are still some flaws in the manuscripts which the authors may consider to change:

1. Method: What is the source of the leaf used for sequencing, such as tree age and specific location.
2. Results : How accurate is this high-throughput genome sequence? Whether there are indicators related to accuracy in the results.
3. In results, the annotation should be more discussed.

Is the rationale for creating the dataset(s) clearly described?

Partly

Are the protocols appropriate and is the work technically sound?

Partly

Are sufficient details of methods and materials provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Genetic diversity and phylogeography

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research