

Oral presentation

Open Access

Ritonavir inhibits NF-AT activation through modulation of the PI-3 kinase/Akt Ppathway

Shibani Pati¹, Anh Thu Nguyen², William Reid¹, J Scott Foulke Jr¹, Anjana Yadav¹, Fabio Romerio¹, Frank Weichold³ and Marvin Reitz^{*3}

Address: ¹Institute of Human Virology, University of Maryland Biotechnology Institute, Baltimore, Maryland, ²University of Maryland, Baltimore County, USA and ³Morgan State University, Baltimore, Maryland, USA

* Corresponding author

from 2006 International Meeting of The Institute of Human Virology
Baltimore, USA. 17–21 November, 2006

Published: 21 December 2006

Retrovirology 2006, **3**(Suppl 1):S68 doi:10.1186/1742-4690-3-S1-S68

© 2006 Pati et al; licensee BioMed Central Ltd.

The HIV protease inhibitor ritonavir has activities apparently unrelated to its inhibition of the HIV protease, including anti-tumor activity in vivo and in vitro, induction of lipodystrophy in vivo, proteasome inhibition, and inhibition of NFκ. Here we show that ritonavir inhibits activation of NF-AT induced by PMA plus ionomycin and by the human herpes virus-8 chemokine receptor homologue, vGPCR. Inhibition occurred by modulation of the PI-3 kinase/Akt/GSK-3 pathway. Ritonavir treatment led to decreased Akt phosphorylation and a resultant decrease in GSK-3 phosphorylation and failed to inhibit NF-AT in GSK-3β^{-/-} knockout cells. Inhibition of multiple signaling pathways by ritonavir may partly explain its anti-tumor activities as well as other effects of ritonavir that are unrelated to its anti-retroviral activity. Taken together, the data suggest that ritonavir may have intrinsic immunomodulatory activities. This work was supported in part by grant RO1 CA099905-01 from NCI.