

Liver Transplantation in Cases with Acute Liver Failure

¹Mustafa Yalçın, ¹Göksel Bengi, ¹Mesut Akarsu, ²Tarkan Ünek, ²İbrahim Astarcioglu

ABSTRACT

Background: Acute liver failure (ALF) is a rare, life-threatening clinical condition that is characterized by severe hepatocellular necrosis, jaundice, coagulopathy and encephalopathy. The aim of this study was to evaluate patients who underwent liver transplantation at Dokuz Eylül University of Medicine Faculty (DEUMF) due to ALF.

Materials and methods: The patients who underwent liver transplantation at DEUMF due to ALF were evaluated retrospectively.

Results: All of the liver transplantations performed in this study were successful; toxicity was the major cause of ALF in these patients (84%).

Conclusion: Results of this study may not be generalizable to all of Turkey since the patients included in this study were only from one region. However, our study is in accordance with others that show that liver transplantation is a very safe and effective method for the treatment of ALF.

Keywords: Acute liver failure, Liver transplantation, Treatment of ALF

How to cite this article: Yalçın M, Bengi G, Akarsu M, Ünek T, Astarcioglu İ. Liver Transplantation in Cases with Acute Liver Failure. *Euroasian J Hepato-Gastroenterol* 2015;5(2):80-82.

Source of support: Nil

Conflict of interest: None

Copyright and License information: Copyright © 2015; Jaypee Brothers Medical Publishers (P) Ltd. This work is licensed under a Creative Commons Attribution 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/3.0/>

Received Date: 03-03-2015

Accepted Date: 05-05-2015

INTRODUCTION

Acute liver failure (ALF) is a rare, life-threatening clinical condition that is characterized by severe and instant hepatocellular necrosis, jaundice, coagulopathy and encephalopathy. Acute liver failure cases have no history of liver disease and the symptoms of ALF appear within 26 weeks.^{1,2} In developed countries, the major cause of ALF is toxicity; in undeveloped countries, ALF is caused by viral factors.

Acute liver failure is related to high morbidity and mortality. Interactions between the patient, genetics, the cause of the hepatic injury and the liver play a role in the prognosis of ALF.³ Currently, the survival of ALF patients without liver transplantation is unclear. The MELD score, INR and King's College criteria are methods of predicting survival. Recently, there has been limited development in the prognostic scoring system and laboratory tests. The most practical scoring system is the King's College criteria, which predicts mortality caused by paracetamol and other causes.

A meta-analysis consisting of 18 studies and 1105 patients indicated that the sensitivity, specificity and odds ratio of the King's College criteria in ALF caused by nonparacetamol causes were 68 [95% confidence interval (CI) 59–77], 82% (95% CI 75–88) and 12.6 (95% CI 6.5–26.1), respectively.⁴ Today, 45% of adult ALF patients recover spontaneously, 25% undergo liver transplantation, and 30% of the adult patients pass before the liver transplantation due to absolute contraindications. In children with ALF, 56% recover spontaneously, 30% undergo liver transplantation, and 13% pass before the liver transplantation.⁵

The aim of this study is to evaluate the patients who underwent liver transplantation at Dokuz Eylül University of Medicine Faculty (DEUMF) due to ALF.

MATERIALS AND METHODS

The patients who underwent liver transplantation at DEUMF due to ALF were evaluated retrospectively. Sex, age, preoperative biochemical test results, causes of ALF and the conditions of the donors (alive or cadaver) were

¹Department of Gastroenterology, Dokuz Eylül University Faculty of Medicine, İzmir, Turkey

²Department of General Surgery, Dokuz Eylül University Faculty of Medicine, İzmir, Turkey

Address reprint requests to: Göksel Bengi, Associate Professor, Department of Gastroenterology, Özel Çankaya Tıp Merkezi, Gazi Bulvarı No: 56 Çankaya-İzmir, Phone: 0-532-4626972, Fax: 0-232-4451404, e-mail: dr.gokselbengi@hotmail.com

recorded. Data were examined using Microsoft Excel software.

RESULTS

Five of the cases (38.4%) who underwent liver transplantation at DEUMF due to ALF were male and eight (61.6%) were female. The mean age of the subjects was 29 (5–61) years. Three of the donors (23%) were cadaver and 10 of them (77%) were living. The mean values of the patients were as follows: Total Bilirubin: 31 mg/dl, AST: 766U/L, ALT: 787U/L, INR: 3.2, albumin: 3 gr/dl, MELD score: 31.5, encephalopathy stage: 2.5. When the etiology of ALF was examined, it was found that ALF was caused by drugs in six patients, caused by fireworks (due to the content of phosphorus) in two patients, caused by mushrooms in two patients, and caused by herbalin (*herba centaurii*) in one patient. One of the patients underwent liver transplantation due to autoimmune hepatitis and

another for cryptogenic ALF (Tables 1 and 2). When we examine drug induced hepatotoxicity cases, Ornidazole and amoxicillin-clavulanate were found to be responsible for one patient. On the other hand, active substance could not be identified due to multiple drug use. Liver transplantation was successful in the treatment of all patients.

DISCUSSION

Acute liver failure is a rare, life-threatening clinical condition that is characterized by severe and instant hepatocellular necrosis, jaundice, coagulopathy and encephalopathy. Acute liver failure cases have no history of liver disease, and the symptoms of ALF appear within 26 weeks.^{1,2} Toxicity is the major cause of ALF in developed countries, while viral factors are the major cause in developing countries. Kayaalp et al analyzed 308 Turkish patients and found that the most common causes of ALF were hepatitis A in children (20.9%) and hepatitis B in adults (34.7%). Eighteen percent of ALS cases are due to cryptogenic causes, while 14% are due to metabolic causes. Wilson disease is the most common cause of metabolic diseases, while toxic liver failure is most often due to mushroom poisoning (13%). Poisoning from fireworks and phosphorus are regional factors, while anti-tuberculosis agents (3.2% of ALF causes) are the most common cause of drug-based ALF. Paracetamol is responsible for only 0.7% of ALF cases.⁶ In our current study, toxicity was the major cause of ALF in the patients who underwent liver transplantation (84%). None of the patients who underwent liver transplantation had ALF caused by viral factors. We believe that the results of this study do not reflect the patient profile of all of Turkey, but reflects only the profile of the patients who applied to our hospital.

Acute liver failure is related to high morbidity and mortality. Interactions between the patient, genetics,

Table 1: Characteristics of the cases

Cases (N)	Sex	Age (years)	Factor	Donor
N1	F	5	Autoimmune	Alive
N2	F	22	Cryptogenic	Alive
N3	F	47	Toxic	Cadaver
N4	M	20	Toxic	Alive
N5	M	28	Toxic	Alive
N6	F	5	Toxic	Alive
N7	F	8	Toxic	Alive
N8	M	33	Toxic	Alive
N9	M	43	Toxic	Alive
N10	F	36	Toxic	Alive
N11	F	61	Toxic	Cadaver
N12	F	44	Toxic	Cadaver
N13	M	28	Toxic	Alive
Total 13	61% F 39% M	Average age 29.2	11 Toxic 1 Autoimmune 1 Cryptogenic	23% Cadaver 77% Alive

F: Female; M: Male

Table 2: Laboratory test results of cases

Case	AST (u/l)	ALT (u/l)	T. Bil (mg/dl)	Cr. (mg/dl)	Na (meq/l)	Alb (g/dl)	INR	MELD	Encephalopathy grade	Assit
N1	57	73	17.29	0.52	143	5.3	2.933	29	4	None
N2	622	756	20.56	0.5	139	2.3	3.406	32	4	Low
N3	1507	1668	45	0.2	139	3.4	4.42	37	4	Low
N4	625	504	48	1	137	2.8	2.9	33	1	Low
N5	1723	1967	42	0.6	138	2.7	6.05	41	3	None
N6	2210	1188	26.5	0.2	142	3.4	2.3	28	3	Low
N7	1747	1150	3.2	1	132	1.8	2.3	20	3	None
N8	126	197	43.92	0.56	137	3	2.55	31	1	Low
N9	87	1540	19.92	0.81	144	3.2	3.18	31	3	Low
N10	622	659	26.06	0.33	140	2.7	2.84	30	1	None
N11	120	70	39.01	0.63	137	2.8	5	38	2	None
N12	205	134	46.7	0.2	133	2.9	2.18	30	1	Low
N13	313	331	33.8	0.39	149	3.1	2.5	30	3	None
Total 13	766.4	787.4	31.6	0.5	139.2	3.0	3.2	31.5	2.5	

F: Female; M: Male; T. bil: Total bilirubin; Cr: Creatine; Alb: Albumine; MELD: Model for end-stage liver disease

the cause of the hepatic injury play a role in the prognosis of ALF.³ Today, 45% of adult patients recover spontaneously, 25% undergo liver transplantation, and 30% are lost before the liver transplantation due to absolute contraindications. On the other hand, 56% of children with ALF recover spontaneously, 30% undergo liver transplantation, and 13% are lost before the liver transplantation.⁵ In our hospital, 13 cases of acute liver failure were treated via liver transplantation (three from cadaver, 10 from living donor). Liver transplantation is a very successful method for the treatment of ALF.

Results of this study indicate that toxicity is the major cause of ALF in patients who underwent liver transplantation at DEUMF. However, this result is not in accordance with the general situation in Turkey. Our current results only reflect the characteristics of our region. However, liver transplantation was successful in all patients, and there were no complications during or after the surgery. As previous studies have shown, liver

transplantation is a very safe and effective method for the treatment of ALF.

REFERENCES

1. Larson AM. Diagnosis and management of acute liver failure. *Curr Opin Gastroenterol* 2010;26(3):214-221.
2. Bernal W, Auzinger G, Dhawan A, Wendon J. Acute liver failure. *Lancet* 2010;376(9736):190-201.
3. Strnad P, Zhou Q, Hanada S, Lazzeroni LC, Zhong BH, So P, Davern TJ, Lee WM. Acute Liver Failure Study Group, Omary MB. Keratin variants predispose to acute liver failure and adverse outcome: race and ethnic associations. *Gastroenterol* 2010;139(3):828-835.
4. Mc Phail MJ, Wendon JA, Bernal W. Meta-analysis of performance of King's College Hospital Criteria in prediction of outcome in nonparacetamol-induced acute liver failure. *J Hepatol* 2010;53(3):492-499.
5. Lee WM, Squires RH Jr, Nyberg SL, Doo E, Hoofnagle JH. Acute liver failure: summary of a workshop. *Hepatol* 2008; 47(4):1401-1415.
6. Kayaalp C, Ersan V, Yılmaz S. Acute liver failure in Turkey: a systematic review. *Turk J Gastroenterol* 2014; 25(1):35-40.