

The Effect of Multifocal Soft Contact Lens Wear on Axial and Peripheral Eye Elongation in the BLINK Study

Donald O. Mutti,¹ Loraine T. Sinnott,¹ David A. Berntsen,² Lisa A. Jones-Jordan,¹ Danielle J. Orr,¹ and Jeffrey J. Walline¹; for the BLINK Study Group

¹The Ohio State University College of Optometry, Columbus, Ohio, United States

²University of Houston College of Optometry, Houston, Texas, United States

Correspondence: Donald O. Mutti, The Ohio State University College of Optometry, West 10th Avenue, Columbus, OH 43210-1280, USA; mutti.2@osu.edu.

Received: March 31, 2022

Accepted: September 5, 2022

Published: September 28, 2022

Citation: Mutti DO, Sinnott LT, Berntsen DA, Jones-Jordan LA, Orr DJ, Walline JJ, for the BLINK Study Group. The effect of multifocal soft contact lens wear on axial and peripheral eye elongation in the BLINK study. *Invest Ophthalmol Vis Sci.* 2022;63(10):17. <https://doi.org/10.1167/iovs.63.10.17>

PURPOSE. The purpose of this study was to compare axial and peripheral eye elongation during myopia therapy with multifocal soft contact lenses.

METHODS. Participants were 294 children (177 [60.2%] girls) age 7 to 11 years old with between -0.75 diopters (D) and -5.00 D of myopia (spherical component) and less than 1.00 D astigmatism at baseline. Children were randomly assigned to Biofinity soft contact lenses for 3 years: D-designs with a $+2.50$ D addition, $+1.50$ D addition, or single vision. Five measurements of eye length were averaged at the fovea, $\pm 20^\circ$, and $\pm 30^\circ$ in the horizontal and vertical meridians of the right eye using the Haag-Streit Lenstar LS 900.

RESULTS. Axial elongation over 3 years with single vision contact lenses was greater than peripheral elongation in the superior and temporal retinal quadrants by 0.07 mm (95% confidence interval [CI] = 0.05 to 0.09 mm) and 0.06 mm (95% CI = 0.03 to 0.09 mm) and similar in the inferior and nasal quadrants. Axial elongation with $+2.50$ D addition multifocal contact lenses was similar to peripheral elongation in the superior retinal quadrant and less than peripheral elongation in the inferior and nasal quadrants by -0.04 mm (95% CI = -0.06 to -0.01 mm) and -0.06 mm (95% CI = -0.09 to -0.02 mm).

CONCLUSIONS. Wearing $+2.50$ D addition multifocal contact lenses neutralized or reversed the increase in retinal steepness with single vision lenses. The mismatch between greater inhibition of elongation at the fovea than peripherally despite greater peripheral myopic defocus suggests that optical myopia therapy may operate through extensive spatial integration or mechanisms other than local defocus.

Keywords: myopia, refractive error, contact lenses, peripheral refraction

A considerable amount of evidence shows that the growth of the eye and the development of refractive error can be modified by changes in visual experience. Lid occlusion and form deprivation in animal models, primarily chicks and monkeys, result in highly myopic refractive errors in a short amount of time.^{1,2} These animal species are also responsive to the sign and magnitude of defocus created by lenses, compensating for the imposed refractive error by means of accelerated or inhibited ocular elongation.³⁻⁵ Animal models have been useful in explaining some aspects of human refractive error development. Disruption of normal, high contrast vision in human infants results in myopic refractive errors.^{6,7} Human infants are also responsive to their own native hyperopic refractive errors, with modulation of growth that results in near-emmetropia from an axial length well matched to ocular refractive power.⁸⁻¹⁰

A key feature of the visual control of eye growth is local control, that ocular responses to changes in the visual environment are spatially specific. The portion of the visual field exposed to form deprivation or defocus in animal studies corresponds to the portion of the eye that

exhibits the compensating change in growth.¹¹⁻¹³ Separating the eye from the central nervous system through optic nerve section does not prevent myopic responses to form deprivation or compensation for refractive errors imposed by minus lenses.^{12,14} Whereas the central and peripheral retina are both sensitive to changes in the visual environment,¹¹ responses to peripheral manipulations are not always confined to the periphery. The ability of the retinal periphery to alter growth at the fovea has been well documented in foveal ablation experiments in the monkey.^{15,16} Foveal elongation accelerates when the retinal periphery is preferentially exposed to form deprivation or blur using apertures,^{15,17} but may not be affected if the exposure is too far into the periphery relative to the central retina.¹⁸ Because these studies have only used animal models, the extent of spatial integration of a visual signal by the retinal periphery and its ability to influence foveal growth in children have yet to be determined.

There is good evidence that the peripheral visual environment in children may affect growth at the fovea. Children wearing progressive addition spectacle lenses experience

less myopia progression.^{19–21} These effects were assumed to be due to reduced hyperopic defocus at the fovea during near vision. However, a report from the Study of Theories about Myopia Progression project suggested that the beneficial effect of progressive addition lenses might be due to superior retinal myopic defocus from the inferiorly placed addition power.^{21,22} Myopia control through center-distance multifocal contact lenses or overnight orthokeratology presents this additional plus power 360° around the periphery. The optical effect of these multifocal contact lenses is to reduce the peripheral hyperopia that is present when children wear a traditional single vision correction, such as glasses or contact lenses. Multifocal contact lens clinical trial results suggest that the influence of the periphery outweighs the local foveal visual signal; despite a clear foveal image and good visual acuity, altering the peripheral optical profile results in slower myopia progression and less axial elongation in children.^{23,24} A theory of growth based on local control would predict that peripheral expansion might be inhibited more than axial elongation during optical myopia control. Different local effects might also be expected between ocular meridians. Baseline results without contact lenses showed that the Bifocal Lenses In Near-sighted Kids (BLINK) study participants had about +1.8 D of relative peripheral hyperopia at 40° in both the nasal and temporal visual fields. Interestingly, the superior and inferior visual fields at 30° showed the opposite sign of defocus, about –0.5 D of relative peripheral myopia.²⁵ Local control would predict that inhibition of peripheral ocular expansion would be greater in the vertical compared to the horizontal meridian due to greater peripheral myopia. The BLINK study cross-sectional results argue against this meridional local control. Both meridians showed steeper retinas at higher levels of myopia, but the difference between meridians remained similar throughout the range of refractive errors.²⁵

The purpose of the current report is to investigate whether the effect of wearing center-distance multifocal soft contact lenses on axial and peripheral elongation of the eye exhibits more local versus global effects compared to single-vision soft contact lenses. Baseline data have been reported previously.²⁵ The primary analyses in this report address the pattern of inhibition from +2.50 D addition contact lenses compared to +1.50 D addition and single vision contact lenses as a function of retinal meridian, quadrant, and eccentricity using longitudinal data collected over 3 years. Local control of ocular growth would predict that elongation in the vertical meridian would be less than in the horizontal meridian and that peripheral elongation would be inhibited more than axial elongation when wearing +2.50 D addition contact lenses. Similar levels of elongation in both meridians and similar (or greater) inhibition of elongation at the fovea than the periphery would indicate a more global response to wearing multifocal contact lenses.

METHODS

Detailed methods used in the BLINK study, and those used for peripheral biometry and refractive error measurements, have been published previously.^{25,26} To summarize, the BLINK study enrolled 294 children between the ages of 7 and 11 years old (inclusive) with between –0.75 D and –5.00 D of myopia in the most hyperopic meridian, less than 1.00 D of astigmatism, and 2.00 D or less of anisometropia into a 3-year randomized clinical trial to determine if center-

distance multifocal contact lenses slowed the progression of myopia more than single vision contact lenses. Children were randomized in equal numbers (98 per group) to wear one of three Biofinity soft contact lens designs (CooperVision; Pleasanton, CA): single-vision, multifocal D with +1.50 D addition power, or multifocal D with +2.50 D addition power. The research adhered to the tenets of the Declaration of Helsinki, was reviewed and approved by independent ethical review boards at the University of Houston and The Ohio State University, conformed with the principles and applicable guidelines for the protection of human subjects in biomedical research, and was monitored by an independent data and safety monitoring committee. Assent from children and parental permission were obtained from each participant and participant's parent/guardian, respectively. The registration for this clinical trial can be found at ClinicalTrials.gov (Identifier: NCT02255474).

All central and peripheral measurements of eye length and refractive error for this report were made on the right eye only under cycloplegia. Subjects received one drop of either 0.5% tetracaine or 0.5% proparacaine followed by 2 single drops of 1% tropicamide 5 minutes apart in each eye. Central and peripheral eye length were the average of 5 valid measurements (unflagged by the instrument) at each retinal location using the Lenstar LS 900 optical biometer (Haag-Streit USA, Mason, OH). Axial length was measured along the line of sight, then peripheral eye length was measured by having the subject turn the eye to fixate small targets on the face of the instrument at eccentricities of 20° and 30°, both horizontally in nasal and temporal gaze and vertically in superior and inferior gaze. Central and peripheral refractive error were measured with and without contact lens correction using the open-view Grand Seiko WAM-5500 binocular autorefractor/keratometer (AIT Industries, Bensenville, IL). The central value was the average of 10 valid readings, whereas the peripheral values were the average of 5 valid readings (within ±1.0 D of the median value for sphere and cylinder). Central refractive error was measured along the line of sight, then horizontal peripheral refractive error was measured in nasal and temporal gaze at eccentricities of 20°, 30°, and 40°. Vertical peripheral refractive error was measured in superior and inferior gaze at 20° and 30°. Measurements of peripheral refractive error with the contact lenses in place were only made in the horizontal meridian because contact lens decentration could be avoided by having participants turn their head laterally during measurement instead of their eyes; this was not possible for vertical measurements. Peripheral refraction in the vertical meridian with contact lenses in place was estimated under an assumption of rotational symmetry. The differences in peripheral measurements between eye-only and the eye wearing a lens at each horizontal eccentricity were added to the corresponding eye-only values at each vertical eccentricity.

Statistical analyses were completed using SAS, version 9.4 for Windows (SAS Institute, Cary, NC). The modeled 3-year changes in elongation at 20° and 30° were analyzed by treatment group and within each of the 4 quadrants. The modeled 3-year change in axial elongation was compared to peripheral elongation, again by treatment group and within each of the 4 quadrants. In order to avoid the many comparisons generated by multiple eccentricities, and to create analyses similar to those for baseline data, individual participant peripheral eye length data were also fit by quadratic equations as a function of gaze angle, one horizontal and one vertical, for each study year.²⁵ This approach has been

validated against magnetic resonance imaging (MRI) and schematic eye retinal contours.²⁷⁻²⁹ Models for each of these analytic approaches included treatment group, study year (categorical variable), and their interactions adjusted for sex, age group, and study site. The *P* values for eccentricity and quadrant comparisons were adjusted for multiple comparisons using the step-down Bonferroni method of Holm.³⁰

RESULTS

As previously reported for the BLINK study, approximately 60% of the participants were girls, the average age was 10.3 ± 1.2 years, 60% were 10 or 11 years old at baseline, 26% were Hispanic or Latino, and 68% were White.²⁴ The average baseline cycloplegic refractive error was -2.39 ± 1.00 D with only small amounts of astigmatism (average ± standard deviation: $J_0 = +0.06 \pm 0.19$ D; $J_{45} = +0.05 \pm 0.14$ D). Of the 294 enrolled subjects, 287 (97.6%) completed the 3-year visit. The multifocal addition power resulted in a different peripheral refractive error profile at the baseline visit compared to wearing single vision contact lenses. The +2.50 D addition power decreased peripheral hyperopia in the horizontal meridian seen with single vision contact lenses, resulting in small amounts of peripheral myopia across ±30° but not at 40° for either the nasal or the temporal retina. The +2.50 D addition power increased the estimated amount of peripheral myopia in the vertical meridian compared to single vision contact lenses. Peripheral refractive error for the +1.50 D addition power was intermediate between the +2.50 D addition power and single vision contact lenses (Fig. 1).

Randomization resulted in balanced peripheral eye lengths between treatment groups with no significant differences at baseline at any corresponding quadrant or eccentricity (All *P* > 0.37). Eyes elongated each year at every measured eccentricity in both meridians and in all three treatment groups. The pattern of elongation kept the peripheral eye length profile generally parallel to the profile at baseline (Fig. 2). Changes in foveal and peripheral eye length during each study year are depicted in Supplementary Figure

S1. This general symmetry in elongation can be seen in the similar amounts of change across eccentricities between each study year and in each meridian. Figure 2 and Supplementary Figure S1 also illustrate the inhibition in elongation produced by wearing +2.50 D addition contact lenses compared to +1.50 D addition and single vision contact lenses. The pattern of inhibition across ±30° of the retina also appears to be generally symmetric, with a parallel but lower amounts of change across the retina, particularly between baseline and year 1, for the compressed pattern for eye lengths in both meridians (Figs. 2A, 2B) and lower amounts of change across the retina, particularly between baseline and year 1 (Supplementary Figs. S1A, S1B), for the +2.50 D addition multifocal contact lenses than +1.50 D addition and single vision contact lenses (Supplementary Figs. S1A, S1B).

The results for elongation at the fovea of the right eye only in this report are similar to those previously reported for both eyes: children randomized to wearing +2.50 D addition contact lenses had less axial elongation over 3 years (0.39 mm, 95% confidence interval [CI] = 0.33 to 0.46 mm) than those wearing either +1.50 D addition (0.55 mm, 95% CI = 0.47 to 0.63 mm) or single vision contact lenses (0.63 mm, 95% CI = 0.57 to 0.70 mm; *P* < 0.001 for comparisons between +2.50 D addition and either +1.50 D addition or single vision control).²⁴ The +2.50 D addition had its greatest effect in year 1 when eye growth was most rapid (*P* < 0.001) and continued to significantly inhibit elongation in year 2 (*P* < 0.001). However, there was no significant difference in the rate of axial elongation between treatment groups in year 3 (*P* = 0.59).

The cumulative 3-year changes in eye length again showed generally symmetric elongation with overlapping 95% CIs across the retinal eccentricities within a quadrant and treatment group (Fig. 3). These similar rates of elongation occurred despite baseline horizontal and vertical asymmetries. Eye length in the temporal retinal quadrant was already relatively shorter at baseline by 0.40 mm at 30° than the nasal retinal quadrant (Fig. 2A) and eye length in the inferior retinal quadrant was 0.15 mm




FIGURE 1. Peripheral refractive error with contact lenses in place at the baseline visit (single vision, +1.50 D addition power, and +2.50 D addition power) in (A) the horizontal meridian, and (B) estimated values for the vertical meridian. Error bars represent the standard error of the mean (some obscured; vertical standard errors estimated from vertical peripheral refraction).




FIGURE 2. Eye lengths measured across $\pm 30^\circ$ of the (A) horizontal meridian and (B) vertical meridian in each year of the study. Error bars represent standard errors of the mean.

shorter at $+30^\circ$ than the superior retinal quadrant at -30° (Fig. 2B).²⁵ Although elongation was generally symmetric within a treatment group, statistically significant interactions between treatment group and both quadrant (interaction $P = 0.005$) and eccentricity (interaction $P = 0.0002$) indicated that treatment resulted in small, but important, departures from this symmetry. For children wearing single vision contact lenses, elongation was greater at 20° than at 30° in 3 out of 4 quadrants by an average of 0.05 mm (95% CI = 0.03 to 0.08 mm; $P < 0.001$; Figs. 3A, 3C, 3D). Only the temporal quadrant had similar amounts of elongation at 20° and 30° (Fig. 3B). Axial elongation was greater than peripheral elongation in the superior and temporal quadrants (axial-superior = 0.07 mm, [95% CI = 0.05 to 0.09 mm; $P < 0.001$]; axial-temporal = 0.06 mm [95% CI = 0.03 to 0.09 mm; $P = 0.002$]; Figs. 3B, 3C). Axial and peripheral elongation were similar in the inferior and nasal quadrants (Figs. 3A, 3D). This pattern with single vision contact lenses was similar in children wearing +1.50 D addition contact lenses. There were no significant differences in elongation at either 20° or 30° in either meridian between children wearing +1.50 D addi-

tion power and those wearing single vision contact lenses. Axial elongation was again greater than peripheral elongation in the superior and temporal quadrants and similar to each other in the inferior and nasal quadrants (axial-superior = 0.04 mm [95% CI = 0.02 to 0.06 mm; $P = 0.008$]; axial-temporal = 0.08 mm [95% CI = 0.05 to 0.11 mm; $P < 0.001$]).

Compared to wearing single vision lenses, children wearing +2.50 D addition multifocal contact lenses showed significantly less elongation at all corresponding peripheral points by 0.16 to 0.28 mm ($P < 0.001$ to 0.049) with the exception of 30° in the superior retina (Fig. 3C). The largest effects of treatment with +2.50 D multifocal contact lenses occurred at 20° and the fovea. The significant differences in elongation between 20° and 30° with single vision lenses were not present with +2.50 D addition multifocal contact lenses. Axial elongation was similar to peripheral elongation in the superior quadrant and less than peripheral elongation in the inferior and nasal quadrants, neutralizing or reversing the pattern seen with single vision contact lenses (axial-inferior = -0.04 mm [95% CI = -0.06 to -0.01 mm; $P = 0.051$]; axial-nasal = -0.06 mm [95% CI = -0.09 to -0.02 mm; $P = 0.005$; Figs. 3A, 3C, 3D). Only the temporal quad-




FIGURE 3. Three-year changes in eye length by eccentricity and retinal/visual field quadrant. Error bars represent the 95% confidence interval.

rant showed greater axial than peripheral elongation with both single vision (as above) and +2.50 D addition contact lenses (axial-temporal = 0.06 mm [95% CI = 0.03 to 0.09 mm; $P < 0.001$]; Fig. 3B).

The results based on quadratic fits to the peripheral eye length profiles of individual participants were consistent with those above by eccentricity and quadrant. A negative quadratic coefficient in Figure 4A indicates a downward-turned parabola, with more negative values associated with steeper retinas (i.e. shorter eye length with greater eccentricity). As previously reported, coefficients at baseline were more negative in the horizontal meridian than in the vertical meridian.²⁵ The quadratic coefficients for eye length profile for children wearing single vision contact lenses became more negative over the 3 years of the study in both the horizontal ($P < 0.003$) and the vertical meridians ($P < 0.001$; Fig. 4A), consistent with greater axial than peripheral elongation noted above. Axial elongation at the fovea when wearing single vision contact lenses was 0.63 mm, greater than the quadratic model estimates of peripheral elongation at 30° in the superior (0.54 mm) and temporal (0.56 mm) retinal quadrants and similar to foveal elongation in the nasal

and inferior retina (0.61 mm; Fig. 5). In contrast to the pattern seen with single vision contact lenses, the quadratic coefficients for the +2.50 D addition group either underwent no significant change over 3 years in the horizontal meridian ($P = 0.42$) or became less negative in the vertical meridian ($P = 0.006$). Results for the +1.50 D addition group were intermediate, with no significant change in the quadratic coefficients in either meridian (all $P > 0.06$). Inhibition of elongation with +2.50 D addition contact lenses was most often greatest at the fovea. The treatment effect (estimated by the difference between the green and the blue curves in Fig. 5) was 0.24 mm at the fovea, but only 0.12 mm, 0.14 mm, and 0.16 mm at 30° in the superior, nasal, and inferior quadrants, respectively. The effect of treatment was only similar to that at the fovea at 30° in the temporal retina (0.22 mm; Fig. 5A). These results suggest that wearing +2.50 D addition multifocal contact lenses neutralized the pattern of increasing retinal steepness seen with single vision contact lenses in the horizontal meridian and reversed it in the vertical meridian toward becoming flatter.

The asymmetry in eye length profile is described by the linear term of the quadratic fit. The linear coeffi-




FIGURE 4. (A) Average quadratic coefficients fit to eye length data across $\pm 30^\circ$ of the retina in the horizontal and vertical meridian by treatment group and study year. Error bars represent 95% confidence intervals. (B) Average linear coefficients from the quadratic equations fit to eye length data across $\pm 30^\circ$ of the retina in the horizontal and vertical meridian by treatment group and study year. Error bars represent 95% confidence intervals.

coefficients were consistent with the nasal-temporal and superior-inferior asymmetries noted above at baseline (Fig. 4B). The negative value in each meridian indicates a shorter eye length in the temporal and inferior retinal quadrants, respectively. The linear terms showed significant changes in the negative direction over the 3 years of the study in the horizontal meridian in all 3 treatment groups ($P < 0.008$) and in the positive direction in the vertical meridian for children wearing single vision ($P < 0.001$) and +1.50 D addition contact lenses ($P = 0.005$). The linear coefficient for the vertical meridian showed no significant change over 3 years in children wearing the +2.50 D addition multifocal contact lenses (Fig. 4B; $P = 0.14$).

DISCUSSION

The eyes of children wearing single vision contact lenses for 3 years elongated more at the fovea than in the periphery, resulting in a steeper retina than at baseline. The eyes of children wearing +2.50 D addition center-distance contact lenses showed either similar or less elongation at the fovea compared to the periphery in 3 of the 4 measured quadrants. As a result, the retinal steepening seen with single vision contact lenses was either not observed in the horizontal meridian or reversed toward a flatter retina than at baseline in the vertical meridian while wearing +2.50 D addition contact lenses. This result is unlikely to be the simple product of less myopia progression and less axial elongation in the +2.50 D addition group. Scaling the coefficients in Figure 4A according to the relative treatment effects between groups would result in negative changes in the coefficients over 3 years in both meridians instead of a statistically insignificant positive change in the horizontal meridian and a statistically significant positive change in the vertical meridian.

There are few longitudinal studies of peripheral eye length to compare to the BLINK study results. A study of 140 mostly emmetropic children also found elongation in all 4 retinal quadrants at 20° after 30 months of follow-

up.³¹ Greater foveal than peripheral elongation found in the current study aligns well with a cross-sectional result from MRI scans showing that axial length exceeds eye width and height, and by greater amounts at higher amounts of myopia.³² The report most relevant to the BLINK study is from a recent study of 58 myopic Chinese children 8 to 12 years of age who self-selected to be fit with either overnight orthokeratology or single vision spectacles for 1 year.³³ The eye elongated at the fovea and at all peripheral points in the single vision spectacle group, as expected. Elongation in the orthokeratology group was unexpectedly greater for the nasal retina than with single vision spectacles, and by a substantial amount in one year of up to 0.21 mm.³³ The BLINK study also showed greater inhibition of elongation in the temporal compared to the nasal retina, but there was evidence of inhibition of elongation at the most peripheral points including nasally, even after 1 year (Figs. 2, 3). Peripheral elongation was never greater in children wearing +2.50 D addition contact lenses than with single vision lenses. Both orthokeratology and the center-distance multifocal contact lenses used in the BLINK study (Fig. 1) expose the peripheral retina to myopic defocus,^{34,35} making it difficult to pinpoint the reason for the difference between the results for the two studies.

The peripheral optical profiles of single vision and +2.50 D addition contact lenses were asymmetric by both quadrant and eccentricity (Fig. 1), providing an opportunity to evaluate global versus local responses to defocus. Single vision contact lenses were associated with large and nearly equal amounts of peripheral hyperopia in both nasal and temporal horizontal quadrants, whereas the vertical meridian was consistently emmetropic. Therefore, local control of eye growth by defocus would not explain the nasal-temporal asymmetry in elongation nor the generally symmetric pattern of elongation between the horizontal and vertical meridians with single vision lenses. The +2.50 D addition reduced peripheral hyperopia in the horizontal meridian, particularly at 20° over the temporal retina, and produced a substantial amount of peripheral myopia




FIGURE 5. The model quadratic fits to central and peripheral eye lengths across $\pm 30^\circ$ at baseline and at 3 years for +2.50 D addition and single vision contact lenses in (A) the horizontal meridian and (B) the vertical meridian. The numbers given represent differences in eye length between adjacent curves centrally and peripherally at 20° and 30° (indicated by the arrows). Results for children wearing +1.50 D addition contact lenses are omitted for clarity.

in the vertical meridian (Fig. 1). This altered profile was associated with a positive treatment benefit of reduced elongation across the $\pm 30^\circ$ of the retina in all quadrants. However, local defocus would not explain the lower degree of inhibition of elongation peripherally in three of the four quadrants compared to the fovea, the portion of the retina with the least change in its image. Local defocus would also not explain the similar amounts of inhibition between the horizontal and vertical ocular meridians. Local control of ocular growth would be more consistent with less elongation in the vertical than the horizontal meridian and greater inhibition of peripheral than axial elongation. Given that these two outcomes did not occur in the BLINK study, the eye seems to exhibit a more global than local response to the optical profile created by the +2.50 D multifocal contact lenses.

Local defocus cues might still be the effective signal but perhaps are being integrated over an extensive amount of the peripheral retina. Another possibility to explore is that similar growth between meridians may be the result of reduced sensitivity in the vertical meridian to inhibition by peripheral myopia because of its habitual exposure to that sign of defocus.^{25,36} The global response seems more

similar to an emmetropic pattern of elongation, namely more symmetric change between fovea and periphery as opposed to the greater axial than the peripheral elongation seen during myopic progression with standard single vision corrections. Uniformity of change between the fovea and periphery as measured by eye length or peripheral refraction is seen in children likely to remain emmetropic.^{31,37,38} This pattern may also be characteristic of a positive treatment effect during myopia control. The 2-year results for children wearing Defocus Incorporated Multiple Segments (DIMS) lenses showed a more uniform pattern of change in peripheral refraction than single vision lenses across the horizontal $\pm 30^\circ$ of the retina.^{39,40} The recent study of peripheral eye lengths following orthokeratology also found that the pattern of growth was more symmetric across the retina than with single vision glasses.³⁵ It should be recognized that this slower growth during myopia control is still much faster than would be expected in 7 to 11 year old emmetropic children. Axial elongation over 3 years from the age of 10 to 13 years might be 0.20 to 0.22 mm in children without myopia compared to 0.39 mm seen in the BLINK study participants wearing +2.50 D addition contact lenses.^{41,42} Current approaches to myopia control may change the pattern and

rate of eye elongation, but more successful myopia control should have even greater levels of inhibition as a goal. Greater levels of inhibition across the retina would also be worthwhile. The differences in elongation between +2.50 D addition contact lenses and single vision contact lenses were less peripherally than at the fovea by as much as 50%, or 0.12 mm in the superior retina compared to the 0.24 mm of treatment effect seen at the fovea. This result suggests the need to evaluate elongation at locations more peripheral than 30°. The $\pm 30^\circ$ retinal eccentricity corresponds to a sagittal depth of about 2 to 3 mm within a vitreous chamber that might be 17 mm deep in a young, myopic eye. Current optical biometric techniques for measuring eye length cannot assess more extreme eccentricities, leaving most of the retinal periphery unmeasured. Inhibition of elongation should extend throughout the entire retinal periphery if myopia control truly results in a smaller globe than an untreated eye. A large part of the motivation for initiating myopia control is because the increasing prevalence of myopia is expected to increase ocular morbidity from excessive eye enlargement.^{43–45} The axial benefit of myopia control loses considerable value if it does not result in an overall smaller eye.

CONCLUSIONS

Wearing multifocal contact lenses with a +2.50 D addition reduced that rate of elongation at every peripheral point in a generally global, symmetric pattern. However, the presence of minor deviations from completely symmetric inhibition of elongation has important implications. Foveal and peripheral eye elongation did not follow the pattern predicted by local defocus cues. Less elongation in the vertical meridian than the horizontal and less elongation peripherally than axially were not observed. The degree of inhibition of elongation was similar between the horizontal and vertical meridians and greater at the fovea than in the periphery when wearing +2.50 D center-distance multifocal contact lenses. If local control does not explain the more global response of the eye to multifocal contact lenses, future research should attempt to better characterize the optical signals and ocular mechanisms that are responsible for the treatment benefit produced by this form of myopia control. Assessments of elongation beyond 30° during myopia control are also needed.

Acknowledgments

Supported by the National Institutes of Health (NIH) grants NIH U10 EY023204; NIH U10 EY023206; NIH U10 EY023208; NIH U10 EY023210; NIH P30 EY007551; and NIH UL1 TR002733.

BLINK Study Group Executive Committee

Jeffrey J. Walline (Study Chair, The Ohio State University College of Optometry), David A. Berntsen (UH Clinic Principal Investigator, University of Houston College of Optometry), Donald O. Mutti (OSU Clinic Principal Investigator, The Ohio State University College of Optometry), Lisa A. Jones-Jordan (Data Coordinating Center Director, The Ohio State University College of Optometry), Donald F. Everett (NEI Program Official), Jimmy Le (NEI Program Official, 2019-present).

Chair's Center

Kimberly J. Shaw (Study Coordinator, The Ohio State University College of Optometry), Juan Huang (Investigator, The Ohio State University College of Optometry), Bradley E. Dougherty

(Survey Consultant, The Ohio State University College of Optometry).

Data Coordinating Center

Loraine T. Sinnott (Biostatistician, The Ohio State University College of Optometry).

University of Houston Clinic Site (University of Houston College of Optometry).

Laura Cardenas (Clinic Coordinator), Krystal L. Schulle (Unmasked Examiner, 2014–2019), Dashaini V. Retnasothie (Unmasked Examiner, 2014–2015), Amber Gaume Giannoni (Masked Examiner), Anita Tićak (Masked Examiner), Maria K. Walker (Masked Examiner), Moriah A. Chandler (Unmasked Examiner, 2016–present), Mylan Nguyen (Data Entry, 2016–2017), Lea Hair (Data Entry, 2017–2018), Augustine N. Nti (Data Entry, 2019–2020).

Ohio State University Clinic Site (The Ohio State University College of Optometry)

Jill A. Myers (Clinic Coordinator), Alex D. Nixon (Unmasked Examiner, 2014–2019), Katherine M. Bickle (Unmasked Examiner, 2014–2020), Gilbert E. Pierce (Unmasked Examiner, 2014–2019), Kathleen S. Reuter (Masked Examiner, 2014–2019), Dustin J. Gardner (Masked Examiner, 2014–2016), Andrew D. Pucker (Masked Examiner, 2015–2016), Matthew Kowalski (Masked Examiner, 2016–2017), Ann Morrison (Masked Examiner, 2017–2019), Danielle J. Orr (Unmasked Examiner, 2018–present).

Data Safety and Monitoring Committee

Janet T. Holbrook (Chair, Johns Hopkins Bloomberg School of Public Health), Jane Gwiazda (Member, New England College of Optometry), Timothy B. Edrington (Member, Southern California College of Optometry), John Mark Jackson (Member, Southern College of Optometry), Charlotte E. Joslin (Member, University of Illinois at Chicago).

Disclosure: **D.O. Mutti**, Bausch & Lomb (F), Vyluma (C); **L.T. Sinnott**, Bausch & Lomb (F); **D.A. Berntsen**, Bausch & Lomb (F), Visioneering Technologies (advisory board) (C, R); **L.A. Jones-Jordan**, Bausch & Lomb (F); **D.J. Orr**, None; **J.J. Walline**, Bausch & Lomb (F)

References

- Wallman J, Turkel J, Trachtman J. Extreme myopia produced by modest changes in early visual experience. *Science*. 1978;201:1249–1251.
- Wiesel TN, Raviola E. Myopia and eye enlargement after neonatal lid fusion in monkeys. *Nature*. 1977;266:66–68.
- Irving EL, Sivak JG, Callender MG. Refractive plasticity of the developing chick eye. *Ophthalmic Physiol Opt*. 1992;12:448–456.
- Schaeffel F, Glasser A, Howland HC. Accommodation, refractive error, and eye growth in chickens. *Vision Res*. 1988;28:639–657.
- Smith EL, 3rd, Hung LF. The role of optical defocus in regulating refractive development in infant monkeys. *Vision Res*. 1999;39:1415–1435.
- Gee SS, Tabbara KF. Increase in ocular axial length in patients with corneal opacification. *Ophthalmology*. 1988;95:1276–1278.
- Miller-Meeks MJ, Bennett SR, Keech RV, Blodi CF. Myopia induced by vitreous hemorrhage. *Am J Ophthalmol*. 1990;109:199–203.
- Ehrlich DL, Braddick OJ, Atkinson J, et al. Infant emmetropization: Longitudinal changes in refraction components from nine to twenty months of age. *Optom Vis Sci*. 1997;74:822–843.
- Pennie FC, Wood IC, Olsen C, White S, Charman WN. A longitudinal study of the biometric and refractive changes in full-term infants during the first year of life. *Vision Res*. 2001;41:2799–2810.

10. Mutti DO, Mitchell GL, Jones LA, et al. Axial growth and changes in lenticular and corneal power during emmetropization in infants. *Invest Ophthalmol Vis Sci.* 2005;46:3074–3080.
11. Wallman J, Gottlieb MD, Rajaram V, Fugate-Wentzek LA. Local retinal regions control local eye growth and myopia. *Science.* 1987;237:73–77.
12. Troilo D, Wallman J. The regulation of eye growth and refractive state: An experimental study of emmetropization. *Vision Res.* 1991;31:1237–1250.
13. Smith EL, Huang J, Hung LF, et al. Hemiretinal form deprivation: Evidence for local control of eye growth and refractive development in infant monkeys. *Invest Ophthalmol Vis Sci.* 2009;50:5057–5069.
14. Wildsoet C. Neural pathways subserving negative lens-induced emmetropization in chicks—insights from selective lesions of the optic nerve and ciliary nerve. *Curr Eye Res.* 2003;27:371–385.
15. Smith EL, Kee CS, Ramamirtham R, Qiao-Grider Y, Hung LF. Peripheral vision can influence eye growth and refractive development in infant monkeys. *Invest Ophthalmol Vis Sci.* 2005;46:3965–3972.
16. Smith EL, Ramamirtham R, Qiao-Grider Y, et al. Effects of foveal ablation on emmetropization and form-deprivation myopia. *Invest Ophthalmol Vis Sci.* 2007;48:3914–3922.
17. Smith EL, Hung LF, Huang J. Relative peripheral hyperopic defocus alters central refractive development in infant monkeys. *Vision Res.* 2009;49:2386–2392.
18. Schippert R, Schaeffel F. Peripheral defocus does not necessarily affect central refractive development. *Vision Res.* 2006;46:3935–3940.
19. Gwiazda J, Hyman L, Hussein M, et al. A randomized clinical trial of progressive addition lenses versus single vision lenses on the progression of myopia in children. *Invest Ophthalmol Vis Sci.* 2003;44:1492–1500.
20. Correction of Myopia Evaluation Trial 2 Study Group for the Pediatric Eye Disease Investigator Group. Progressive addition lenses versus single vision lenses for slowing progression of myopia in children with high accommodative lag and near esophoria. *Invest Ophthalmol Vis Sci.* 2011;52:2749–2757.
21. Berntsen DA, Sinnott LT, Mutti DO, Zadnik K. A randomized trial using progressive addition lenses to evaluate theories of myopia progression in children with a high lag of accommodation. *Invest Ophthalmol Vis Sci.* 2012;53:640–649.
22. Berntsen DA, Barr CD, Mutti DO, Zadnik K. Peripheral defocus and myopia progression in myopic children randomly assigned to wear single vision and progressive addition lenses. *Invest Ophthalmol Vis Sci.* 2013;54:5761–5770.
23. Chamberlain P, Peixoto-de-Matos SC, Logan NS, et al. A 3-year randomized clinical trial of MiSight lenses for myopia control. *Optom Vis Sci.* 2019;96:556–567.
24. Walline JJ, Walker MK, Mutti DO, et al. Effect of high add power, medium add power, or single-vision contact lenses on myopia progression in children: The BLINK randomized clinical trial. *JAMA.* 2020;324:571–580.
25. Mutti DO, Sinnott LT, Reuter KS, et al. Peripheral refraction and eye lengths in myopic children in the bifocal lenses in nearsighted kids (BLINK) study. *Transl Vis Sci Technol.* 2019;8:17.
26. Walline JJ, Gaume Giannoni A, Sinnott LT, et al. A randomized trial of soft multifocal contact lenses for myopia control: Baseline data and methods. *Optom Vis Sci.* 2017;94:856–866.
27. Verkicharla PK, Suheimat M, Schmid KL, Atchison DA. Peripheral refraction, peripheral eye length, and retinal shape in myopia. *Optom Vis Sci.* 2016;93:1072–1078.
28. Verkicharla PK, Suheimat M, Pope JM, et al. Validation of a partial coherence interferometry method for estimating retinal shape. *Biomed Opt Express.* 2015;6:3235–3247.
29. Atchison DA, Charman WN. Can partial coherence interferometry be used to determine retinal shape? *Optom Vis Sci.* 2011;88:E601–E607.
30. Holm S. A simple sequentially rejective multiple test procedure. *Scand J Statist.* 1979;6:65–70.
31. Schmid GF. Association between retinal steepness and central myopic shift in children. *Optom Vis Sci.* 2011;88:684–690.
32. Atchison DA, Jones CE, Schmid KL, et al. Eye shape in emmetropia and myopia. *Invest Ophthalmol Vis Sci.* 2004;45:3380–3386.
33. Huang Y, Li X, Ding C, et al. Orthokeratology reshapes eyes to be less prolate and more symmetric. *Cont Lens Anterior Eye.* 2022;45(4):101532.
34. Kang P, Swarbrick H. Peripheral refraction in myopic children wearing orthokeratology and gas-permeable lenses. *Optom Vis Sci.* 2011;88:476–482.
35. Queiros A, Gonzalez-Meijome JM, Jorge J, Villa-Collar C, Gutierrez AR. Peripheral refraction in myopic patients after orthokeratology. *Optom Vis Sci.* 2010;87:323–329.
36. Atchison DA, Pritchard N, Schmid KL. Peripheral refraction along the horizontal and vertical visual fields in myopia. *Vision Res.* 2006;46:1450–1458.
37. Mutti DO, Hayes JR, Mitchell GL, et al. Refractive error, axial length, and relative peripheral refractive error before and after the onset of myopia. *Invest Ophthalmol Vis Sci.* 2007;48:2510–2519.
38. Lee TT, Cho P. Relative peripheral refraction in children: Twelve-month changes in eyes with different ametropias. *Ophthalmic Physiol Opt.* 2013;33:283–293.
39. Zhang HY, Lam CSY, Tang WC, Leung M, To CH. Defocus incorporated multiple segments spectacle lenses changed the relative peripheral refraction: A 2-year randomized clinical trial. *Invest Ophthalmol Vis Sci.* 2020;61:53.
40. Lam CSY, Tang WC, Tse DY, et al. Defocus incorporated multiple segments (DIMS) spectacle lenses slow myopia progression: A 2-year randomised clinical trial. *Br J Ophthalmol.* 2020;104:363–368.
41. Jones LA, Mitchell GL, Mutti DO, et al. Comparison of ocular component growth curves among refractive error groups in children. *Invest Ophthalmol Vis Sci.* 2005;46:2317–2327.
42. Rozema J, Dankert S, Iribarren R, Lanca C, Saw SM. Axial growth and lens power loss at myopia onset in Singaporean children. *Invest Ophthalmol Vis Sci.* 2019;60:3091–3099.
43. Flitcroft DI. The complex interactions of retinal, optical and environmental factors in myopia aetiology. *Prog Retin Eye Res.* 2012;31:622–660.
44. Saw SM, Gazzard G, Shih-Yen EC, Chua WH. Myopia and associated pathological complications. *Ophthalmic Physiol Opt.* 2005;25:381–391.
45. Bullimore MA, Brennan NA. Myopia control: Why each diopter matters. *Optom Vis Sci.* 2019;96:463–465.