

POSTER PRESENTATION

Open Access

Right Ventricular Volumes and Function normalized to body surface area, age and sex in a large cohort of well-treated Thalassemia Major without myocardial iron overload

Antonella Meloni^{1*}, Maria Chiara Dell'Amico¹, Brunella Favilli¹, Giovanni D Aquaro¹, Ait-Ali Lamia¹, Elisabetta Chiodi², Claudio Ascoti³, Stefania Renne³, Pier Paolo Bitti⁴, Calogera Gerardi⁵, Vincenzo Positano¹, Massimo Lombardi¹, Alessia Pepe¹

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Introduction

Cardiovascular Magnetic Resonance CMR has provided the opportunity to quantify right ventricular (RV) parameters with excellent reproducibility and accuracy. The role of the RV is gaining ground in thalassemia major (TM) patients and this population could experience different "normal" RV values due to chronic anemia and eventually pre-existing iron burdens. In literature, there are no data on the ranges for "normal" RV volumes and ejection fraction (EF) in TM patients. The aim of this study was to establish the ranges for normal RV volumes, mass and ejection fraction normalized to the influence of body surface area (BSA), age and sex from CMR in a large cohort of well-treated TM patients without myocardial iron overload.

Materials

Among the 923 TM patients enrolled in the Myocardial Iron Overload (MIOT) network who underwent CMR for the assessment of cardiac iron overload, function and fibrosis, we selected 142 patients with no known risk factors or history of cardiac disease, normal electrocardiogram, no myocardial iron overload (all the cardiac segments with a normal T2* value) and no myocardial fibrosis. All patients had been regularly transfused and chelated since early childhood. Moreover, we studied 71 healthy subjects matched for age and sex. RV function parameters were quantitatively evaluated in a standard

way by SSFP cine images using MASS[®] software. RV end-diastolic volume (EDV), end-systolic volume (ESV) and stroke volume (SV) were normalized by body surface area (EDVI, ESVI, SVI).

Results

Table 1 shows the comparison of the CMR parameters with differentiation for sex and age in TM patients and healthy subjects and the cut-off of normality defined as mean - 2 standard deviation (SD). TM patients showed significantly lower BSA than the controls ($P < 0.0001$). TM males (except age group 14-20 yrs) showed significantly higher RV EF compared to controls. In TM patients all LV volumes indexes were significantly larger in males than in females ($P < 0.0001$ in all age groups). The EF was not different between the sexes. In males as well as in females the RV volumes were no significant different among the age groups, while in males the EF was significant different ($P = 0.004$).

Conclusion

In a large cohort of well-treated TM patients males showed significantly higher RV EF compared to controls. Appropriate "normal" reference ranges normalized to BSA, sex and age should be used to avoid misdiagnosis of cardiomyopathy in the clinical arena in TM patients.

*G. Monasterio¹ Foundation and Institute of Clinical Physiology, CNR, Pisa, Italy
Full list of author information is available at the end of the article

Table 1

	< 14			14-20			20-30			30-40			>=40		
	TM	H	P	TM	H	P	TM	H	P	TM	H	P	TM	H	P
Males	N=7	N=7		N=6	N=6		N=25	N=15		N=23	N=11		N=6	N=5	
EDVI (ml/m ²)	88 ± 16 (56)	80 ± 12	0.309	89 ± 16 (57)	89 ± 17	0.946	96 ± 21 (54)	103 ± 17	0.245	91 ± 16 (59)	82 ± 13	0.139	90 ± 13 (64)	81 ± 9	0.228
ESVI (ml/m ²)	26 ± 3 (20)	31 ± 8	0.273	36 ± 7 (22)	35 ± 9	0.931	34 ± 8 (18)	44 ± 12	0.0004	33 ± 8 (17)	33 ± 8	0.952	28 ± 6 (16)	35 ± 3	0.045
SVI (ml/m ²)	61 ± 16 (29)	49 ± 6	0.108	53 ± 11 (31)	54 ± 10	0.836	61 ± 18 (25)	60 ± 9	0.882	58 ± 11 (36)	46 ± 17	0.010	62 ± 8 (46)	46 ± 8	0.007
EF (%)	68 ± 6 (56)	62 ± 5	0.048	59 ± 5 (49)	61 ± 5	0.579	64 ± 4 (56)	58 ± 6	0.001	64 ± 5 (54)	61 ± 7	0.189	69 ± 3 (63)	57 ± 5	<0.0001
Females	N=2	N=2	N=2	N=8	N=6		N=24	N=6		N=33	N=8		N=8	N=5	
EDVI (ml/m ²)	53 ± 16 (21)	61 ± 1	0.587	77 ± 8 (61)	79 ± 9	0.866	78 ± 14 (50)	78 ± 12	0.990	73 ± 13 (47)	77 ± 9	0.427	74 ± 11 (52)	70 ± 16	0.648
ESVI (ml/m ²)	18 ± 1 (17)	20 ± 4	0.656	29 ± 5	30 ± 5	0.777	28 ± 8 (12)	31 ± 9	0.378	24 ± 7	26 ± 6	0.317	26 ± 5 (16)	25 ± 12	0.922
SVI (ml/m ²)	35 ± 5 (25)	41 ± 3	0.635	47 ± 4 (39)	49 ± 7	0.673	51 ± 9 (33)	47 ± 7	0.384	49 ± 8 (30)	51 ± 10	0.738	47 ± 6 (35)	45 ± 5	0.517
EF (%)	64 ± 9 (46)	68 ± 6	0.687	62 ± 3 (56)	62 ± 6	0.997	65 ± 6 (53)	61 ± 7	0.171	68 ± 5 (58)	66 ± 8	0.329	67 ± 7 (55)	66 ± 8	0.862

Author details

¹G. Monasterio¹ Foundation and Institute of Clinical Physiology, CNR, Pisa, Italy.
²Arcispedale "S. Anna", Ferrara, Italy. ³P.O. "Giovanni Paolo II", Lamezia Terme, Italy. ⁴P. O. San Francesco. ASL Nuoro, Nuoro, Italy. ⁵Ospedale "Sant'Elia", Caltanissetta, Italy.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P299

Cite this article as: Meloni *et al.*: Right Ventricular Volumes and Function normalized to body surface area, age and sex in a large cohort of well-treated Thalassemia Major without myocardial iron overload. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P299.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

