



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Acute retrograde type A intramural hematoma during severe acute respiratory syndrome coronavirus 2 pandemic

Maria Katsarou, MD,^a Viviana Grassi, MD,^a Chiara Lomazzi, MD,^a Maurizio Domanin, MD,^{a,b} and Santi Trimarchi, MD, PhD,^{a,b} Milan, Italy


Acute intramural hematomas (IMHs) occur in ~6% of patients with acute dissections (ADs) and mostly affect the descending aorta.¹ Type A IMHs involve the ascending aorta and type B IMHs do not involve the ascending aorta. Retrograde type A IMHs (retro-TAIMHs) originate in the descending aorta and extend into the arch or ascending aorta. TAIMHs with distal AD carry an in-hospital mortality risk of 12% to 26%.^{1,2}

We report the case of an 85-year-old woman with acute retro-TAIMH and distal AD. The patient provided consent for the report of her case. She was admitted to the emergency room with acute onset dyspnea and chest pain but no evidence of malperfusion. Emergency computed tomography angiography identified a retro-TAIMH with AD and a proximal entry tear above the celiac axis (A/Cover).

The patient was hemodynamically stable. She was treated with hypotensive and analgesic therapy and hospitalized for intensive monitoring. Follow-up computed tomography angiography was performed at 24 hours (B) and 7 days (C) showing progressive to complete thrombosis of the entry tear, with a reduction in the aortic diameter, the most important predictor of IMH regression and positive outcomes.³ Complete symptom regression occurred. The event was observed during the peak of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic in Lombardy, Italy. The patient had tested positive for SARS-CoV-2 at 5 days after symptom onset, with progressive dyspnea and worsening findings on a chest radiograph (D). She died of pulmonary complications at 19 days postoperatively.

Hybrid treatment with ascending aortic replacement and distal thoracic aortic endovascular repair (TEVAR) or using the frozen elephant trunk procedure is the most appropriate treatment for acute retro-TAIMH. TEVAR is a valid alternative only for patients with prohibitive surgical risk, although the landing zones could be unsuitable and the risk of neurologic and cardiac complications can be high.⁴ Medical treatment appears to be appropriate for asymptomatic patients, those with noncomplicated retro-TAIMHs, and patients with high open surgical and TEVAR risks.⁴

Considering both the absence of end-organ malperfusion and the advanced age of our patient, we chose medical treatment, which can reduce mortality by 67% to 95%.⁵ This choice proved effective with symptom recovery and clinical stability of the present patient, until the deadly overlap of SARS-CoV-2.



REFERENCES

1. Harris KM, Braverman AC, Eagle KA, Woznicki EM, Reed EP, Myrmet T, et al. Acute aortic intramural hematoma: an analysis from the International Registry of Acute Aortic Dissection. *Circulation* 2012;126:S91-6.

From the Section of Vascular Surgery, Fondazione IRCCS Ca'Granda Ospedale Maggiore Policlinico^a; and the Department of Clinical and Community Sciences, University of Milan.^b

Author conflict of interest: none.

E-mail: santi.trimarchi@unimi.it.

The editors and reviewers of this article have no relevant financial relationships to disclose per the JVS policy that requires reviewers to decline review of any manuscript for which they may have a conflict of interest.

J Vasc Surg 2021;73:2166-7

0741-5214

Copyright © 2020 by the Society for Vascular Surgery. Published by Elsevier Inc.

<https://doi.org/10.1016/j.jvs.2020.09.019>

2. Tolenaar JL, Harris KM, Upchurch GR Jr, Evangelista A, Moll FL, Di Eusnio M, et al. The differences and similarities between intramural hematoma of the descending aorta and acute type B dissection. *J Vasc Surg* 2013;58:1498-504.
3. Evangelista A, Dominguez R, Sebastia C, Salas A, Permanyer-Miralda G, Avegliano G, et al. Prognostic value of clinical and morphologic findings in short-term evolution of aortic intramural haematoma: therapeutic implications. *Eur Heart J* 2004;25:81-7.
4. Nauta F, De Beaufort H, Mussa FF, De Vincentiis C, Omura A, Matsuda H, et al. Management of retrograde type A IMH with acute arch tear/type B dissection. *Ann Cardiothorac Surg* 2019;8:531-9.
5. Ince H, Nienaber CA. Diagnosis and management of patients with aortic dissection. *Heart* 2007;93:266-70.

Submitted Jul 12, 2020; accepted Sep 11, 2020.