

Atraumatic acute forearm compartment syndrome due to systemic heparin

Gustavo Chavez,¹ Jeff Choi ,² Nathaniel Fogel,³ Joshua D Jaramillo,² Matthew Murphy,⁴ David Spain²

¹School of Medicine, Stanford University, Stanford, California, USA

²Department of Surgery, Division of General Surgery, Stanford University, Stanford, California, USA

³Department of Orthopedic Surgery, Stanford University, Stanford, California, USA

⁴Department of Surgery, Division of Plastic and Reconstructive Surgery, Stanford University, Stanford, California, USA

Correspondence to

Dr Jeff Choi; jc2226@stanford.edu

Received 13 October 2019

Accepted 24 October 2019

CASE PRESENTATION

A 65-year-old woman was admitted to the hospital for acute heart failure exacerbation. Her history was notable for insulin-dependent diabetes mellitus and mitral valve regurgitation requiring mechanical mitral valve replacement. Found to have mitral paravalvular leak, the patient's home warfarin was held and heparin infusion was started in preparation for valvular surgery.

Eleven days after starting heparin infusion, the patient awoke with severe right forearm pain that worsened throughout the day and by the afternoon had developed decreased sensation at the small finger. Visually, the right forearm had mild swelling and ecchymoses, but was without evidence of traumatic injury, venipuncture, erythema, or fluctuance. On forearm examination, the patient was remarkably tender to palpation over both dorsal and volar compartments, endorsed pain with passive extension of the fingers, and both dorsal and volar compartments were rigid to palpation. The distal compartments of the hand remained soft and non-tender without evidence of increased compartment pressure or ischemia. Vascular examination demonstrated palpable radial and ulnar pulses and a capillary refill of less than 2 seconds.

WHAT WOULD YOU DO?

- Provide analgesics and continue to monitor.
- Obtain an ultrasound of the right forearm over the region of tenderness.
- Obtain a CT angiography of the right upper extremity.
- Obtain forearm compartment pressures.
- Proceed to the operating room for fasciotomy.

WHAT DID WE DO AND WHY?

E. Proceed to the operating room for fasciotomy

After rapid hand surgery consultation, the patient was brought to the operating room to undergo urgent fasciotomy due to concern for atraumatic acute compartment syndrome (ACS). A regional block was pursued secondary to her cardiovascular disease and the related risk associated with a general anesthetic. Once complete, a curvilinear incision was made over the volar forearm, in addition to a limited dorsal incision at the junction between the mobile wad and dorsal compartments. Through these incisions, all compartments were completely released. The volar musculature was noted to be congested, and a bleeding arterial branch within the flexor digitorum superficialis muscle belly was identified and ligated. At time of discharge 2 weeks

after surgery, the patient had a clean and dry incision with preserved motor-sensory function with the exception of persistent decreased sensation of the small finger distal to the palmar digital crease.

Forearm ACS occurs when increase in forearm compartment pressures impede venous and lymphatic drainage, devolving into ischemia and irreversible damage to compartment tissues.^{1,2} Of the forearm compartments, the muscles of the deep volar compartment (flexor pollicis longus and flexor digitorum profundus muscles) and the median nerve are most frequently affected. The most common causes of forearm ACS include supracondylar humerus fractures, distal radial fractures, and both bone forearm fractures.³ Though classic ACS symptoms include pain with passive stretch of muscles, paresthesia, pulselessness, pallor, and paralysis, clinical suspicion should provoke intervention before presentation of the onset of sensory and vascular deficits.²

Atraumatic ACS is a challenging diagnosis as the instigating factors are not well known. They have been associated with hematologic conditions such as hemophilia, disseminated intravascular coagulation, deep vein thrombosis, atypical localized infections, and medications including thrombolytics and anticoagulants.⁴⁻¹⁵ The purported mechanism is spontaneous hemorrhage that leads to increased compartment pressures. There have been four reported cases of heparin-induced or low molecular weight heparin-induced atraumatic ACS of the thigh, and two reported cases of warfarin-induced atraumatic ACS of the upper arm and forearm.^{8,12-15}

This case emphasizes the possibility of atraumatic etiologies for extremity ACS. Patients with known risk factors and symptoms of ACS, even in the absence of trauma, should undergo immediate evaluation and necessary consultation for operative management.

Acknowledgements Dr. Choi would like to recognize the Neil and Claudia Doerhoff fund for support of his scholarly activities.

Contributors GC conducted literature review, drafting, and incorporation of revisions by other authors. JC provided drafting, organizing, revisions, and analysis. NF and JDJ provided revisions and analysis. DS and MM provided revisions and guidance.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Chavez G, Choi J, Fogel N, et al. *Trauma Surg Acute Care Open* 2019;4:e000399.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Jeff Choi <http://orcid.org/0000-0003-1639-8781>

REFERENCES

- 1 Matsen III FA. Compartmental syndromes: an unified concept. *Clin Orthop Relat Res* 1975;113:8–14.
- 2 Friedrich JB, Shin AY. Management of forearm compartment syndrome. *Hand Clin* 2007;23:245–54.
- 3 Kalyani BS, Fisher BE, Roberts CS, Giannoudis PV. Compartment syndrome of the forearm: a systematic review. *J Hand Surg Am* 2011;36:535–43.
- 4 Luce EA, Futrell JW, Wilgus EF, Hoopes JE. Compression neuropathy following brachial arterial puncture in anticoagulated patients. *J Trauma* 1976;16:717–21.
- 5 Macon WL, Futrell JW. Median-nerve neuropathy after percutaneous puncture of the brachial artery in patients receiving anticoagulants. *N Engl J Med* 1973;288:1396.
- 6 Fadel FS, Benyahia M. Compartment syndrome in upper arm in anticoagulant therapy after minor trauma. *Ugeskr Laeger* 2010;172:2149–50.
- 7 Hay SM, Allen MJ, Barnes MR. Acute compartment syndromes resulting from anticoagulant treatment. *BMJ* 1992;305:1474–5.
- 8 Zimmerman DC, Kapoor T, Elfond M, Scott P, et al. Spontaneous compartment syndrome of the upper arm in a patient receiving anticoagulation therapy. *J Emerg Med* 2013;44:e53–6.
- 9 Nadeem RD, Clift BA, Martindale JP, Hadden WA, Ritchie IK, et al. Acute compartment syndrome of the thigh after joint replacement with anticoagulation. *J Bone Joint Surg Br* 1998;80-B:866–8.
- 10 Halpern AA, Mochizuki R, Long CE. Compartment syndrome of the forearm following radial-artery puncture in a patient treated with anticoagulants. *J Bone Joint Surg Am* 1978;60:1136–7.
- 11 Burghardt RD, Siebenlist S, Döbele S, Lucke M, Stöckle U, et al. Compartment syndrome of the thigh. A case report with delayed onset after stable pelvic ring fracture and chronic anticoagulation therapy. *BMC Geriatr* 2010;10:2–6.
- 12 Kakkar R, Ellis M, Fearon PV. Compartment syndrome of the thigh as a complication of anticoagulant therapy in a patient with a left ventricular assist device (Berlin heart). *Gen Thorac Cardiovasc Surg* 2010;58:477–9.
- 13 Limberg RM, Dougherty C, Mallon WK. Enoxaparin-induced bleeding resulting in compartment syndrome of the thigh: a case report. *J Emerg Med* 2011;41:e1–4.
- 14 McLaughlin JA, Paulson MM, Rosenthal RE. Delayed onset of anterior tibial compartment syndrome in a patient receiving low-molecular-weight heparin. A case report. *J Bone Joint Surg Am* 1998;80:1789–90.
- 15 Griffiths D, Jones DHA. Spontaneous compartment syndrome in a patient on long-term anticoagulation. *J Hand Surg Am* 1993;18:41–2.