

RESEARCH ARTICLE

Which Medication Is the Patient Taking at Admission to the Emergency Ward? Still Unclear Despite the Swedish Prescribed Drug Register

Ida Engqvist¹, Katja Wyss¹, Charlotte Asker-Hagelberg^{2,3}, Ulf Bergman^{4,5}, Ingegerd Odar-Cederlöf^{4*}, Carl-Olav Stiller², Jessica Fryckstedt^{1*}

1 Karolinska Institutet, Department of Medicine, Department of Emergency Medicine, Karolinska University Hospital Solna, SE 171 76, Stockholm, Sweden, **2** Karolinska Institutet, Department of Medicine, Clinical Pharmacology Unit, Karolinska University Hospital, Stockholm, Solna, SE 171 76, Stockholm, Sweden, **3** Medical Products Agency, P.O. box 26, SE 751 03, Uppsala, Sweden, **4** Karolinska Institutet, Division of Clinical Pharmacology, Department of Laboratory Medicine, Karolinska University Hospital, Huddinge, SE 141 86, Stockholm, Sweden, **5** Karolinska Institutet, Centre for Pharmacoepidemiology, Department of Medicine, Karolinska University Hospital Solna, SE 171 76, Stockholm, Sweden

* ingegerd.odar-cederlof@karolinska.se (IOC); jessica.fryckstedt@karolinska.se (JF)



OPEN ACCESS

Citation: Engqvist I, Wyss K, Asker-Hagelberg C, Bergman U, Odar-Cederlöf I, Stiller C-O, et al. (2015) Which Medication Is the Patient Taking at Admission to the Emergency Ward? Still Unclear Despite the Swedish Prescribed Drug Register. PLoS ONE 10(6): e0128716. doi:10.1371/journal.pone.0128716

Academic Editor: Alfonso Carvajal, Universidad de Valladolid, SPAIN

Received: October 1, 2014

Accepted: April 29, 2015

Published: June 12, 2015

Copyright: © 2015 Engqvist et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This project received a grant (800 000 SEK about 85 000 EURO) from the National Board of Health and Welfare, administrated by the Stockholm county council. It was mainly used for salaries (for IE, KW and JF). The remaining authors contributed within their ordinary positions in the county.

Competing Interests: CAH is employed by the Swedish Medical Products Agency with academic affiliation to Karolinska Institutet. This publication is a

Abstract

Introduction

Correct information on patients' medication is crucial for diagnosis and treatment in the Emergency Department. The aim of this study was to investigate the concordance between the admission chart and two other records of the patient's medication.

Methods

This cohort study includes data on 168 patients over 18 years admitted to the Emergency Ward between September 1 and 30, 2008. The record kept by the general practitioner and the patient record of dispensed drugs in the Swedish Prescribed Drug Register were compared to the admission chart record.

Results

Drug record discrepancies of potential clinical significance between the admission chart record and the Swedish Prescribed Drug Register or general practitioner record were present in 79 and 82 percent, respectively. For 63 percent of the studied patients the admission chart record did not include all drugs registered in the Swedish Prescribed Drug Register. For 62 percent the admission chart record did not include all drugs registered in the general practitioner record. In addition, for 32 percent of the patients the admission chart record included drugs not registered in the Swedish Prescribed Drug Register and for 52 percent the admission chart record included drugs not found in the general practitioner record. The most discordant drug classes were cardiovascular and CNS-active drugs. Clinically

result of academic research and does not necessarily reflect the view of the Medical Products Agency. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials. None of the other authors state any conflict of financial or non-financial interest.

significant drug record discrepancies were more frequent in older patients with multiple medication and caregivers.

Conclusion

The apparent absence of an accurate record of the patient's drugs at admission to the Emergency Ward constitutes a potential patient safety hazard. The available sources in Sweden, containing information on the drugs a particular patient is taking, do not seem to be up to date. These results highlight the importance of an accurate list of currently used drugs that follows the patient and can be accessed upon acute admission to the hospital.

Introduction

Many physicians in the Emergency Department have experienced the difficulty in obtaining a correct medication history. Incomplete knowledge of the patients' medication may result in incorrect diagnosis and inadequate treatment. In addition, adverse drug reactions, unwanted drug interactions, abrupt discontinuation of medication and other drug related problems (DRPs) may not be identified [1–6]. Physicians can access information about prescribed and dispensed drugs via the Swedish Prescribed Drug Register (SPDR) [7–8]. However, the utility of the SPDR in an Emergency practice has not been evaluated before. Most patients have a general practitioner (GP), who keeps a medication record (GP record—GPr). Upon admission, the emergency physician registers medications in an admission chart record after interviewing the patient, and/or care-takers without prior consultation of the SPDR or the patients' GP. Our previous studies have addressed the therapeutic problems generated by insufficient knowledge on the currently used drugs [2–6]. The aim of this study was not to analyse the clinical consequences of drug record discrepancies in our patient cohort, but rather to quantify major discrepancies of potential clinical importance between different sources of information: the admission chart record, the GP record and drugs in the SPDR.

Methods

The Karolinska University Hospital Solna is one of seven emergency hospitals in Stockholm (2 million inhabitants), Sweden. The Emergency Department has 80 000 visits and 2700 admissions to the Emergency Ward (EW) annually.

Upon admission to the EW, the physician records the patient's current medication in the electronic patient file [9]. This information is derived from previous hospitalizations, the patient directly, relatives or other caregivers. A commonly used source of information is a medication record acquired from the patient's GP. Thirdly, as mentioned, via the Swedish Prescribed Drug Register (SPDR) data on all dispensed drugs, but not over-the-counter (OTC) drugs (unless they were prescribed), can now be made available to prescribing physicians after informed consent of the patient [7,8].

Patients and hospital setting

The study was conducted between September 1 and 30, 2008. Patients considered for inclusion were above 18 years, admitted to the EW during office hours on weekdays. Weekend and night admissions were reviewed on the first subsequent workday if the patients were still present in the ward. Patients signed the consent form after receiving verbal and written information of

the study. The following data were recorded: age, sex, medications on admission, cause and length of hospitalization and caregivers outside of the hospital. Patients who declared no medication were also included since we hypothesised that this information may not be correct.

The admission chart record (ACR) was collected from drugs listed in the admission chart (107 patients) or printed from the electronic health record (61 patients) linked to the electronic patient file [9]. Data from the SPDR was acquired for a period of minimum six months prior to admission since most drugs are prescribed on a three or six month basis in Sweden (monthly elsewhere). Patients named their GP, who was informed of the study purpose and requested to provide a record of the patients' current medication. The GPs were reminded twice. The patients were asked if they consulted several doctors for medications, thus assessing the number of care-givers each patient had.

Comparison between information on the patients' medication

Two investigators (IE and KW) collaborated in comparing the generic substances by name in the three records. No comparison was made between the GP record and the SPDR. Dietary supplements, food agents, non-prescription local treatment e.g. tear substitutes and moisturisers, were excluded. A drug that was used according to one record, but was not listed in the other record was classified as a discrepancy.

Criteria for assessment of the potential clinical importance

A discrepancy between records was judged as of potentially clinical significant by the following criteria:

- The drug was used to treat a condition that needs continuous treatment (for example hypertension).
- The patient could potentially develop a drug related problem when receiving the wrong dose/regimen of the drug (for example analgesics).
- The patient could develop symptoms on abrupt withdrawal of the drug (for example benzodiazepines or beta-receptor blocking agents).

The drug record discrepancies that did not meet these criteria were analysed by specialists in Internal Medicine and Clinical Pharmacology (JF and CAH).

The drug record discrepancies identified by comparison of the admission chart record and the SPDR were classified according to a system previously described in the pilot study [6].

- A_{admission chart-/SPDR+} a drug not included in the admission chart record, but dispensed at pharmacy according to the SPDR.
- B_{admission chart +/-SPDR-} a drug in the admission chart record but not dispensed at pharmacy according to the SPDR.
- C a drug found in both records with significantly different dose or dosing regimen.
- D a drug found in both records but the prescribed amount according to the SPDR is insufficient to last until the admission date or up to 14 days before admission date.

The drug list discrepancies that could arise in the comparison between the admission chart record and the GP record were classified in a similar manner:

- A_{admission chart-/GP+} a drug not included in the admission chart record, but registered in the GP record.

- B admission chart+/GP- a drug in the admission chart record but not included in the GP record.
- C a drug found in both records with significantly different dose or dosing regimen.

Drugs were grouped according to WHO's Anatomical Therapeutic Chemical (ATC) classification system [10].

Statistical Analysis

Graph Pad Prism version 5.02, San Diego, CA was used for statistical analysis. Contingency tables were analysed by Fishers exact test. For pair wise comparison Mann-Whitney U-test was used. Statistical significance was set at $p < 0.05$.

Ethical considerations

This study was approved by the Regional Ethics Review Board (Regionala Etikprövningsnämnden 2008/3:6) at Karolinska Institutet.

Results

Patients

Of the 247 patients admitted to the EW during the study period, 47 were admitted and discharged outside office hours, five declined participation, 25 were unable to give consent and two were foreign citizens. Thus, 168 patients were included. Cardiovascular diseases and dyspnoea accounted for half of the admissions (Table 1a). The median age was 68 years (range 19–93), sex distribution was even (male 83 and female 85), median number of drugs on admission was six (range 0–29) and the median length of hospital stay was three days (range 1–49).

When comparing the admission chart record and the SPDR, 675 drug record discrepancies were identified, 533 (79%) classified as clinically significant. At least one drug record discrepancy was noted in 141 patients (84%), and 133 patients (79%) had at least one clinically significant drug record discrepancy. The mean number of clinically significant drug record discrepancies per patient were 3.2 (median 2.5, range 0–13) (Table 2).

Comparison of admission chart record with the GP record

According to 131 patients, a GP prescribed their medications. Twelve GPs did not provide a medication record, leaving 119 patients available for evaluation.

Of the 626 identified drug record discrepancies, 534 (85%) were classified as of potential clinical significance. In 106 patients (89%), at least one discrepancy was noted, of which 82% (98 patients) were judged to be clinically significant rendering an average of four clinically significant drug record discrepancies per patient (median 3, range 0–23) (Table 2).

A drug present in the admission chart record but not in the GP record (Type B admission chart +/GP- discrepancies) was most common and accounted for 50% of the clinically significant drug record discrepancies, affecting 62 (52%) patients. A drug being present in the GP record but not in the admission chart record (Type A admission chart-/GP+), accounted for 40% of the clinically significant drug record discrepancies, affecting 74 (62%) patients. A significant difference in dose or dose regime (Type C discrepancies) made up 10% of the clinically significant drug record discrepancies, affecting 35 (30%) patients. (Table 3)

Patients at risk

Patients with at least one clinically significant drug record discrepancy of the admission chart record and the SPDR or the GP record were older ($p < 0.05$, Mann Whitney test) and

Table 1. a. Reasons for admission to the Emergency Ward and b. clinically relevant discrepancies.

a		b.		
Symptoms/diagnosis	No. Patients (%)	Other relevant diagnoses	No. Clinically relevant discrepancies ACR/SDPR (%)	No. Clinically relevant discrepancies ACR/GPr (%)
Bleeding	8 (4.7%)	Cardiovasc 12, HT:19, HF:5, AF:13, DM:9, Ca pulm:2, ethylism:1, haematologic disease:2, ulcerationGI:1, renal failure:1, autoimmune disease:1	27,(5.1%)	34 (6.4%)
Cardiovascular diseases MI, arrhythmias, angina, HF	47 (28%)	COPD:15, Asthma:1, acutebronchitis:1, embolipulm:1, metpulm.5, HF:5,cardiovasc 2, HT:3, AF:3, empyema: 1	133 (25%)	135 (25%)
Diabetes Mellitus	3 (1.8%)	Syncope:11, Ca corpora uteri:1, SLE, nephritis:1, HT:3, Venous ulcer:1,COPD:1, osteoporos:1, anemia 1, pleuritis:1, op heart valve: 1, anxiety:1 depression:1, Alzheimer:1, met pulm:2	13 (2.4%)	2 (0.4%)
Dizziness Vertigo, Syncope, Falls	19 (11%)	Migraine:1, hepatitis C:1, liver cirrhosis 2, methadone pancreatitis:2, ethylism:1, appendicitis:1, DM:2, HT:2, HF:2, AF:2 angina:1, RA:1, post stroke:1, pneumonia:1, ulcuceration: 2, gastritis:1	50 (9.4%)	53 (9.9%)
Dyspnea	38 (23%)	Cardiovascular dis: 5, Mb Crohn:2, COPD:2, DM:1, liver ca:1, ALS:1, hematologic disease: 1, KLL:1, angina p:1,	152 (28%)	180 (34%)
Impaired general condition	18 (11%)	Renal failure: 2, ulcerative legs: 3, perianal abscess:1, hyper- hypoglycemia:3, hypothermia:1, cough:2, traffic incidence:1, nausea:1	59 (11%)	70 (13%)
Infections	10 (5.9%)	Cancer:2, DM:1, COPD:2, AF:1, HF:2, HT:3, cardiovascular disease:1, vertebral fracture: 1, renal failure:3	25 (4.7%)	11 (2.1%)
Intoxications Unconsciousness	6 (3.6%)	SLE nephritis:1, DM:3, HT:3, Psychiatric diseases:3, TX:1, HD:1	19 (3.6%)	17 (3.2%)
Miscellaneous	6 (3.6%)	Rectal bleeding:3 ulcerative colitis:1, RA:1, GI bleedings:2, haematemesis:1, urinary cancer:1, DM:1, COPD:1	26 (4.9%)	16 (3.0%)
Pain	13 (7.7%)	HT:1, renal failure:1 vascular disease:1, RA:1, HT:1, AF:1	29 (5.4%)	16 (3.0%)
Total	168 (100%)		533 (100%)	534 (100%)

Abbreviations: No. = number of patients with this diagnosis at admission (Table 1a) and number of clinically relevant discrepancies (Table 1b). COPD = chronic pulmonary disease, HT = hypertension, HF = heart failure, AF = atrial fibrillation, MI = myocardial infarction RA = rheumatoid arthritis, SLE = systemic lupus erythematosus, TX = transplantation, DM = diabetes mellitus, HD = hemodialysis, ALS = amyotrophic lateral sclerosis, ca = cancer, met = metastases, pulm = pulmonary

doi:10.1371/journal.pone.0128716.t001

medicated with more drugs ($p < 0.0001$, Mann Whitney test) compared to patients without clinically significant drug record discrepancies (Table 4). Cardiovascular symptoms/diseases and dyspnoea were the most common reasons and accounted for half of all admittances at the EW (Table 1b).

These symptoms were also linked to the highest rate of discrepancies, which generally were of the same order of magnitude in the GPr and the SPDR (Table 1b). Multiple caregivers were more common in patients with clinically significant drug record discrepancies ($p = 0.025$, Fishers exact test). There was no difference between men and women regarding the presence of drug record discrepancies ($p = 0.5714$, Fishers exact test). (Table 4)

Table 2. Number of discrepancies comparing the records from the admission chart (ACR), the Swedish Prescribed Drug Register (SPDR), and the General Practitioner register (GPr).

No. of drug list discrepancies	ACR / SPDR No. of patients (%)	ACR / GPr No. of patients (%)
0	35 (21%)	21 (18%)
1–3	70 (42%)	29 (24%)
4–6	40 (24%)	40 (34%)
7–9	15 (9%)	13 (11%)
>10	8 (5%) max 13	16 (13%) max 23
sum	168 (100%)	119 (100%)

The most common (54%) clinically significant drug record discrepancy was drugs dispensed according to the SPDR but not found in the admission chart record

(type A admission chart-/ SPDR +). This concerned 106 patients (63%).

Drugs being registered in the admission chart record but not present in the SPDR

(B admission chart+/SPDR-) constituted 18% of the total number of clinically significant drug record discrepancies, affecting 53 (32%) of the patients.

doi:10.1371/journal.pone.0128716.t002

Table 3. Distribution of different types of discrepancies between records from the admission chart (ACR), the Swedish Prescribed Drug Register (SPDR), and the General Practitioner register (GPr).

Types of discrepancies	ACR / SPDR % of discrepancies	ACR / GPr % of discrepancies
Type A	54%	40%
Type B	18%	50%
Type C	21%	10%
Type D	6%	-

Type A: medication present in SPDR or GPr but not in ACR.

Type B: medication present in ACR but not in SPDR or GPr.

Type C+D: discrepancy regarding dose/regime (C) or amount prescribed insufficient to last until admission (D) in medication present in both records.

doi:10.1371/journal.pone.0128716.t003

Drugs involved in clinically significant drug record discrepancies

In the comparison between admission chart record and the SPDR, CNS-active drugs were most commonly identified (in total 28%, whereof 62% neuroleptics, sedatives and hypnotics (N05) and 31% analgesics (N02)), followed by cardiovascular drugs (in total 22%, of which 28% were diuretics (C03), 20% beta-blocking agents (C07), and 20% renin angiotensin

Table 4. Potential risk factors for drug discrepancies when comparing records from the admission chart (ACR), the Swedish Prescribed Drug Register (SPDR), and the General Practitioner register (GPr).

	ACR / SPDR		ACR / GPr	
	≥ 1 drug list discrepancies (n = 133)	No drug list discrepancies (n = 35)	≥ 1 drug list discrepancies (n = 98)	No drug list discrepancies (n = 21)
Age, years (mean (SD))	68.2 (SD 16.2)	55. (SD 16.3)	68.9 (SD 15.2)	57.0 (SD 21.7)
Number of medications at admission (mean (SD))	8.3 (SD 6.5)	1.7 (SD 3.1)	7.5 (SD 6.4)	1.5 (SD 2.6)
>1 health care provider	34.6%	14.3%	40.8%	14.3%
Sex, male	48.1%	54.3%	47.6%	46.9%

doi:10.1371/journal.pone.0128716.t004

Table 5. Drugs classified by pharmacological groups according to the ATC classification system [10] involved in discrepancies in records from the admission chart (ACR), the Swedish Prescribed Drug Register (SPDR), and the General Practitioner register (GPr) as the numbers and the percentages of discrepancies.

ATC Drug Class	ACR / SPDR No. of Discrepancies (%)	ACR / GPr No. of Discrepancies (%)
A. Alimentary tract and metabolism	76 (14%)	93 (17%)
B. Blood and blood forming organs	41 (7.7%)	50 (9.4%)
C. Cardiovascular system	118 (22%)	144 (27%)
D. Dermatological	5 (0.9%)	3 (0.6%)
G. Genito-urinary tract and sex-hormones	18 (3.4%)	14 (2.6%)
H. Hormones	13 (2.4%)	15 (2.8%)
J. Infectious diseases	9 (1.7%)	7 (1.3%)
L. Tumors and immune system	13 (2.4%)	5 (0.9%)
M. Muscular-skeletal system	18 (3.4%)	24 (4.5%)
N. Nervous System	148 (28%)	106 (20%)
P. Anti-parasitic	1 (0.2%)	0 (0.0%)
R. Respiratory system	66 (12%)	64 (12%)
S. Sensory organs	3 (0.6%)	4(0.7%)
V. Various	4 (0.8%)	5 (0.9%)
Total	533 (100%)	534 (100%)

doi:10.1371/journal.pone.0128716.t005

blocking agents (C09)). A similar picture was seen comparing the admission chart and the GP record, where cardiovascular drugs were the largest group (in total 27%, of which 29% belonged to diuretics (C03), 20% beta-blocking agents (C07), and 20% cardiac therapy (C01), e.g. nitroglycerine) followed by CNS-active drugs (in total 20%, 59% neuroleptics, sedatives and hypnotics (N05) and 29% analgesics (N02)) (Table 5).

Discussion

As part of ongoing studies of drug related problems at the Karolinska University Hospital, Huddinge and Solna [2–6] we have quantified drug record discrepancies of potential clinical importance between different sources of information regarding current medication for patients admitted to the emergency ward.

We found that 79% of the included 168 patients had at least one clinically significant drug list discrepancy when comparing the admission chart record to the SPDR.

A similar picture was seen in the 119 patients for whom the admission chart was compared to the GP record. This indicates that a majority of patients are subject to a continuance gap in their preadmission medication regimen. This corresponds well with the findings of Ekedahl et al [11] who compared the admission chart medication list and the medications reported by the patients, to the SPDR in a similar Swedish population.

In our study, older patients using many drugs prescribed by multiple caregivers were more prone to clinically significant discrepancies. It can be hypothesized that these factors contribute to the comparatively large number of ADRs found in elderly patients [2, 4–6].

Patients admitted because of cardiovascular symptoms/diseases or dyspnoea represented the largest percentage of discrepancies (Table 1b). Correspondingly, among the medications used, cardiovascular and CNS active drugs were the most common ones and may be seen as risk factors. These medications are also among the most commonly prescribed drugs in the general population [12], frequently causing DRPs [2–5] which highlights the importance of these particular discrepancies (Table 5).

Some discrepancies may arise because many of these medications may need titration by the GP by verbal instruction, while others are prescribed for use when needed (e.g. nitroglycerine or hypnotics) and may therefore be “forgotten” by the patient when in the EW.

The SPDR is a unique initiative in Sweden, and its usability has not yet been evaluated in an emergency setting. The lack of correct information on the patient's medication and the consequences of this have, however, previously been reported. Several studies have evaluated the accuracy of the admission chart record compared with more thorough medication reconciliation strategies, such as second interviews [1, 2, 6, 9, 13], records from local pharmacies [14], accounts of drug dispensers collected from the patients' home [14–15] and/or records from primary care physicians [15–18]. In accordance with our findings, these have shown that discrepancies are common, seen in between 54–96% of studied patients. Between 27–59% of these drug record discrepancies were considered as potential medical risks. Our results lie in the upper end of the spectrum, possible reasons as to which will be discussed below.

Discrepancies between the admission chart record and the SPDR

The SPDR does not register OTC drugs (unless they are prescribed), but this only partly explains the large number of discrepancies between these two records. We found that more than 60% of the patients lacked a clinically important prescription drug in the admission chart record. This highlights an important limitation with the SPDR; if a treatment was stopped, the dose adjusted or the drug replaced by another, the SPDR will no longer be accurate.

The opposite problem, a drug present in the admission chart record, but not in the SPDR, was seen in 32% of the patients. Non-compliance, OTC drugs or sporadic medications could explain this. Another possibility is that the admission chart record was based on a previously recorded medication record erroneously assumed by the admitting physician to be correct.

Discrepancies between the admission chart record and the GP record

One possible reason for the drug record discrepancies could be the organisation of the health care in Sweden. The GP does not have unique responsibility for patients' medication. Infrequent consultations, multiple caregivers and poor transmission of information from hospital to GP, could explain the lack of concordance between records. In this study we only contacted the patients' GP, but not other outpatient caregivers.

Multiple caregivers may be a problem in urban areas since health care often is provided by several GPs, private specialists and hospital based doctors. The GPs in the Stockholm area accounted for only one third of medical consultations and prescriptions issued to the population [19]. The GPs share of total consultations and prescriptions is considerably higher in countries where the GPs have a gatekeeper role such as Denmark and the UK, where GP prescribing accounts for the majority of National Health Service drug prescribing [20–21]. All types of drug list discrepancies could also derive from transcription errors and non compliance.

Limitations of the study

This study was conducted in a tertiary care university hospital where many patients with multiple diagnoses such as advanced heart disease, cancer, renal and liver failure are treated and this may not be representative for hospitals in general. The patient number is small, but representative of the category of patients who are admitted to the EW in this hospital [2–6].

It was not our aim to determine which of the records that was most accurate. This would have required a more complex reconciliation process involving a second interview with the patient or a close caregiver, if at all possible.

In an attempt to assess the patient selection we compared the demographic characteristics of the excluded patients ($n = 27$) with the study population. No significant differences in age, average hospitalization-days or the number of medications upon admission were identified (data not shown). This comparison does not include 47 patients who were admitted and discharged outside office hours. We cannot rule out the possibility of selection bias. Patients admitted outside office hours but still present in the ward on the subsequent weekday were, however, included in the study, partly compensating for this.

The SPDR automatically registers prescribed drugs that are dispensed [7]. Data was acquired for a period of minimum six months prior to admission, based on the assumption that most drug prescriptions are renewed every 3–6 months. By doing so we might have missed prescriptions renewed annually, but also avoiding registering old and no longer relevant prescriptions. Acquiring data from a shorter time period from SPDR, we could have missed important prescriptions renewed less frequently.

This study did not determine if the drug discrepancies resulted in negative clinical outcome. The maximal follow up period at the hospital was three days and we did not trace the patients after that. In order to assess the clinical consequences of drug record discrepancies we would need a matched control group of patients with a correct list of drugs. In order to have sufficient statistical power, a study like that would have to be of considerable size.

Consequences

Clinically significant drug record discrepancies can lead to a number of errors in patient care, both during hospitalization and after discharge. DRPs, such as double prescriptions, may arise or simply remain undetected. The SPDR provides accurate information with regard to prescribed drugs only at the moment of drug dispensation, but is not necessarily correct and complete at the time of admission to the EW. National projects, where the most recent prescriptions and changes in dosage of the individual patient's medication can be accessed by all caregivers at all times, are on the way and much needed [22]. Drug related morbidity is a global problem and one of the most common reasons for hospitalization, particularly in the elderly [23–25].

The availability of data on dispensed drugs such as in the Swedish Prescribed Drug Register and similarly in all Nordic countries [26] is, however, not yet available in many countries. In an extensive review of Nordic publications we have not found any similar studies outside Sweden to compare our results with [27]. However, we have reasons to believe that this is a global problem, maybe less pronounced in settings where the GP has a gatekeeper function [20–21]. Relying on the SPDR as the only source of drug record may potentially jeopardize patient care. The extent of the problem has not yet been determined.

Every patient should have an accurate list of drugs that can be assessed not only upon admission at the emergency ward but also when consulting other healthcare units. Obtaining this goal should have a high priority within any healthcare system.

Acknowledgments

We want to express gratitude to RN Elin Kimland for bibliographic help.

Author Contributions

Performed the experiments: IE KW CAH COS JF. Analyzed the data: IE KW CAH COS JF. Contributed reagents/materials/analysis tools: IE KW. Wrote the paper: IE KW CAH UB COS IOC JF. Included patients, compared medical records from the different sources: IE KW.

References

1. Cornish PL, Knowles SR, Marchesano R, Tam V, Shadowitz S, MD, Juurlink DN et al. Unintended medication discrepancies at the time of hospital admission. *Arch Int Med* 2005; 165(4):424–9
2. von Euler M, Eliasson E, Öhlén G, Bergman U. Adverse drug reactions causing hospitalization can be monitored from computerized medical records and thereby indicate the quality of drug utilization. *Pharmacoepidemiol Drug Saf* 2006; 15:179–184 PMID: [16184621](#)
3. Fryckstedt J, Asker-Hagelberg C. Drug related problems are common in the emergency department of internal medicine. The cause of admission in almost every third patient according to a quality follow-up. [Article in Swedish] *Läkartidningen* 2008; 105: 894–8
4. Odar-Cederlöf I, Oskarsson P, Öhlén G, Tesfa Y, Bergendal A, Helldén A et al. Adverse drug effect as cause of hospital admission. Common drugs are the major part according to the cross-sectional study. [Article in Swedish] *Läkartidningen* 2008; 105: 890–893.
5. Paul E, End-Rodrigues T, Thylén P, Bergman U. Adverse drug reactions a common cause of hospitalization of the elderly. A clinical retrospective study. [Article in Swedish] *Läkartidningen* 2008; 105: 2338–2342.
6. Oskarsson P, Odar-Cederlöf I, Bergendal A, Hentschke M, Asker-Hagelberg C, Fryckstedt J et al. Marked differences about drugs purchased at pharmacies and current drug treatment at admission to an internal medicine emergency department—a pilot study. *Svenska Läkaresällskapets handlingar Hygiea* 2007; 116(1):72 [Abstract in Swedish] doi: [10.1093/humrep/deu149](#) PMID: [24939957](#)
7. Wettermark B, Hammar N, Fored CM, Leimanis A, Otterblad Olausson P, Bergman U, et al. The new Swedish Prescribed Drug Register—opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf* 2007; 16(7):726–35. PMID: [16897791](#)
8. Eriksson U, Nilsson AL. 2007 Läkemedelsförteckningen—Utvärderingsrapport. Stockholm: Sveriges Kommuner och Landsting;. Translation in English: Register of dispensed drugs assessment report. Swedish Associations of Local Authorities and Regions (SKL) 2007 [Article in Swedish]
9. Cars T, Wettermark B, Malmström ER, Ekevinger G, Vikström B, Bergman U et al. Extraction of electronic health record data in a hospital setting: comparison of automatic and semi-automatic methods using anti-TNF therapy as model. *Basic & Clinical Pharmacol & Toxicol*; 2013 Feb [Epub ahead of print]
10. WHO Collaborating Centre for Drug Statistics Methodology. Guidelines for ATC classification and DDD assignment 2008. Oslo 2009
11. Ekedahl A, Broselius H, Jönsson J, Karlsson H, Yngvesson M. Discrepancies between the electronic medical record, the prescriptions in the Swedish national prescription repository and the current medication reported by the patients. *Pharmacoepidemiol Drug Saf* 2011; 20:1177–1183. doi: [10.1002/pds.2226](#) PMID: [21858899](#)
12. Loikas D, Wettermark B, von Euler M, Bergman U, Schenck-Gustafsson K: Differences in drug utilization between men and women: a cross-sectional analysis of all dispensed drugs in Sweden. *BMJ Open* 2013; 3:e002378. doi: [10.1136/bmjopen-2012-002378](#) PMID: [23645921](#)
13. Lubowski TJ, Cronin LM, Pavelka MS, Briscoe-Dwyer LA, Briceland LL, Hamilton RA. Effectiveness of a Medication Reconciliation Project Conducted by PharmD Students. *Am J Pharm Educ* 2007; 71(5): 94. PMID: [17998991](#)
14. Miller SE, Miller S, Balon J, Helling TS. Medication Reconciliation in a Rural Trauma Population. *Ann Emerg Med* 2008; 52(5): 483–91. doi: [10.1016/j.annemergmed.2008.03.021](#) PMID: [18550221](#)
15. Nassaralla CL, Naessens JM, Chaudhry R, Hansen MA, Scheitel SM. Implementation of a medication reconciliation process in an ambulatory internal medicine clinic, *Qual Saf Health Care* 2007; 16: 90–94. PMID: [17403752](#)
16. Bedell SE, Jabbour S, Goldberg R, Glaser H, Gobble S, Young-Xu Y et al. Discrepancies in the use of medications: their extent and predictors in an outpatient practice. *Arch Intern Med*;2000; 160(14): 2129–34. PMID: [10904455](#)
17. Wagner MM, Hogan WR. The accuracy of medication data in an outpatient electronic medical record. *J Am Med Inform Assoc* 1996; 3: 234–44. PMID: [8723614](#)
18. Akwagyriam I, Goodyer LI, Harding L, Khakoo S, Millington H.; Drug history taking and the identification of drug related problems in an accident and emergency department. *J Accid Emerg Med* 1996; 13(3): 166–8 PMID: [8733649](#)
19. Wettermark B, Bergman U, Krakau I. Using aggregate data on dispensed drugs to evaluate the quality of prescribing in urban primary health care in Sweden. *Public Health* 2006; 120: 451–461 PMID: [16513149](#)
20. Bjerrum L, Bergman U. 2006 Wide variation in the number of different drugs prescribed by general practitioners. A prescription database study. *Scand J Prim Health Care* 2000; 18: 94–8. PMID: [10944063](#)

21. Department of Health. Prescriptions dispensed in the community: statistics for 1989 to 1999. Statistical bulletin 2000/20. London: Department of Health; 2000.
22. Eriksson Ulrika. Nationell läkemedelsstrategi. [http://www.skl.se/web/Lakemedel - Nationell lakemedelsstrategi.aspx](http://www.skl.se/web/Lakemedel_-_Nationell_lakemedelsstrategi.aspx). 2009; Accessed November 2011.
23. Lindley CM, Tully MP, Paramsothy V et al. Inappropriate medication is a major cause of adverse drug reactions in elderly patients. *Age Ageing* 1992; 21(4):294–300. PMID: [1514459](#)
24. Pirmohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* 2004; 329;15–19. PMID: [15231615](#)
25. Hellén A, Bergman U, von Euler M, Hentschke M, Odar-Cederlöf I, Öhlén G. Adverse Drug Reactions and Impaired Renal Function in Elderly Patients Admitted to the Emergency Department. A Retrospective Study. *Drugs Aging* 2009; 26 (7): 595–606 doi: [10.2165/11315790-000000000-00000](#) PMID: [19655826](#)
26. Furu K, Wettermark B, Sørensen HT, Martikainen J, Almarsdóttir AB, Andersen M. The Nordic Countries as a cohort for pharmacoepidemiological research. *Basic Clin Pharmacol Toxicol* 2010; 106:86–94 doi: [10.1111/j.1742-7843.2009.00494.x](#) PMID: [19961477](#)
27. Wettermark B; Zoëga H; Furu K; Korhonen M; Hallas J; Nørgaard M et al. The Nordic Prescription Databases as a Resource for Pharmacoepidemiological Research—A Literature Review. *Pharmacoepidemiology and Drug Safety* 2013 Published online in Wiley Online Library (wileyonlinelibrary.com) doi: [10.1002/pds.3457](#)