

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

***rac-cis*-Cyclohexane-1,2-dicarboxylic acid–isoquinoline (1/1)**

Graham Smith\* and Urs D. Wermuth

Faculty of Science and Technology, Queensland University of Technology, GPO Box 2434, Brisbane, Queensland 4001, Australia  
Correspondence e-mail: g.smith@qut.edu.au


Received 29 July 2011; accepted 29 July 2011

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.038;  $wR$  factor = 0.090; data-to-parameter ratio = 14.3.

In the crystal structure of the title molecular adduct,  $\text{C}_9\text{H}_7\text{N}\cdot\text{C}_8\text{H}_{12}\text{O}_4$ , the two species are linked through a carboxylic acid–isoquinoline  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bond. These molecular pairs then inter-associate through the second acid group of the *cis*-cyclohexane-1,2-dicarboxylic acid molecules, forming a classic centrosymmetric cyclic head-to-head carboxylic acid–carboxyl  $\text{O}-\text{H}\cdots\text{O}$  hydrogen-bonding association [graph-set  $R_2^2(8)$ ], giving a zero-dimensional (cluster) structure, consisting of two of each species.

## Related literature

For the structure of racemic *cis*-cyclohexane-1,2-dicarboxylic acid, see: Benedetti *et al.* (1970). For the structures of the racemic 1:1 ammonium and 2-aminopyridinium salts of this acid, see: Smith & Wermuth (2011*a,b*). For the structure of the 1:1 adduct with 4,4'-bipyridine, see: Bhogala *et al.* (2005). For hydrogen bonding in carboxylic acids and graph-set analysis, see: Leiserowitz (1976); Etter *et al.* (1990).



## Experimental

## Crystal data

$\text{C}_9\text{H}_7\text{N}\cdot\text{C}_8\text{H}_{12}\text{O}_4$   
 $M_r = 301.33$   
 Triclinic,  $P\bar{1}$   
 $a = 6.2459$  (3) Å

$b = 11.4238$  (6) Å  
 $c = 11.9970$  (6) Å  
 $\alpha = 64.082$  (5)°  
 $\beta = 77.793$  (4)°

$\gamma = 82.756$  (4)°  
 $V = 751.95$  (7) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation

$\mu = 0.10$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.40 \times 0.28 \times 0.20$  mm

## Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer  
 Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)  
 $T_{\min} = 0.974$ ,  $T_{\max} = 0.990$

9094 measured reflections  
 2952 independent reflections  
 2463 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.022$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.090$   
 $S = 1.02$   
 2952 reflections  
 207 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.17$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.18$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O11}-\text{H11}\cdots\text{O12}^i$	0.96 (2)	1.68 (2)	2.6362 (14)	171.7 (18)
$\text{O22}-\text{H22}\cdots\text{N2A}$	0.98 (2)	1.69 (2)	2.670 (2)	174.5 (19)

Symmetry code: (i)  $-x + 1, -y + 2, -z + 2$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *WinGX* (Farrugia, 1999); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON* (Spek, 2009).

The authors acknowledge financial support from the Australian Research Council, the Faculty of Science and Technology and the University Library, Queensland University of Technology.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2108).

## References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.  
 Benedetti, E., Pedone, C. & Allegra, G. (1970). *J. Phys. Chem.* **74**, 512–516.  
 Bhogala, B. R., Basavoju, S. & Nangia, A. (2005). *CrystEngComm*, **7**, 551–562.  
 Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.  
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
 Leiserowitz, L. (1976). *Acta Cryst.* **B32**, 775–802.  
 Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Smith, G. & Wermuth, U. D. (2011*a*). *Acta Cryst.* **E67**, o174.  
 Smith, G. & Wermuth, U. D. (2011*b*). *Acta Cryst.* **E67**, o1900.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2011). E67, o2261 [ doi:10.1107/S1600536811030613 ]

## *rac-cis*-Cyclohexane-1,2-dicarboxylic acid-isoquinoline (1/1)

G. Smith and U. D. Wermuth

### Comment

Although the structure of racemic *cis*-cyclohexane-1,2-dicarboxylic acid is known (Benedetti *et al.*, 1970), together with its 1:1 adduct with 4,4'-bipyridine (Bhogala *et al.*, 2005), there are few examples of salts of this *cis*-acid in the crystallographic literature. We have previously reported the structures of the anhydrous 1:1 ammonium salt (Smith & Wermuth, 2011*a*) and the 2-aminopyridinium salt (Smith & Wermuth, 2011*b*). Our 1:1 stoichiometric interaction of cyclohexane-1,2-dicarboxylic anhydride with isoquinoline in 50% ethanol–water solution gave minor crystals of the 1:1 adduct C<sub>8</sub>H<sub>12</sub>O<sub>4</sub>.C<sub>9</sub>H<sub>7</sub>N, formed in a residual oil, and the structure is reported here.

In the structure of the title adduct (Fig. 1), the two molecular species are interlinked through a carboxylic acid *O*—H $\cdots$ N<sub>isoquinoline</sub> hydrogen bond (Table 1). The molecule pairs then associate through the second acid group, forming a classic centrosymmetric cyclic head-to-head carboxylic acid–carboxyl *O*—H $\cdots$ O hydrogen-bonding interaction (Leiserowitz, 1976) [graph set *R*<sup>2</sup><sub>2</sub>(8) (Etter *et al.*, 1990)] giving a zero-dimensional structure (Fig. 2).

### Experimental

The title compound was synthesized by heating a solution of 1 mmol of cyclohexane-1,2-dicarboxylic anhydride and 1 mmol of isoquinoline in 50 ml of 1:1 ethanol–water under reflux for 10 min. After concentration to 30 ml the solution was allowed to evaporate at room temperature, giving a viscous oil which eventually gave minor colourless crystals (m.p. 439–441 K) from which a specimen was cleaved for the X-ray analysis.

### Refinement

The carboxylic acid H atoms were located by difference methods and their positional and isotropic displacement parameters were refined. Other H-atoms were included in the refinement at calculated positions [C—H = 0.93–0.98 Å and with *U*<sub>iso</sub>(H) = 1.2*U*<sub>eq</sub>(aromatic C), or 1.5*U*<sub>eq</sub>(aliphatic C), using a riding-model approximation.

### Figures




Fig. 1. Atom numbering scheme for the two molecules in the title adduct. The inter-species hydrogen bond is shown as a dashed line and displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. The cyclic carboxylic acid hydrogen-bonding interactions between the acid–base molecular pairs, showing hydrogen bonds as dashed lines. Non-associative H atoms are omitted. For symmetry codes, see Table 1.

## *rac-cis*-Cyclohexane-1,2-dicarboxylic acid–isoquinoline (1/1)

### Crystal data

$C_9H_7N \cdot C_8H_{12}O_4$	$Z = 2$
$M_r = 301.33$	$F(000) = 320$
Triclinic, $PT$	$D_x = 1.331 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Melting point = 439–441 K
$a = 6.2459 (3) \text{ \AA}$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$b = 11.4238 (6) \text{ \AA}$	Cell parameters from 4474 reflections
$c = 11.9970 (6) \text{ \AA}$	$\theta = 3.3\text{--}28.7^\circ$
$\alpha = 64.082 (5)^\circ$	$\mu = 0.10 \text{ mm}^{-1}$
$\beta = 77.793 (4)^\circ$	$T = 200 \text{ K}$
$\gamma = 82.756 (4)^\circ$	Block, colourless
$V = 751.95 (7) \text{ \AA}^3$	$0.40 \times 0.28 \times 0.20 \text{ mm}$

### Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer	2952 independent reflections
Radiation source: Enhance (Mo) X-ray source graphite	2463 reflections with $I > 2\sigma(I)$
Detector resolution: $16.077 \text{ pixels mm}^{-1}$	$R_{\text{int}} = 0.022$
$\omega$ scans	$\theta_{\text{max}} = 26.0^\circ$ , $\theta_{\text{min}} = 3.3^\circ$
Absorption correction: multi-scan ( <i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	$h = -7 \rightarrow 7$
$T_{\text{min}} = 0.974$ , $T_{\text{max}} = 0.990$	$k = -14 \rightarrow 14$
9094 measured reflections	$l = -14 \rightarrow 14$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.038$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.090$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.02$	$w = 1/[\sigma^2(F_o^2) + (0.0354P)^2 + 0.20P]$
2952 reflections	where $P = (F_o^2 + 2F_c^2)/3$
207 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 0.17 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.18 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O11	0.70832 (16)	1.07685 (11)	0.86864 (10)	0.0405 (4)
O12	0.37261 (16)	1.01881 (10)	0.88359 (9)	0.0371 (3)
O21	0.49576 (18)	0.88369 (10)	0.70388 (12)	0.0476 (4)
O22	0.17799 (18)	0.95466 (11)	0.63667 (11)	0.0468 (4)
C1	0.5909 (2)	1.13098 (13)	0.67706 (12)	0.0272 (4)
C2	0.4067 (2)	1.11424 (13)	0.61893 (12)	0.0272 (4)
C3	0.1978 (2)	1.19528 (13)	0.63551 (13)	0.0298 (4)
C4	0.2466 (2)	1.33837 (14)	0.58905 (14)	0.0338 (4)
C5	0.4233 (2)	1.35310 (13)	0.65186 (13)	0.0314 (4)
C6	0.6329 (2)	1.27692 (13)	0.62795 (13)	0.0309 (4)
C11	0.5440 (2)	1.06979 (12)	0.81900 (13)	0.0271 (4)
C21	0.3661 (2)	0.97179 (14)	0.65964 (13)	0.0308 (4)
N2A	0.1006 (2)	0.70353 (12)	0.71708 (12)	0.0366 (4)
C1A	0.2157 (2)	0.61693 (14)	0.79877 (14)	0.0343 (5)
C3A	-0.0816 (3)	0.66352 (15)	0.69941 (14)	0.0375 (5)
C4A	-0.1451 (2)	0.53805 (15)	0.76063 (14)	0.0357 (5)
C5A	-0.0839 (3)	0.31194 (15)	0.92123 (15)	0.0390 (5)
C6A	0.0362 (3)	0.22778 (16)	1.00839 (16)	0.0434 (5)
C7A	0.2218 (3)	0.26877 (16)	1.02899 (15)	0.0421 (5)
C8A	0.2845 (2)	0.39437 (15)	0.96060 (14)	0.0365 (5)
C9A	0.1621 (2)	0.48468 (14)	0.87014 (13)	0.0304 (4)
C10A	-0.0255 (2)	0.44335 (14)	0.84987 (13)	0.0303 (4)
H1	0.72500	1.08880	0.64990	0.0410*
H2	0.46140	1.14950	0.52810	0.0410*
H11	0.669 (3)	1.037 (2)	0.959 (2)	0.080 (7)*
H22	0.158 (3)	0.862 (2)	0.663 (2)	0.078 (6)*
H31	0.09420	1.18840	0.58900	0.0450*
H32	0.13100	1.16090	0.72380	0.0450*
H41	0.29490	1.37600	0.49850	0.0510*
H42	0.11370	1.38550	0.60730	0.0510*
H51	0.37080	1.32150	0.74180	0.0470*
H52	0.45410	1.44450	0.61900	0.0470*
H61	0.74230	1.28670	0.66960	0.0460*
H62	0.68990	1.31220	0.53830	0.0460*

## supplementary materials

---

H1A	0.34010	0.64420	0.81040	0.0410*
H3A	-0.16580	0.72460	0.64320	0.0450*
H4A	-0.26770	0.51450	0.74370	0.0430*
H5A	-0.20530	0.28290	0.90850	0.0470*
H6A	-0.00490	0.14190	1.05510	0.0520*
H7A	0.30170	0.21020	1.08930	0.0510*
H8A	0.40850	0.42070	0.97350	0.0440*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O11	0.0362 (6)	0.0521 (7)	0.0326 (6)	-0.0141 (5)	-0.0111 (5)	-0.0120 (5)
O12	0.0331 (6)	0.0448 (6)	0.0305 (5)	-0.0113 (5)	-0.0061 (4)	-0.0105 (5)
O21	0.0450 (6)	0.0329 (6)	0.0715 (8)	0.0055 (5)	-0.0211 (6)	-0.0251 (6)
O22	0.0487 (7)	0.0313 (6)	0.0680 (8)	-0.0050 (5)	-0.0269 (6)	-0.0197 (6)
C1	0.0240 (7)	0.0283 (7)	0.0305 (7)	-0.0012 (5)	-0.0032 (5)	-0.0141 (6)
C2	0.0303 (7)	0.0285 (7)	0.0247 (7)	-0.0036 (6)	-0.0036 (5)	-0.0128 (6)
C3	0.0272 (7)	0.0299 (7)	0.0326 (7)	-0.0030 (6)	-0.0061 (6)	-0.0126 (6)
C4	0.0332 (7)	0.0284 (7)	0.0380 (8)	-0.0008 (6)	-0.0084 (6)	-0.0116 (6)
C5	0.0372 (8)	0.0241 (7)	0.0319 (7)	-0.0056 (6)	-0.0045 (6)	-0.0106 (6)
C6	0.0300 (7)	0.0322 (8)	0.0298 (7)	-0.0085 (6)	-0.0029 (6)	-0.0116 (6)
C11	0.0275 (7)	0.0228 (7)	0.0334 (7)	0.0003 (5)	-0.0081 (6)	-0.0131 (6)
C21	0.0345 (7)	0.0326 (8)	0.0300 (7)	-0.0022 (6)	-0.0053 (6)	-0.0174 (6)
N2A	0.0422 (7)	0.0331 (7)	0.0390 (7)	-0.0029 (5)	-0.0073 (6)	-0.0189 (6)
C1A	0.0326 (8)	0.0379 (8)	0.0420 (8)	-0.0036 (6)	-0.0051 (6)	-0.0256 (7)
C3A	0.0438 (9)	0.0391 (8)	0.0354 (8)	0.0015 (7)	-0.0132 (7)	-0.0190 (7)
C4A	0.0350 (8)	0.0431 (9)	0.0393 (8)	-0.0030 (6)	-0.0101 (6)	-0.0249 (7)
C5A	0.0368 (8)	0.0376 (8)	0.0493 (9)	-0.0062 (6)	-0.0046 (7)	-0.0247 (8)
C6A	0.0498 (10)	0.0316 (8)	0.0468 (9)	-0.0026 (7)	-0.0025 (8)	-0.0172 (7)
C7A	0.0483 (9)	0.0386 (9)	0.0417 (9)	0.0091 (7)	-0.0118 (7)	-0.0201 (7)
C8A	0.0347 (8)	0.0414 (9)	0.0429 (9)	0.0045 (6)	-0.0108 (7)	-0.0262 (7)
C9A	0.0308 (7)	0.0347 (8)	0.0333 (7)	0.0003 (6)	-0.0037 (6)	-0.0225 (6)
C10A	0.0312 (7)	0.0345 (8)	0.0330 (7)	-0.0015 (6)	-0.0025 (6)	-0.0226 (6)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O11—C11	1.3175 (17)	C4—H41	0.9700
O12—C11	1.2237 (17)	C5—H51	0.9700
O21—C21	1.2082 (19)	C5—H52	0.9700
O22—C21	1.3178 (18)	C6—H62	0.9700
O11—H11	0.96 (2)	C6—H61	0.9700
O22—H22	0.98 (2)	C1A—C9A	1.415 (2)
N2A—C3A	1.366 (2)	C3A—C4A	1.361 (2)
N2A—C1A	1.314 (2)	C4A—C10A	1.413 (2)
C1—C2	1.5356 (19)	C5A—C6A	1.361 (3)
C1—C11	1.5081 (19)	C5A—C10A	1.415 (2)
C1—C6	1.543 (2)	C6A—C7A	1.408 (3)
C2—C21	1.519 (2)	C7A—C8A	1.365 (3)
C2—C3	1.532 (2)	C8A—C9A	1.414 (2)

C3—C4	1.527 (2)	C9A—C10A	1.420 (2)
C4—C5	1.5243 (19)	C1A—H1A	0.9300
C5—C6	1.525 (2)	C3A—H3A	0.9300
C1—H1	0.9800	C4A—H4A	0.9300
C2—H2	0.9800	C5A—H5A	0.9300
C3—H31	0.9700	C6A—H6A	0.9300
C3—H32	0.9700	C7A—H7A	0.9300
C4—H42	0.9700	C8A—H8A	0.9300
C11—O11—H11	109.3 (12)	C6—C5—H51	110.00
C21—O22—H22	110.7 (12)	C4—C5—H51	109.00
C1A—N2A—C3A	118.09 (14)	H51—C5—H52	108.00
C2—C1—C6	110.04 (11)	C6—C5—H52	110.00
C6—C1—C11	109.54 (12)	C1—C6—H61	109.00
C2—C1—C11	113.10 (11)	C1—C6—H62	109.00
C1—C2—C21	112.12 (11)	C5—C6—H62	109.00
C3—C2—C21	113.55 (11)	H61—C6—H62	108.00
C1—C2—C3	113.23 (12)	C5—C6—H61	109.00
C2—C3—C4	111.43 (11)	N2A—C1A—C9A	124.04 (13)
C3—C4—C5	111.18 (12)	N2A—C3A—C4A	122.78 (15)
C4—C5—C6	110.73 (13)	C3A—C4A—C10A	120.23 (14)
C1—C6—C5	111.33 (11)	C6A—C5A—C10A	120.47 (17)
O11—C11—C1	112.94 (11)	C5A—C6A—C7A	121.01 (17)
O12—C11—C1	124.77 (12)	C6A—C7A—C8A	120.05 (16)
O11—C11—O12	122.29 (13)	C7A—C8A—C9A	120.39 (14)
O22—C21—C2	112.83 (13)	C1A—C9A—C8A	122.95 (13)
O21—C21—C2	123.78 (13)	C1A—C9A—C10A	117.55 (13)
O21—C21—O22	123.33 (16)	C8A—C9A—C10A	119.49 (14)
C2—C1—H1	108.00	C4A—C10A—C5A	124.13 (14)
C11—C1—H1	108.00	C4A—C10A—C9A	117.27 (14)
C6—C1—H1	108.00	C5A—C10A—C9A	118.58 (14)
C1—C2—H2	106.00	N2A—C1A—H1A	118.00
C21—C2—H2	106.00	C9A—C1A—H1A	118.00
C3—C2—H2	106.00	N2A—C3A—H3A	119.00
C2—C3—H31	109.00	C4A—C3A—H3A	119.00
C2—C3—H32	109.00	C3A—C4A—H4A	120.00
C4—C3—H32	109.00	C10A—C4A—H4A	120.00
H31—C3—H32	108.00	C6A—C5A—H5A	120.00
C4—C3—H31	109.00	C10A—C5A—H5A	120.00
C3—C4—H42	109.00	C5A—C6A—H6A	120.00
C5—C4—H41	109.00	C7A—C6A—H6A	119.00
C5—C4—H42	109.00	C6A—C7A—H7A	120.00
H41—C4—H42	108.00	C8A—C7A—H7A	120.00
C3—C4—H41	109.00	C7A—C8A—H8A	120.00
C4—C5—H52	110.00	C9A—C8A—H8A	120.00
C3A—N2A—C1A—C9A	-0.3 (2)	C3—C4—C5—C6	57.52 (15)
C1A—N2A—C3A—C4A	-1.4 (2)	C4—C5—C6—C1	-58.31 (15)
C6—C1—C2—C3	-52.58 (14)	N2A—C1A—C9A—C8A	-177.79 (15)
C11—C1—C2—C21	-59.79 (16)	N2A—C1A—C9A—C10A	1.2 (2)

## supplementary materials

---

C6—C1—C2—C21	177.35 (11)	N2A—C3A—C4A—C10A	2.1 (3)
C11—C1—C2—C3	70.29 (16)	C3A—C4A—C10A—C5A	177.60 (16)
C2—C1—C11—O11	175.82 (13)	C3A—C4A—C10A—C9A	-1.0 (2)
C2—C1—C11—O12	-4.5 (2)	C10A—C5A—C6A—C7A	-0.5 (3)
C6—C1—C11—O11	-61.03 (16)	C6A—C5A—C10A—C4A	-177.82 (16)
C6—C1—C11—O12	118.67 (16)	C6A—C5A—C10A—C9A	0.8 (2)
C11—C1—C6—C5	-69.79 (14)	C5A—C6A—C7A—C8A	-0.4 (3)
C2—C1—C6—C5	55.15 (14)	C6A—C7A—C8A—C9A	1.0 (3)
C21—C2—C3—C4	-178.03 (11)	C7A—C8A—C9A—C1A	178.30 (16)
C1—C2—C3—C4	52.62 (15)	C7A—C8A—C9A—C10A	-0.7 (2)
C3—C2—C21—O21	-148.08 (14)	C1A—C9A—C10A—C4A	-0.5 (2)
C3—C2—C21—O22	34.64 (16)	C1A—C9A—C10A—C5A	-179.24 (14)
C1—C2—C21—O21	-18.2 (2)	C8A—C9A—C10A—C4A	178.53 (14)
C1—C2—C21—O22	164.55 (12)	C8A—C9A—C10A—C5A	-0.2 (2)
C2—C3—C4—C5	-54.27 (15)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
O11—H11 $\cdots$ O12 <sup>i</sup>	0.96 (2)	1.68 (2)	2.6362 (14)	171.7 (18)
O22—H22 $\cdots$ N2A	0.98 (2)	1.69 (2)	2.670 (2)	174.5 (19)
C3—H32 $\cdots$ O12	0.97	2.54	3.1126 (17)	117
C6—H61 $\cdots$ O11	0.97	2.53	2.8841 (18)	101

Symmetry codes: (i)  $-x+1, -y+2, -z+2$ .



Fig. 1




Fig. 2

