

Role of STIM1 in neurodegeneration

Carlos Pascual-Caro, Noelia Espinosa-Bermejo, Eulalia Pozo-Guisado, Francisco Javier Martin-Romero

ORCID number: Carlos Pascual-Caro (0000-0002-2345-393X); Noelia Espinosa-Bermejo (0000-0002-6976-0871); Eulalia Pozo-Guisado (0000-0003-3486-6411); Francisco Javier Martin-Romero (0000-0001-6796-7396).

Author contributions: Martin-Romero FJ wrote the initial draft; Pascual-Caro C, Espinosa-Bermejo N and Pozo-Guisado E revised and reformatted the final version of the manuscript, together with Martin-Romero FJ; all authors approved the version to be published.

Supported by the Spanish Ministerio de Ciencia, Innovación y Universidades, No. BFU2017-82716-P.

Conflict-of-interest statement: The authors declare that they have no conflict of interests.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Francisco Javier Martin-Romero, PhD, Associate Professor, Department of

Carlos Pascual-Caro, Noelia Espinosa-Bermejo, Francisco Javier Martin-Romero, Department of Biochemistry and Molecular Biology, School of Life Sciences and Institute of Molecular Pathology Biomarkers, University of Extremadura, Badajoz 06006, Spain

Eulalia Pozo-Guisado, Department of Cell Biology, School of Medicine and Institute of Molecular Pathology Biomarkers, University of Extremadura, Badajoz 06006, Spain

Abstract

STIM1 is an endoplasmic reticulum (ER) protein with a key role in Ca²⁺ mobilization. Due to its ability to act as an ER-intraluminal Ca²⁺ sensor, it regulates store-operated Ca²⁺ entry (SOCE), which is a Ca²⁺ influx pathway involved in a wide variety of signalling pathways in eukaryotic cells. Despite its important role in Ca²⁺ transport, current knowledge about the role of STIM1 in neurons is much more limited. Growing evidence supports a role for STIM1 and SOCE in the preservation of dendritic spines required for long-term potentiation and the formation of memory. In this regard, recent studies have demonstrated that the loss of STIM1, which impairs Ca²⁺ mobilization in neurons, risks cell viability and could be the cause of neurodegenerative diseases. The role of STIM1 in neurodegeneration and the molecular basis of cell death triggered by low levels of STIM1 are discussed in this review.

Key words: Calcium; Neurodegeneration; Parkinson's disease; Alzheimer's disease; STIM1; Voltage-operated Ca²⁺ channels

©The Author(s) 2018. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: STIM1 is an endoplasmic reticulum protein that regulates store-operated Ca²⁺ entry, which is a Ca²⁺ influx pathway involved in a wide variety of signalling pathways. Growing evidence supports a role for this protein, STIM1, in long-term potentiation and the formation of memory. In this regard, the loss of STIM1 observed in brain tissue from Alzheimer's disease patients risks cell viability and could be the cause of neurodegenerative diseases. This is the reason for discussing the role of STIM1 in neurodegeneration in this review.

Pascual-Caro C, Espinosa-Bermejo N, Pozo-Guisado E, Martin-Romero FJ. Role of STIM1 in neurodegeneration. *World J Biol Chem* 2018; 9(2): 16-24

URL: <https://www.wjgnet.com/1949-8454/full/v9/i2/16.htm>

DOI: <https://dx.doi.org/10.4331/wjbc.v9.i2.16>

Biochemistry and Molecular Biology, School of Life Sciences and Institute of Molecular Pathology Biomarkers, University of Extremadura, Avenida de Elvas s/n, Badajoz 06006, Spain.

fjmartin@unex.es

Telephone: +34-92-4489971

Received: August 15, 2018

Peer-review started: August 17, 2018

First decision: September 11, 2018

Revised: October 8, 2018

Accepted: October 23, 2018

Article in press: October 23, 2018

Published online: November 16, 2018

STIM1 AND CALCIUM MOBILIZATION

STIM1 (stromal interaction molecule 1) is a type I transmembrane protein located mainly in the endoplasmic reticulum (ER), with a significant pool of approximately 20% at the plasma membrane. Due to its Ca²⁺-sensitive EF-hand domain close to the N-terminus, STIM1 acts as an ER-intraluminal Ca²⁺ sensor^[1,2]. This EF-hand domain shows an apparent dissociation constant for Ca²⁺ of 250 μmol/L^[3]. The decrease of the ER-intraluminal Ca²⁺ concentration, with the subsequent dissociation of Ca²⁺ from the EF-hand domain, triggers the oligomerization and the conformational change of STIM1. These two events are critical for STIM1 activation.

The rapid decrease of the ER-intraluminal Ca²⁺ concentration is a common event in cells under diverse stimuli, such as the activation of growth factor receptors or the activation of G protein-coupled receptors. In both cases, phosphoinositide-specific phospholipase C hydrolyzes phosphatidylinositol 4,5-bisphosphate to generate inositol 1,4,5-trisphosphate (IP₃) and diacylglycerol. The generation of IP₃ activates its receptor at the ER, with the subsequent release of Ca²⁺ through this channel/receptor and the rise of cytosolic free Ca²⁺ concentration ([Ca²⁺]_c). As mentioned above, the emptying of intracellular Ca²⁺ stores (mainly the ER) activates STIM1, which is then able to bind and activate STIM1-dependent Ca²⁺ channels^[4], such as ORAI1^[5]. The activation of ORAI1 leads to the transient increase of Ca²⁺ influx and to the rise of [Ca²⁺]_c, which is required for the refilling of the ER and for the sustainability of this system in successive stimulations. Thus, STIM1 protein and STIM1-dependent Ca²⁺ channels ensure Ca²⁺ mobilization and the stimulation of Ca²⁺-dependent signaling pathways by activating the “store-operated Ca²⁺ entry” (SOCE), *i.e.*, the Ca²⁺ influx pathway activated by the decrease of the ER-intraluminal Ca²⁺ level.

The activation of plasma membrane Ca²⁺ channels by STIM1 is carried out in ER-plasma membrane contact sites (ER-PM junctions)^[6], where STIM1 relocates in response to Ca²⁺ store depletion. When the ER-intraluminal Ca²⁺ concentration is high, STIM1 remains bound to the growing tip of microtubules and moves freely on the ER surface^[7]. However, activated STIM1 becomes phosphorylated at three ERK1/2-target sites (Ser575, Ser608, and Ser621) and this phosphorylation is critical for enhancing the dissociation from microtubules^[8,9]. Oligomers of active STIM1 are less mobile and phospho-STIM1 is found at the cell periphery^[10], close to the plasma membrane, where it binds ORAI1. Because STIM1 and ORAI1 are ubiquitous, they are involved in a wide range of signaling pathways that regulate many cellular functions^[11]. However, the number of studies about the role of STIM1 in neuronal tissue is much more limited.

STIM1 EXPRESSION AND FUNCTION IN NEURONAL CELLS

STIM1 is widely expressed in the brain according to databases such as Expression Atlas (from the European Bioinformatics Institute, <http://www.ebi.ac.uk/gxa>) or UniGene (from the National Center for Biotechnology Information, <https://www.ncbi.nlm.nih.gov/unigene>). Indeed, it is well known that STIM1 becomes activated upon depletion of intracellular Ca²⁺ stores in the brain in a similar fashion to that found in any other cell or tissue^[12,13]. The role of STIM1 in neuronal function was initially suggested in *Drosophila melanogaster* neurons. Shortly after the description of STIM1 as the main regulator of SOCE, it was proved that STIM1 was required for normal flight and associated patterns of rhythmic firing of the flight motoneurons^[14], and that SOCE regulates spatial and temporal Ca²⁺ mobilization in vertebrate photoreceptor cones, suggesting a role in the generation of excitatory signals across the retinal synapse^[15].

A key finding was reported in 2010 by Ricardo Dolmetsch's and Donald L. Gill's labs. They found that STIM1 directly suppresses depolarization-induced opening of the voltage-operated Ca²⁺ channel (VOCC) Ca_v1.2^[16,17]. What was striking was the fact that STIM1 binds to Ca_v1.2 through the same domain that activates ORAI1, the Ca²⁺ release-activated Ca²⁺ activation domain, and also triggers the internalization of the channel from the membrane. These findings provided the molecular explanation for the shared control of Ca²⁺ entry through ORAI1 and Ca_v1.2, making it possible for them to operate independently. In HEK293 cells, it was later reported that Homer proteins are required for the binding between STIM1 and Ca_v1.2 channels upon Ca²⁺ store-depletion conditions triggered by thapsigargin^[18], an inhibitor of the ER-Ca²⁺ pump.

T-type VOCCs, such as Ca_v3.1, are also modulated by STIM1. This was first observed not in neurons but in cardiomyocytes, where it was reported that STIM1 co-precipitated with Ca_v1.3 channels, and that the knocking-down of STIM1 expression

increased $\text{Ca}_v1.3$ surface expression and the current density of T-type VOCCs^[19].

Given the abundance of STIM1 and STIM2 in neuronal tissues and their role in Ca^{2+} mobilization, it is not surprising to learn that they have a direct impact on cognitive functions. In mice with conditional deletion of *Stim1* or *Stim2* genes in the forebrain (conditional knock-outs or cKO), the analysis of spatial reference memory revealed a mild learning delay in *Stim1* cKO mice, no effect in *Stim2* cKO mice, and a deep impairment in spatial learning in the double cKO^[20]. This striking effect was explained by the regulation of the phosphorylation of the AMPA receptor subunit GluA1, the transcriptional regulator CREB and the $\text{Ca}_v1.2$ on protein kinase A-target sites, leading to the proposal that the upregulation of cAMP/PKA signaling impairs the development of spatial memory^[20]. Kuznicki's lab reported that STIM1 protein in neurons can control AMPA-induced Ca^{2+} entry, based on the inhibition of Ca^{2+} entry observed with AMPA receptors (AMPA) inhibitors and the finding that STIM1 physically binds GluA1/GluA2 AMPAR^[21].

On the other hand, in transgenic mice overexpressing STIM1 in neurons it was reported a reduction of long-term depression in hippocampal slices, as well as a decrease in anxiety-like behavior and an increase in contextual learning improvement^[22]. All of this further confirms the role of STIM1 in the modulation of synaptic strength and memory formation.

Closely related to the above statement, the control of L-type VOCCs by STIM1 has functional consequences that were reported for dendritic spine structural plasticity. In hippocampal neurons, depolarization by the neurotransmitter glutamate activates postsynaptic N-methyl-D-aspartate receptors and L-type VOCC-dependent Ca^{2+} influx, as well as the release of Ca^{2+} from the ER. The consequent activation of STIM1 inhibits VOCCs, an event that leads to the enlargement of ER content in spines, which is believed to help in the stabilization of mushroom spines that have become enlarged during long-term potentiation^[23].

STIM1 IN NEURONAL CELL DEATH

There are some examples of the involvement of STIM1 and SOCE in neuronal injury. For instance, cell death due to diffuse axonal injury is preceded by an increase of STIM1 expression in neurons of the rat cerebral cortex after lateral head rotational injury^[24]. In this regard, STIM1 expression was significantly increased in a traumatic brain injury model, and STIM1 knock-down inhibited apoptotic cell death after traumatic injury by decreasing the upregulation of mGluR1-dependent Ca^{2+} signaling^[25]. However, Berna-Ero *et al*^[26] demonstrated that STIM2, but not STIM1, was essential for ischemia-induced cytosolic Ca^{2+} accumulation in neurons using hypoxic conditions for culturing neurons from wild-type and *Stim2*^{-/-} mice, hippocampal slice preparations, as well as in *Stim2*^{-/-} mice subjected to focal cerebral ischemia.

Oxytosis, a type of cell death characterized by an increase of reactive oxygen species (ROS) and augmented Ca^{2+} influx, can be triggered in neurons in culture by depleting reduced glutathione content. Henke *et al*^[27], reported that the Ca^{2+} -influx pathway in this cell death could be mediated by ORAI1, the CRAC channel activated by STIM1. Similarly, in PC12 cells exposed to 6-hydroxydopamine (6-OHDA), an experimental model to trigger ROS-dependent cell death, the knockdown of STIM1 was able to attenuate apoptotic cell death by limiting the mitochondrial Ca^{2+} uptake induced by 6-OHDA. This resulted in the protection of PC12 cells against the oxidative stress generated by ER stress and mitochondrial dysfunction^[28]. On the other hand, the inhibition of SOCE or the knock-down of STIM1 limited ROS production and the activation of apoptosis in PC12 cells exposed to 1-methyl-4-phenylpyridinium or MPP⁺^[29], the toxic metabolite of MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine), a well-known inducer of Parkinsonism. These studies revealed that oxidative stress induces an increase of $[\text{Ca}^{2+}]_i$, mediated by the activation of SOCE. Indeed, STIM1 is a redox-sensitive protein, and it is known that Cys56 becomes S-glutathionylated during oxidative stress^[30], a residue located near its luminal EF-hand domain. Hawkins *et al*^[30], demonstrated that S-glutathionylation lowered the affinity of STIM1 for Ca^{2+} , thereby activating STIM1 in a store-independent fashion. Similarly, ORAI1 is a redox sensor through the Cys195 located in the second extracellular loop. Although it was initially shown that the oxidation of this Cys residue inhibited Ca^{2+} current through this channel^[31], other researchers found that exposure to H_2O_2 increased influx through ORAI1^[32], suggesting that ROS has multiple redox-sensitive targets in the SOCE machinery.

Mitochondrial dysfunction is an early event in neurotoxicity triggered by massive Ca^{2+} influx, as observed during glutamate neurotoxicity^[33]. Following acute increase in

$[Ca^{2+}]_i$, Ca^{2+} uptake by mitochondria contributes to the protection against cell death. However, Ca^{2+} overload in mitochondria triggers the opening of the mitochondrial permeability transition pore (mPTP), an event that led to cell death in different neuronal cell types^[34,35]. It is accepted that the overproduction of ROS modulates the opening of the mPTP, but it has also been shown this opening at physiological levels of ROS. Recently, Agarwal *et al.*^[36], reported that astrocytes show transient cytosolic Ca^{2+} spikes generated by the Ca^{2+} release from mitochondria when the mPTP opens by a mechanism that involves ROS generated during the electron transfer in the respiratory systems. Electron transport rates are strongly dependent on the availability of NADH, and therefore dependent on the Krebs cycle status, which is tightly controlled by the mitochondrial $[Ca^{2+}]$. Therefore, there is a strong correlation between dysregulation of Ca^{2+} entry through ORAI1, mitochondrial Ca^{2+} overload, ROS generation, mPTP opening and cell death.

STIM1 IN NEURODEGENERATIVE DISEASES

Alzheimer's disease

Taking into consideration the information summarized above, it should not be surprising that the dysregulation of STIM1 could underlie the pathogenesis of some of the most frequent neurodegenerative diseases. In 1907 Alzheimer^[37] described a disease in a 51-year-old woman with presenile dementia who displayed diffuse cortical atrophy, nerve cell loss, plaques, and tangles. Nowadays, Alzheimer's disease (AD) patients are classified within 3 groups: Early-onset AD (up to 5% of all patients with AD), late-onset or sporadic AD (the most common form of the disease), and familial AD (FAD, less than 1% of AD patients). FAD is linked to known genes, such as the amyloid beta precursor protein gene (*APP*), the apolipoprotein E gene (*APOE*), presenilin1/2 genes (*PSEN1*, *PSEN2*), or the alpha-2-macroglobulin gene (*A2M*), and most early-onset AD patients are FAD patients. There is no significant pathological difference between sporadic AD and FAD, but symptoms progress more rapidly in FAD^[38].

The major risk for sporadic AD is aging, which increases the difficulty of finding a suitable model animal that recapitulates all the hallmarks of the human disease in the absence of mutated genes as in FAD. However, there is a growing consensus regarding the hypothesis that Ca^{2+} dysregulation is in the pathogenesis of AD^[39-42]. This hypothesis is supported by evidence that revealed how diverse Ca^{2+} mobilization systems are impaired in AD, including VOCCs, IP_3 receptors, store-operated Ca^{2+} channels (SOCs), and mitochondrial Ca^{2+} transporters^[43].

Regarding STIM1 and SOCE, it is known that SOCE is reduced and that STIM1 and ORAI1 expression are downregulated in long-term cultures of hippocampal neurons, an experimental approach intended to mimic *in vivo* neuronal aging^[44]. Also, reduced expression of STIM2 was observed in hippocampal neurons from the presenilin-1 M146V knock-in mouse model of FAD. As it is assumed that STIM2 and the activation of the calmodulin-dependent protein kinase II (CaMKII) mediates the stabilization of mushroom spines, this decrease in STIM2 levels could explain the loss of dendritic spines and the defects in the development of long-term potentiation LTP and memory development in AD patients^[45]. In this regard, it is known that the gamma-secretase protein complex interacts with STIM1 in SH-SY5Y neuroblastoma cells, skin fibroblasts from FAD patients, and in mouse primary cortical neurons^[46]. Tong *et al.*^[46], also reported that cultured hippocampal neurons expressing the mutant PSEN1 M146L, showed reduced dendritic spines, together with diminished SOCE. Because the wild-type phenotype was rescued by overexpressing STIM1, or by inhibiting gamma-secretase activity, they hypothesized that STIM1 could be a substrate for the gamma-secretase complex. Finally, they proved that the transmembrane domain of STIM1 shows a target domain for the proteolytic activity of the gamma-secretase complex and that the reduced SOCE in PSEN1-mutant neurons was due to the higher rates of STIM1 proteolysis. Although this proteolysis needs to be studied further to confirm cleavage sites on STIM1, this data fits well with the recent observation that there is a sharp decline of STIM1 protein levels in brain tissue from non-familial (sporadic) AD patients^[47]. This is supporting evidence for a common hallmark in sporadic AD and FAD, *i.e.*, reduced STIM1 could be severely affecting Ca^{2+} mobilization in neurons in both groups of patients. Thus, it is necessary to study the consequences of the reduced STIM1 expression in neurons in order to understand how neuronal cell physiology develops in the absence of STIM1 and to find possible targets for clinical interventions. An approach to studying the patho-physiological consequences of a limited level of STIM1 in neurons has been recently reported by our group^[47]. In this report, we modified *STIM1* gene locus using CRISPR/Cas9-mediated

editing techniques, and we found that the differentiation of SH-SY5Y cells to neuronal-like cells was not impaired by the absence of STIM1. However, the loss of STIM1 triggered significant cell death due to the impairment of mitochondrial respiratory chain complex I, and to reduced mitochondrial Ca^{2+} concentration. These two events led to high levels of senescence. STIM1-KO cells showed potentiation of Ca^{2+} entry through L-type VOCCs^[47], further confirming earlier observations that demonstrated the inhibitory role of STIM1 on $\text{Ca}_v1.2$ channels^[16,17]. Consequently, the knocking-down of *CACNA1C* gene transcripts (for $\text{Ca}_v1.2$ channel) rescued the wild-type phenotype, confirming that the upregulation of Ca^{2+} entry through $\text{Ca}_v1.2$ channels was deleterious in STIM1-deficient cells^[47] (Figure 1). In this regard, higher Ca^{2+} entry through VOCCs had been recorded in CA1 pyramidal neurons from the hippocampus in aged rats^[48], an effect that resulted in the down-regulation of short-term neuronal plasticity.

Accumulation of beta amyloid peptides ($\text{A}\beta$) begins earlier than most of the clinical symptoms associated with FAD. However, clinical interventions to prevent this accumulation have been inconclusive so far. Accumulation of $\text{A}\beta$ directly affects Ca^{2+} mobilization, and the possibility that an increase of PKA-dependent phosphorylation of $\text{Ca}_v1.2$ channels could underlie the upregulation of Ca^{2+} influx through these channels has been discussed^[49,50]. Therefore, an alternative clinical intervention is the blocking of excessive Ca^{2+} entry in neurons. Transgenic mice have been designed to accumulate $\text{A}\beta$ and hyperphosphorylation of tau protein in CA1 pyramidal neurons, as an experimental approach to mimic some clinical features of FAD patients. Using these mice (known as 3xTgAD mice) it has been shown that Ca^{2+} current through L-type VOCCs became higher in these hippocampal neurons, supporting the possible role of VOCCs in neuronal degeneration in FAD patients^[51]. In addition, the long-term treatment of subjects receiving active treatment with L-type VOCCs blockers (nitrendipine) reduced sporadic dementia by 55% during aging^[52], suggesting that the enhanced Ca^{2+} entry through VOCCs could be in the pathogenesis of sporadic AD. Protection against the loss of working memory has also been monitored in rats treated with the VOCC blocker nimodipine, a treatment that reduced Ca^{2+} current through $\text{Ca}_v1.3$ in CA1 neurons^[53]. Finally, isradipine, another dihydropyridine, attenuated $\text{A}\beta$ accumulation toxicity by reducing $\text{Ca}_v1.2$ expression and Ca^{2+} influx in MC65 neuroblastoma cells^[54]. Interestingly, isradipine also showed a neuroprotective effect in models of Parkinson's disease (PD) and stroke^[50,55].

Whereas a decline in STIM1 level is deleterious, in part due to the upregulation of VOCCs, high levels of STIM1 and SOCE might be protective, as suggested by the reduced $\text{A}\beta$ secretion observed in cells expressing a constitutively activated STIM1 mutant (D76A)^[56]. On the other hand, $\text{A}\beta$ seems to affect STIM1-dependent Ca^{2+} entry because knocking-down APP transcripts delayed the binding of STIM1 to ORAI1 in response to store depletion^[57], and SOCE was largely reduced in cultured astrocytes from APP-KO mice^[58], confirming the crosstalk between SOCE and APP.

PD

A recent report showed that neurotoxins that trigger PD symptoms targeted TRPC1 expression and increased Ca^{2+} influx through $\text{Ca}_v1.3$ channels (L-type VOCC) which led to degeneration of dopaminergic (DA) neurons^[59]. Because of the key role of $\text{Ca}_v1.3$ in the regulation of basal single-spike firing in DA neurons^[60], the reported inhibition of $\text{Ca}_v1.3$ by the STIM1-TRPC1 complex^[59] could explain the disruption of neuronal Ca^{2+} homeostasis in PD patients. Indeed, in mice treated with MPTP, the expression of $\text{Ca}_v1.2$ and $\text{Ca}_v1.3$ in the substantia nigra increased after 2 wk of treatment, and isradipine (L-type VOCC blocker) prevented this upregulation and the loss of DA neurons^[61]. Similarly, nimodipine prevented cell death triggered by MPP^+ in SH-SY5Y cell in culture and Parkinsonism in MPTP-treated mice^[62]. Because dihydropyridines are not highly selective in discriminating between $\text{Ca}_v1.2$ and $\text{Ca}_v1.3$, Wang *et al.*^[61], reported a high-throughput screening that led to 1-(3-chlorophenethyl)-3-cyclopentylpyrimidine-2,4,6-(1H,3H,5H)-trione as the first potent and highly selective $\text{Ca}_v1.3$ antagonist with potential utility in clinical approaches. However, Ortner *et al.*^[63], reported later that this specific compound showed inhibitory activity of $\text{Ca}_v1.3$ only in a minority of cells.

On the other hand, antagonists of SOCE and depletion of STIM1 by siRNA increased cell viability, reduced intracellular ROS production as well as lipid peroxidation and prevented mitochondrial dysfunction in MPP^+ -treated PC12 cells^[29], supporting the hypothesis that augmented Ca^{2+} entry through STIM1-activated channels mediates toxicity of MPP^+ . This result, however, is in conflict with the observation that treatment with this neurotoxin decreased TRPC1 expression, TRPC1




Figure 1 Deficiency of STIM1 and neurodegeneration. Neurons expressing the mutant PSEN1 M146L showed higher rates of STIM1 proteolysis, reduced levels of STIM1, reduced store-operated Ca²⁺ entry and diminished dendritic spines⁴⁶. Deficiency of STIM1 has been observed in non-familial (sporadic) Alzheimer's disease (AD) patients, and can be mimicked by genome edition of STIM1 locus in SH-SY5Y cells⁴⁷. Because STIM1 is a negative regulator of Ca_v1.2 channels, this deficiency triggered the upregulation of Ca²⁺ entry through Ca_v1.2 channels which was responsible for the loss of inner mitochondrial membrane polarization, senescence, and cell death⁴⁷. This higher rate of Ca²⁺ influx through Ca_v1.2 channels has also been monitored in 3xTgAD mice⁵¹. The long-term treatment with dihydropyridines, known blockers of Ca_v1.2, reduced sporadic dementia by 55% during aging⁵², pointing out the decrease of STIM1 as a possible mechanism to explain neurodegeneration in sporadic and familial AD.

interaction with STIM1, and Ca²⁺ entry in SH-SY5Y cells⁶⁴, making further study necessary to discover the role of STIM1, SOCE, and VOCCs in the pathogenesis of PD.

CONCLUSION

Neurodegenerative diseases are devastating for the elderly population and no fully efficient therapies are available to treat some of them, particularly AD. However, a growing body of evidence supports a role for excessive Ca²⁺ entry through VOCCs in neurodegeneration. Recent reports proposed that the specific loss of STIM1 in neuronal tissue fully explains the observed Ca²⁺ homeostasis disruption in neurons during sporadic AD and FAD. In this regard, STIM1 deficiency triggered upregulation of Ca²⁺ entry through Ca_v1.2 in differentiated SH-SY5Y cells, which can be explained by the role of STIM1 in the inhibitory control of Ca_v1.2. This augmented Ca²⁺ influx led to the inhibition of the mitochondrial respiratory chain complex activity, mitochondrial inner membrane depolarization, reduced mitochondrial free Ca²⁺ concentration, and to higher levels of senescence and cell death. All these effects were prevented by silencing Ca_v1.2 expression, emphasizing the upregulation of these channels as a major cause of neuronal cell death (Figure 1).

REFERENCES

- 1 **Liou J**, Kim ML, Heo WD, Jones JT, Myers JW, Ferrell JE Jr, Meyer T. STIM is a Ca²⁺ sensor essential for Ca²⁺-store-depletion-triggered Ca²⁺ influx. *Curr Biol* 2005; **15**: 1235-1241 [PMID: 16005298 DOI: 10.1016/j.cub.2005.05.055]
- 2 **Zhang SL**, Yu Y, Roos J, Kozak JA, Deerinck TJ, Ellisman MH, Stauderman KA, Cahalan MD. STIM1 is a Ca²⁺ sensor that activates CRAC channels and migrates from the Ca²⁺ store to the plasma membrane. *Nature* 2005; **437**: 902-905 [PMID: 16208375 DOI: 10.1038/nature04147]
- 3 **Stathopoulos PB**, Li GY, Plevin MJ, Ames JB, Ikura M. Stored Ca²⁺ depletion-induced oligomerization of stromal interaction molecule 1 (STIM1) via the EF-SAM region: An initiation mechanism for capacitive Ca²⁺ entry. *J Biol Chem* 2006; **281**: 35855-35862 [PMID: 17020874 DOI: 10.1074/jbc.M608247200]
- 4 **Zhou Y**, Srinivasan P, Razavi S, Seymour S, Meraner P, Gudlur A, Stathopoulos PB, Ikura M, Rao A, Hogan PG. Initial activation of STIM1, the regulator of store-operated calcium entry. *Nat Struct Mol Biol* 2013; **20**: 973-981 [PMID: 23851458 DOI: 10.1038/nsmb.2625]
- 5 **Prakriya M**, Feske S, Gwack Y, Srikanth S, Rao A, Hogan PG. Orai1 is an essential pore subunit of the CRAC channel. *Nature* 2006; **443**: 230-233 [PMID: 16921383 DOI: 10.1038/nature05122]
- 6 **Ong HL**, Liu X, Tsaneva-Atanasova K, Singh BB, Bandyopadhyay BC, Swaim WD, Russell JT, Hegde RS, Sherman A, Ambudkar IS. Relocalization of STIM1 for activation of store-operated Ca(2+) entry is determined by the depletion of subplasma membrane endoplasmic reticulum

- Ca(2+) store. *J Biol Chem* 2007; **282**: 12176-12185 [PMID: [17298947](#) DOI: [10.1074/jbc.M609435200](#)]
- 7 **Grigoriev I**, Gouveia SM, van der Vaart B, Demmers J, Smyth JT, Honnappa S, Splinter D, Steinmetz MO, Putney JW Jr, Hoogenraad CC, Akhmanova A. STIM1 is a MT-plus-end-tracking protein involved in remodeling of the ER. *Curr Biol* 2008; **18**: 177-182 [PMID: [18249114](#) DOI: [10.1016/j.cub.2007.12.050](#)]
 - 8 **Pozo-Guisado E**, Casas-Rua V, Tomas-Martin P, Lopez-Guerrero AM, Alvarez-Barrientos A, Martin-Romero FJ. Phosphorylation of STIM1 at ERK1/2 target sites regulates interaction with the microtubule plus-end binding protein EB1. *J Cell Sci* 2013; **126**: 3170-3180 [PMID: [23687376](#) DOI: [10.1242/jcs.125054](#)]
 - 9 **Pozo-Guisado E**, Martin-Romero FJ. The regulation of STIM1 by phosphorylation. *Commun Integr Biol* 2013; **6**: e26283 [PMID: [24505502](#) DOI: [10.4161/cib.26283](#)]
 - 10 **Lopez-Guerrero AM**, Tomas-Martin P, Pascual-Caro C, Macartney T, Rojas-Fernandez A, Ball G, Alessi DR, Pozo-Guisado E, Martin-Romero FJ. Regulation of membrane ruffling by polarized STIM1 and ORAI1 in cortactin-rich domains. *Sci Rep* 2017; **7**: 383 [PMID: [28341841](#) DOI: [10.1038/s41598-017-00331-4](#)]
 - 11 **Martin-Romero FJ**, Pascual-Caro C, Espinosa-Bermejo N, Pozo-Guisado E. Regulation of calcium signaling by STIM1 and ORAI1, in Calcium and Signal Transduction. *Buchholz JN, Editor* Intech Open: London; 2018 [DOI: [10.5772/intechopen.78587](#)]
 - 12 **Klejman ME**, Gruszczynska-Biegala J, Skibinska-Kijek A, Wisniewska MB, Misztal K, Blazejczyk M, Bojarski L, Kuznicki J. Expression of STIM1 in brain and puncta-like co-localization of STIM1 and ORAI1 upon depletion of Ca(2+) store in neurons. *Neurochem Int* 2009; **54**: 49-55 [PMID: [19013491](#) DOI: [10.1016/j.neuint.2008.10.005](#)]
 - 13 **Skibinska-Kijek A**, Wisniewska MB, Gruszczynska-Biegala J, Methner A, Kuznicki J. Immunolocalization of STIM1 in the mouse brain. *Acta Neurobiol Exp (Wars)* 2009; **69**: 413-428 [PMID: [20048759](#)]
 - 14 **Venkiteswaran G**, Hasan G. Intracellular Ca²⁺ signaling and store-operated Ca²⁺ entry are required in Drosophila neurons for flight. *Proc Natl Acad Sci USA* 2009; **106**: 10326-10331 [PMID: [19515818](#) DOI: [10.1073/pnas.0902982106](#)]
 - 15 **Szikra T**, Barabas P, Bartoletti TM, Huang W, Akopian A, Thoreson WB, Krizaj D. Calcium homeostasis and cone signaling are regulated by interactions between calcium stores and plasma membrane ion channels. *PLoS One* 2009; **4**: e6723 [PMID: [19696927](#) DOI: [10.1371/journal.pone.0006723](#)]
 - 16 **Park CY**, Shcheglovitov A, Dolmetsch R. The CRAC channel activator STIM1 binds and inhibits L-type voltage-gated calcium channels. *Science* 2010; **330**: 101-105 [PMID: [20929812](#) DOI: [10.1126/science.1191027](#)]
 - 17 **Wang Y**, Deng X, Mancarella S, Hendron E, Eguchi S, Soboloff J, Tang XD, Gill DL. The calcium store sensor, STIM1, reciprocally controls Orai and CaV1.2 channels. *Science* 2010; **330**: 105-109 [PMID: [20929813](#) DOI: [10.1126/science.1191086](#)]
 - 18 **Dionisio N**, Smani T, Woodard GE, Castellano A, Salido GM, Rosado JA. Homer proteins mediate the interaction between STIM1 and Cav1.2 channels. *Biochim Biophys Acta* 2015; **1853**: 1145-1153 [PMID: [25712868](#) DOI: [10.1016/j.bbamcr.2015.02.014](#)]
 - 19 **Nguyen N**, Biet M, Simard E, Béliveau E, Francoeur N, Guillemette G, Dumaine R, Grandbois M, Boulay G. STIM1 participates in the contractile rhythmicity of HL-1 cells by moderating T-type Ca(2+) channel activity. *Biochim Biophys Acta* 2013; **1833**: 1294-1303 [PMID: [23458835](#) DOI: [10.1016/j.bbamcr.2013.02.027](#)]
 - 20 **Garcia-Alvarez G**, Shetty MS, Lu B, Yap KA, Oh-Hora M, Sajikumar S, Bichler Z, Fivaz M. Impaired spatial memory and enhanced long-term potentiation in mice with forebrain-specific ablation of the Stim genes. *Front Behav Neurosci* 2015; **9**: 180 [PMID: [26236206](#) DOI: [10.3389/fnbeh.2015.00180](#)]
 - 21 **Gruszczynska-Biegala J**, Sladowska M, Kuznicki J. AMPA Receptors Are Involved in Store-Operated Calcium Entry and Interact with STIM Proteins in Rat Primary Cortical Neurons. *Front Cell Neurosci* 2016; **10**: 251 [PMID: [27826230](#) DOI: [10.3389/fncel.2016.00251](#)]
 - 22 **Majewski Ł**, Maciąg F, Boguszewski PM, Wasilewska I, Wiera G, Wójtowicz T, Mozrzyk J, Kuznicki J. Overexpression of STIM1 in neurons in mouse brain improves contextual learning and impairs long-term depression. *Biochim Biophys Acta Mol Cell Res* 2017; **1864**: 1071-1087 [PMID: [27913207](#) DOI: [10.1016/j.bbamcr.2016.11.025](#)]
 - 23 **Dittmer PJ**, Wild AR, Dell'Acqua ML, Sather WA. STIM1 Ca²⁺ Sensor Control of L-type Ca²⁺-Channel-Dependent Dendritic Spine Structural Plasticity and Nuclear Signaling. *Cell Rep* 2017; **19**: 321-334 [PMID: [28402855](#) DOI: [10.1016/j.celrep.2017.03.056](#)]
 - 24 **Li Y**, Song J, Liu X, Zhang M, An J, Sun P, Li D, Jin T, Wang J. High expression of STIM1 in the early stages of diffuse axonal injury. *Brain Res* 2013; **1495**: 95-102 [PMID: [23261659](#) DOI: [10.1016/j.brainres.2012.12.005](#)]
 - 25 **Hou PF**, Liu ZH, Li N, Cheng WJ, Guo SW. Knockdown of STIM1 improves neuronal survival after traumatic neuronal injury through regulating mGluR1-dependent Ca(2+) signaling in mouse cortical neurons. *Cell Mol Neurobiol* 2015; **35**: 283-292 [PMID: [25304289](#) DOI: [10.1007/s10571-014-0123-0](#)]
 - 26 **Berna-Erro A**, Braun A, Kraft R, Kleinschnitz C, Schuhmann MK, Stegner D, Wultsch T, Eilers J, Meuth SG, Stoll G. STIM2 regulates capacitive Ca²⁺ entry in neurons and plays a key role in hypoxic neuronal cell death. *Sci Signal* 2009; **2**: ra67 [PMID: [19843959](#) DOI: [10.1126/scisignal.2000522](#)]
 - 27 **Henke N**, Albrecht P, Bouchachia I, Ryazantseva M, Knoll K, Lewerenz J, Kaznacheyeva E, Maher P, Methner A. The plasma membrane channel ORAI1 mediates detrimental calcium influx caused by endogenous oxidative stress. *Cell Death Dis* 2013; **4**: e470 [PMID: [23348584](#) DOI: [10.1038/cddis.2012.216](#)]
 - 28 **Li B**, Xiao L, Wang ZY, Zheng PS. Knockdown of STIM1 inhibits 6-hydroxydopamine-induced oxidative stress through attenuating calcium-dependent ER stress and mitochondrial dysfunction in undifferentiated PC12 cells. *Free Radic Res* 2014; **48**: 758-768 [PMID: [24720513](#) DOI: [10.3109/10715762.2014.905687](#)]
 - 29 **Li X**, Chen W, Zhang L, Liu WB, Fei Z. Inhibition of store-operated calcium entry attenuates MPP(+)-induced oxidative stress via preservation of mitochondrial function in PC12 cells: involvement of Homer1a. *PLoS One* 2013; **8**: e83638 [PMID: [24358303](#) DOI: [10.1371/journal.pone.0083638](#)]

- 10.1371/journal.pone.0083638]
- 30 **Hawkins BJ**, Irrinki KM, Mallilankaraman K, Lien YC, Wang Y, Bhanumathy CD, Subbiah R, Ritchie MF, Soboloff J, Baba Y. S-glutathionylation activates STIM1 and alters mitochondrial homeostasis. *J Cell Biol* 2010; **190**: 391-405 [PMID: 20679432 DOI: 10.1083/jcb.201004152]
- 31 **Bogeski I**, Kummerow C, Al-Ansary D, Schwarz EC, Koehler R, Kozai D, Takahashi N, Peinelt C, Griesemer D, Bozem M. Differential redox regulation of ORAI ion channels: a mechanism to tune cellular calcium signaling. *Sci Signal* 2010; **3**: ra24 [PMID: 20354224 DOI: 10.1126/scisignal.2000672]
- 32 **Grube M**, Myers G, Penner R, Fleig A. Activation of store-operated I(CRAC) by hydrogen peroxide. *Cell Calcium* 2010; **48**: 1-9 [PMID: 20646759 DOI: 10.1016/j.ceca.2010.05.005]
- 33 **Schinder AF**, Olson EC, Spitzer NC, Montal M. Mitochondrial dysfunction is a primary event in glutamate neurotoxicity. *J Neurosci* 1996; **16**: 6125-6133 [PMID: 8815895 DOI: 10.1523/JNEUROSCI.16-19-06125.1996]
- 34 **Murchison D**, Griffith WH. Mitochondria buffer non-toxic calcium loads and release calcium through the mitochondrial permeability transition pore and sodium/calcium exchanger in rat basal forebrain neurons. *Brain Res* 2000; **854**: 139-151 [PMID: 10784115 DOI: 10.1016/S0006-8993(99)02297-0]
- 35 **Limke TL**, Atchison WD. Acute exposure to methylmercury opens the mitochondrial permeability transition pore in rat cerebellar granule cells. *Toxicol Appl Pharmacol* 2002; **178**: 52-61 [PMID: 11781080 DOI: 10.1006/taap.2001.9327]
- 36 **Agarwal A**, Wu PH, Hughes EG, Fukaya M, Tischfield MA, Langseth AJ, Wirtz D, Bergles DE. Transient Opening of the Mitochondrial Permeability Transition Pore Induces Microdomain Calcium Transients in Astrocyte Processes. *Neuron* 2018; **93**: 587-605.e7 [PMID: 28132831 DOI: 10.1016/j.neuron.2016.12.034]
- 37 **Alzheimer A**. Über eine eigenartige Erkrankung der Hirnrinde. *Allg Z Psychiat Med* 1907; 146-148
- 38 **Hardy J**, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002; **297**: 353-356 [PMID: 12130773 DOI: 10.1126/science.1072994]
- 39 **LaFerla FM**. Calcium dyshomeostasis and intracellular signalling in Alzheimer's disease. *Nat Rev Neurosci* 2002; **3**: 862-872 [PMID: 12415294 DOI: 10.1038/nrn960]
- 40 **Berridge MJ**. Calcium regulation of neural rhythms, memory and Alzheimer's disease. *J Physiol* 2014; **592**: 281-293 [PMID: 23753528 DOI: 10.1113/jphysiol.2013.257527]
- 41 **Raza M**, Deshpande LS, Blair RE, Carter DS, Sombati S, DeLorenzo RJ. Aging is associated with elevated intracellular calcium levels and altered calcium homeostatic mechanisms in hippocampal neurons. *Neurosci Lett* 2007; **418**: 77-81 [PMID: 17374449 DOI: 10.1016/j.neulet.2007.03.005]
- 42 **Popugaeva E**, Pchitskaya E, Bezprozvanny I. Dysregulation of neuronal calcium homeostasis in Alzheimer's disease - A therapeutic opportunity? *Biochem Biophys Res Commun* 2017; **483**: 998-1004 [PMID: 27641664 DOI: 10.1016/j.bbrc.2016.09.053]
- 43 **Tong BC**, Wu AJ, Li M, Cheung KH. Calcium signaling in Alzheimer's disease and therapies. *Biochim Biophys Acta Mol Cell Res* 2018; **1865**: 1745-1760 [PMID: 30059692 DOI: 10.1016/j.bbamcr.2018.07.018]
- 44 **Calvo-Rodríguez M**, García-Durillo M, Villalobos C, Núñez L. In vitro aging promotes endoplasmic reticulum (ER)-mitochondria Ca²⁺ cross talk and loss of store-operated Ca²⁺ entry (SOCE) in rat hippocampal neurons. *Biochim Biophys Acta* 2016; **1863**: 2637-2649 [PMID: 27503411 DOI: 10.1016/j.bbamcr.2016.08.001]
- 45 **Sun S**, Zhang H, Liu J, Popugaeva E, Xu NJ, Feske S, White CL 3rd, Bezprozvanny I. Reduced synaptic STIM2 expression and impaired store-operated calcium entry cause destabilization of mature spines in mutant presenilin mice. *Neuron* 2014; **82**: 79-93 [PMID: 24698269 DOI: 10.1016/j.neuron.2014.02.019]
- 46 **Tong BC**, Lee CS, Cheng WH, Lai KO, Foscett JK, Cheung KH. Familial Alzheimer's disease-associated presenilin 1 mutants promote γ -secretase cleavage of STIM1 to impair store-operated Ca²⁺ entry. *Sci Signal* 2016; **9**: ra89 [PMID: 27601731 DOI: 10.1126/scisignal.aaf1371]
- 47 **Pascual-Caro C**, Berrocal M, Lopez-Guerrero AM, Alvarez-Barrientos A, Pozo-Guisado E, Gutierrez-Merino C, Mata AM, Martin-Romero FJ. STIM1 deficiency is linked to Alzheimer's disease and triggers cell death in SH-SY5Y cells by upregulation of L-type voltage-operated Ca²⁺ entry. *J Mol Med (Berl)* 2018; **96**: 1061-1079 [PMID: 30088035 DOI: 10.1007/s00109-018-1677-y]
- 48 **Thibault O**, Hadley R, Landfield PW. Elevated postsynaptic [Ca²⁺]_i and L-type calcium channel activity in aged hippocampal neurons: relationship to impaired synaptic plasticity. *J Neurosci* 2001; **21**: 9744-9756 [PMID: 11739583 DOI: 10.1523/JNEUROSCI.21-24-09744.2001]
- 49 **Hall DD**, Davare MA, Shi M, Allen ML, Weisenhaus M, McKnight GS, Hell JW. Critical role of cAMP-dependent protein kinase anchoring to the L-type calcium channel Cav1.2 via A-kinase anchor protein 150 in neurons. *Biochemistry* 2007; **46**: 1635-1646 [PMID: 17279627 DOI: 10.1021/bi062217x]
- 50 **Anekonda TS**, Quinn JF. Calcium channel blocking as a therapeutic strategy for Alzheimer's disease: the case for isradipine. *Biochim Biophys Acta* 2011; **1812**: 1584-1590 [PMID: 21925266 DOI: 10.1016/j.bbadis.2011.08.013]
- 51 **Wang Y**, Mattson MP. L-type Ca²⁺ currents at CA1 synapses, but not CA3 or dentate granule neuron synapses, are increased in 3xTgAD mice in an age-dependent manner. *Neurobiol Aging* 2014; **35**: 88-95 [PMID: 23932880 DOI: 10.1016/j.neurobiolaging.2013.07.007]
- 52 **Forette F**, Seux ML, Staessen JA, Thijs L, Babarskiene MR, Babeanu S, Bossini A, Fagard R, Gil-Extremera B, Laks T. The prevention of dementia with antihypertensive treatment: new evidence from the Systolic Hypertension in Europe (Syst-Eur) study. *Arch Intern Med* 2002; **162**: 2046-2052 [PMID: 12374512 DOI: 10.1001/archinte.162.18.2046]
- 53 **Veng LM**, Mesches MH, Browning MD. Age-related working memory impairment is correlated with increases in the L-type calcium channel protein alpha1D (Cav1.3) in area CA1 of the hippocampus and both are ameliorated by chronic nimodipine treatment. *Brain Res Mol Brain Res* 2003; **110**: 193-202 [PMID: 12591156 DOI: 10.1016/S0169-328X(02)00643-5]
- 54 **Anekonda TS**, Quinn JF, Harris C, Frahler K, Wadsworth TL, Woltjer RL. L-type voltage-gated calcium channel blockade with isradipine as a therapeutic strategy for Alzheimer's disease. *Neurobiol Dis* 2011; **41**: 62-70 [PMID: 20816785 DOI: 10.1016/j.nbd.2010.08.020]

- 55 **Lenhard SC**, Strittmatter R, Price WJ, Chandra S, White RF, Barone FC. Brain MRI and neurological deficit measurements in focal stroke: rapid throughput validated with isradipine. *Pharmacology* 2008; **81**: 1-10 [PMID: [17726342](#) DOI: [10.1159/000107661](#)]
- 56 **Zeiger W**, Vetrivel KS, Buggia-Prévoit V, Nguyen PD, Wagner SL, Villereal ML, Thinakaran G. Ca²⁺ influx through store-operated Ca²⁺ channels reduces Alzheimer disease β-amyloid peptide secretion. *J Biol Chem* 2013; **288**: 26955-26966 [PMID: [23902769](#) DOI: [10.1074/jbc.M113.473355](#)]
- 57 **Gazda K**, Kuznicki J, Wegierski T. Knockdown of amyloid precursor protein increases calcium levels in the endoplasmic reticulum. *Sci Rep* 2017; **7**: 14512 [PMID: [29109429](#) DOI: [10.1038/s41598-017-15166-2](#)]
- 58 **Linde CI**, Baryshnikov SG, Mazzocco-Spezia A, Golovina VA. Dysregulation of Ca²⁺ signaling in astrocytes from mice lacking amyloid precursor protein. *Am J Physiol Cell Physiol* 2011; **300**: C1502-C1512 [PMID: [21368296](#) DOI: [10.1152/ajpcell.00379.2010](#)]
- 59 **Sun Y**, Zhang H, Selvaraj S, Sukumaran P, Lei S, Birnbaumer L, Singh BB. Inhibition of L-Type Ca²⁺ Channels by TRPC1-STIM1 Complex Is Essential for the Protection of Dopaminergic Neurons. *J Neurosci* 2017; **37**: 3364-3377 [PMID: [28258168](#) DOI: [10.1523/JNEUROSCI.3010-16.2017](#)]
- 60 **Liu Y**, Harding M, Pittman A, Dore J, Striessnig J, Rajadhyaksha A, Chen X. Cav1.2 and Cav1.3 L-type calcium channels regulate dopaminergic firing activity in the mouse ventral tegmental area. *J Neurophysiol* 2014; **112**: 1119-1130 [PMID: [24848473](#) DOI: [10.1152/jn.00757.2013](#)]
- 61 **Wang QM**, Xu YY, Liu S, Ma ZG. Isradipine attenuates MPTP-induced dopamine neuron degeneration by inhibiting up-regulation of L-type calcium channels and iron accumulation in the substantia nigra of mice. *Oncotarget* 2017; **8**: 47284-47295 [PMID: [28521299](#) DOI: [10.18632/oncotarget.17618](#)]
- 62 **Singh A**, Verma P, Balaji G, Samantaray S, Mohanakumar KP. Nimodipine, an L-type calcium channel blocker attenuates mitochondrial dysfunctions to protect against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced Parkinsonism in mice. *Neurochem Int* 2016; **99**: 221-232 [PMID: [27395789](#) DOI: [10.1016/j.neuint.2016.07.003](#)]
- 63 **Ortner NJ**, Bock G, Vandael DH, Mauersberger R, Draheim HJ, Gust R, Carbone E, Tuluc P, Striessnig J. Pyrimidine-2,4,6-triones are a new class of voltage-gated L-type Ca²⁺ channel activators. *Nat Commun* 2014; **5**: 3897 [PMID: [24941892](#) DOI: [10.1038/ncomms4897](#)]
- 64 **Selvaraj S**, Sun Y, Watt JA, Wang S, Lei S, Birnbaumer L, Singh BB. Neurotoxin-induced ER stress in mouse dopaminergic neurons involves downregulation of TRPC1 and inhibition of AKT/mTOR signaling. *J Clin Invest* 2012; **122**: 1354-1367 [PMID: [22446186](#) DOI: [10.1172/JCI61332](#)]

P- Reviewer: Nesci S, Utkin YN

S- Editor: Ji FF **L- Editor:** A **E- Editor:** Yin SY





Published By Baishideng Publishing Group Inc
7901 Stoneridge Drive, Suite 501, Pleasanton, CA 94588, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

