



The Fumarate Reductase of *Bacteroides thetaiotaomicron*, unlike That of *Escherichia coli*, Is Configured so that It Does Not Generate Reactive Oxygen Species

Zheng Lu, James A. Imlay

Department of Microbiology, University of Illinois, Urbana, Illinois, USA

ABSTRACT The impact of oxidative stress upon organismal fitness is most apparent in the phenomenon of obligate anaerobiosis. The root cause may be multifaceted, but the intracellular generation of reactive oxygen species (ROS) likely plays a key role. ROS are formed when redox enzymes accidentally transfer electrons to oxygen rather than to their physiological substrates. In this study, we confirm that the predominant intestinal anaerobe *Bacteroides thetaiotaomicron* generates intracellular ROS at a very high rate when it is aerated. Fumarate reductase (Frd) is a prominent enzyme in the anaerobic metabolism of many bacteria, including *B. thetaiotaomicron*, and prior studies of *Escherichia coli* Frd showed that the enzyme is unusually prone to ROS generation. Surprisingly, in this study biochemical analysis demonstrated that the *B. thetaiotaomicron* Frd does not react with oxygen at all: neither superoxide nor hydrogen peroxide is formed. Subunit-swapping experiments indicated that this difference does not derive from the flavoprotein subunit at which ROS normally arise. Experiments with the related enzyme succinate dehydrogenase discouraged the hypothesis that heme moieties are responsible. Thus, resistance to oxidation may reflect a shift of electron density away from the flavin moiety toward the iron-sulfur clusters. This study shows that the autoxidizability of a redox enzyme can be suppressed by subtle modifications that do not compromise its physiological function. One implication is that selective pressures might enhance the oxygen tolerance of an organism by manipulating the electronic properties of its redox enzymes so they do not generate ROS.

IMPORTANCE Whether in sediments or pathogenic biofilms, the structures of microbial communities are configured around the sensitivities of their members to oxygen. Oxygen triggers the intracellular formation of reactive oxygen species (ROS), and the sensitivity of a microbe to oxygen likely depends upon the rates at which ROS are formed inside it. This study supports that idea, as an obligate anaerobe was confirmed to generate ROS very rapidly upon aeration. However, the suspected source of the ROS was disproven, as the fumarate reductase of the anaerobe did not display the high oxidation rate of its *E. coli* homologue. Evidently, adjustments in its electronic structure can suppress the tendency of an enzyme to generate ROS. Importantly, this outcome suggests that evolutionary pressure may succeed in modifying redox enzymes and thereby diminishing the stress that an organism experiences in oxic environments. The actual source of ROS in the anaerobe remains to be discovered.

Received 7 October 2016 Accepted 10 November 2016 Published 3 January 2017

Citation Lu Z, Imlay JA. 2017. The fumarate reductase of *Bacteroides thetaiotaomicron*, unlike that of *Escherichia coli*, is configured so that it does not generate reactive oxygen species. mBio 8:e01873-16. doi:10.1128/mBio.01873-16.

Editor Matthew R. Chapman, University of Michigan

Copyright © 2017 Lu and Imlay. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to James A. Imlay, jimlay@illinois.edu.

This article is a direct contribution from a Fellow of the American Academy of Microbiology. External solicited reviewers: Gary Ceccini, UCSF School of Medicine; C. Jeffrey Smith, East Carolina University.

The oxygenation of the planet occurred late in evolutionary time (1), and it imposed a crisis upon extant microbes. Molecular oxygen is toxic. It can directly poison specialized free radical and low-potential enzymes that are found in some anaerobes (2, 3). More generally, oxygen also can intercept some of the electrons that flow through redox enzymes, thereby generating superoxide and hydrogen peroxide (4). These species are stronger oxidants than is oxygen itself, and they rapidly oxidize the exposed iron cofactors on families of [4Fe-4S] dehydratases (5–8) and mononuclear iron enzymes (9–11). The oxidized iron atoms dissociate from those enzymes, activities are lost, and their pathways stop working. The outcome is a cessation of metabolism and growth.

Such oxidant-sensitive enzymes are almost universally distributed through the biota, and so aerobic organisms have invented

ways to protect them. The primary defense is the synthesis of superoxide dismutases (SOD) that scavenge O_2^- and of peroxidases and catalases that scavenge H_2O_2 . In the model bacterium *Escherichia coli*, these enzymes are among the most abundant in the cell, yet their collective activity is barely sufficient to keep reactive oxygen species (ROS) concentrations below the threshold for overt toxicity (12). Mutant strains that lack either SOD or catalase/peroxidase activities are unable to grow in a simple glucose medium (13, 14).

Obligate anaerobiosis represents a scenario in which evolution has failed to equip microbes for oxygen exposure. The reason that oxygen poisons anaerobic metabolism is an issue of great interest to investigators of microbial ecology, pathogenesis, and industrial fermentations. Experimental evidence suggests that endogenous




FIG 1 Physical structure and metabolic role of *B. thetaiotaomicron* fumarate reductase. (A) Structural diagrams of *B. thetaiotaomicron* fumarate reductase (Frd), *E. coli* Frd, and *E. coli* succinate dehydrogenase (Sdh). Physiological directions of electron flow are indicated by arrows. MQ, menaquinone; UQ, ubiquinone. Both cluster-proximal and -distal quinone binding sites were visualized in the *E. coli* Frd structure (36), and they are tentatively inferred for the *B. thetaiotaomicron* structure (66). (B) Role of Frd in redox balancing during the fermentation of glucose by *B. thetaiotaomicron*. Glycolysis generates 2 mol of NADH, which are recycled by malate dehydrogenase and by the Frd-dependent electron transport chain. Red crosses indicate the inactivation of pyruvate:ferredoxin oxidoreductase (Por) and fumarase (Fum) when the cell is aerated.

ROS production may be a key element. *Bacteroides thetaiotaomicron* is a dominant obligate anaerobe in the human intestine (15), and it provides an apt contrast to *E. coli*. Both are intestinal microbes that catabolize carbohydrates found in that environment; however, whereas *E. coli* thrives upon excretion into oxic surface waters, *Bacteroides* becomes quiescent. Metabolic analysis indicates that oxygenation inactivates two key enzymes in the *B. thetaiotaomicron* central metabolism: pyruvate:ferredoxin oxidoreductase (POR), which may be directly damaged by oxygen itself, and fumarase (16). The latter enzyme belongs to the iron-sulfur dehydratase family that is especially vulnerable to O₂⁻ and H₂O₂. In *B. thetaiotaomicron*, these enzymes remain inactive for the duration of aeration; when anoxia is restored, the enzymes are reactivated, and growth resumes.

In striking contrast, the *E. coli* fumarase enzymes maintain full activity upon aeration. This discrepancy is unlikely to derive from a difference in the titers of scavenging enzymes in the two organisms. *B. thetaiotaomicron* exhibits SOD activity that is similar to that of *E. coli*, and it has four distinct enzymes that can degrade H₂O₂ (17). Instead, an attractive possibility is that these organisms differ in the rates at which ROS are formed inside the aerated cell. If ROS are formed especially rapidly in *B. thetaiotaomicron*, then standard levels of scavenging enzymes may be insufficient to protect ROS-sensitive enzymes.

Our understanding of the mechanism of ROS production has lagged behind our knowledge of ROS-mediated damage. *In vitro* studies have identified quite a few enzymes that release ROS as inadvertent by-products when they operate in oxic solutions (4, 18, 19). In each case, they are flavin-dependent redox enzymes, and O₂⁻ and H₂O₂ are formed when oxygen collides adventitiously with their flavins at the point in the catalytic cycle when the flavin is reduced. In the adventitious reactions, molecular oxygen competes with the physiological acceptor for the reduced enzyme.

The rates at which different flavoenzymes leak electrons to oxygen vary widely (18), and it seems likely that the organisms that struggle the most with oxygen are those with the highest titers of the leakiest enzymes. To identify such enzymes, it is useful to pinpoint the physical traits that predispose flavoenzymes to react with oxygen. Comparative studies have been performed upon

members of the complex II enzyme family. This family received particular attention because succinate dehydrogenase (Sdh) is a source of ROS within the mammalian respiratory chain (20–24). Sdh is a respiratory enzyme that transfers electrons from succinate to the quinone pool (Fig. 1A); autoxidation occurs when oxygen intercepts electrons from its reduced flavin (24, 25). The Sdh family also includes aspartate oxidase and fumarate reductase, both of which generate ROS markedly more quickly than does Sdh itself (25). Analysis of these enzymes revealed that ROS production is maximized when the flavin is highly solvent exposed, when it has a low reduction potential, and when it is the center of electron density on the reduced enzyme. These studies also revealed that aspartate oxidase and Frd have substantial impacts upon ROS levels in *E. coli*. Aspartate oxidase is a minor enzyme but is responsible for 30% of the H₂O₂ formed in the aerobic cell (26). Fumarate reductase (Frd) is the terminal enzyme for anaerobic respiration, allowing the cell to use fumarate in place of oxygen as an electron acceptor. It is a close structural homologue of Sdh, but its electronic properties make it much more prone to autoxidation (25). Although expression of the *E. coli* Frd is repressed in oxic environments, the autoxidation of the enzyme becomes a significant event when the bacterium moves from anoxic environments, in which Frd is well expressed, to oxic ones. During the initial period of aeration, intracellular ROS production surges, and the preexisting Frd is responsible (26, 27).

Whereas Frd has an ancillary role in the anaerobic metabolism of *E. coli*, it is a key enzyme in the central pathway of *B. thetaiotaomicron* (Fig. 1B). Thus, we suggested (16) that in this anaerobe the titers of Frd might be especially high and that upon aeration its contribution to ROS formation might be proportionately great. If so, then perhaps even high levels of scavenging enzymes might be inadequate to suppress steady-state levels of ROS, and fumarase inactivation might be the consequence. Subsequently, Meehan and Malamy tested this notion using *Bacteroides fragilis*, a relative that behaves as an obligate anaerobe but has features, such as an oxygen-dependent ribonucleotide reductase, that suggest greater oxygen tolerance. They observed that a *B. fragilis* strain with a mutation in a Frd subunit seemed to release H₂O₂ at a diminished rate (28). Their interpretation was that Frd is the primary source




FIG 2 Aeration blocks growth but does not diminish the viability of wild-type *B. thetaiotaomicron*. Cultures were grown in an anaerobic chamber to exponential phase and then transferred to fully oxic environments. The data are representative of three independent experiments.

of ROS in this bacterium and that this abundant ROS might be involved in poisoning metabolism upon aeration.

In the present study, we confirmed that H₂O₂ is produced in aerated *B. thetaiotaomicron* much more rapidly than in *E. coli*. However, biochemical analysis revealed the surprising fact that the Frd of *B. thetaiotaomicron* does not generate either O₂⁻ or H₂O₂. Instead, the iron-sulfur subunit of this enzyme suppresses flavin autoxidation, probably by pulling the electron density away from the solvent-exposed flavin. This example reveals that evolution can modify redox enzymes in a way that suppresses their autoxidation without compromising their physiological function. It is not clear whether this advantage provided the impetus for the electron arrangement of the *B. thetaiotaomicron* enzyme; however, the evolution of enzymes to minimize ROS formation would have been a natural complement to the appearance of other ROS defenses.

RESULTS

Hydrogen peroxide is rapidly formed inside *B. thetaiotaomicron* when it is aerated. The aeration of *Bacteroides thetaiotaomicron* in BHIS medium caused growth to arrest almost immediately (Fig. 2A). During exposure to oxygen for 11 h, the cells failed to grow but remained fully viable. When anoxia was restored, growth resumed after a short (60- to 90-min) delay (Fig. 2B). This ability to recover quickly from oxygen exposure likely helps *B. thetaiotaomicron* to transit from one host to another. In contrast, after the same period of aeration, *oxyR* mutants exhibited a longer lag of several hours. Closer examination revealed that the *oxyR* mutants suffered a substantial loss of viability during the period of oxygen

exposure (Fig. 3A); the lag in subsequent outgrowth reflected the fact that many of the cells were dead.

These results confirm a previous report (17) that *B. thetaiotaomicron*, like its relative *B. fragilis* (29), activates its OxyR response when it is aerated and that this adaptation is critical to its survival. In contrast, the *E. coli* OxyR regulon is not induced upon aeration, and upon aeration, *E. coli oxyR* mutants do not exhibit any survival defect. Since OxyR is activated by hydrogen peroxide, a plausible explanation is that *B. thetaiotaomicron* generates more H₂O₂ when oxygen enters its cytoplasm than does *E. coli*. To test this idea, we examined the rate of intracellular H₂O₂ formation by measuring the rate at which hydroperoxidase-deficient (Hpx⁻) mutants excrete H₂O₂ into the medium. *E. coli* employs two catalases and NADH peroxidase (AhpCF) to degrade hydrogen peroxide, and so *E. coli* Hpx⁻ mutants lack the genes coding for these three enzymes (*katG*, *katE*, and *ahpCF*) (30). *B. thetaiotaomicron* employs a catalase, NADH peroxidase, and two rubrerythrins, and *B. thetaiotaomicron* Hpx⁻ mutants are *katE ahpCF rbr1 rbr2* strains (17). Prior work suggested that Hpx⁻ *B. thetaiotaomicron* excreted H₂O₂ at high rates, although the rates were not quantified under the same conditions as for *E. coli*. In the present study, the Hpx⁻ derivatives of both bacteria were cultured in several anoxic complex and defined media, and the bacteria were then shifted to oxic buffer or medium prior to measurements of H₂O₂ production. Under identical circumstances, the rate of H₂O₂ formation was approximately 10-fold higher for the *B. thetaiotaomicron* strain than for the *E. coli* strain (Fig. 3B). The higher rate of H₂O₂ production offers a likely explanation for why OxyR is induced and H₂O₂-sensitive enzymes are inactivated when *B. thetaiotaomicron* is aerated, in contrast to *E. coli*.

The H₂O₂ that is formed in aerobic *E. coli* comes from a mixture of sources that have not all been identified. During constant aerobiosis, about one-third of endogenous H₂O₂ derives from aspartate oxidase (26), which uses molecular oxygen as an electron acceptor in the first step of nicotinamide biosynthesis. Because this reaction is saturated by moderate levels of oxygen, the rate of H₂O₂ formation by this enzyme has a ceiling that is unaffected by further increases in oxygen concentration. In contrast, other endogenous H₂O₂ is formed by unknown sources and is generated in proportion to oxygen concentration. This property suggests that this H₂O₂ is produced by the adventitious oxidation of redox enzymes by oxygen, therefore displaying chemical rather than Michaelis-Menten kinetics. Similarly, we observed that the high




FIG 3 Endogenous H₂O₂ formation is much more rapid in aerated *B. thetaiotaomicron* than in aerated *E. coli*. (A) Survival of wild-type and *oxyR* mutant strains after aeration. (B) Mutant strains lacking H₂O₂ scavenging enzymes (Hpx⁻) were precultured in the indicated anoxic media, and they were then resuspended in aerobic PBS-glucose or defined media, as described in Materials and Methods. Intracellular H₂O₂ production is inferred from the rate of H₂O₂ release into the medium. The star denotes that *B. thetaiotaomicron* did not grow in LB medium.




FIG 4 The rate of H₂O₂ production in *B. thetaiotaomicron* is proportionate to the ambient O₂ concentration. Log-phase *B. thetaiotaomicron* Hpx⁻ cells were resuspended in PBS-glucose at an OD₆₀₀ of 0.1. H₂O₂ production was monitored at different levels of dissolved O₂.

rate of H₂O₂ formation by *B. thetaiotaomicron* was proportionate to the concentration of dissolved oxygen (Fig. 4). The mechanistic implication is that H₂O₂ is produced by the accidental oxidation of an enzyme(s). In terms of organismal biology, it means that the oxidative stress experienced by the bacterium will be less severe in micro-oxic environments than in fully oxic ones.

The rate of intracellular H₂O₂ formation can be derived from the rate at which it accumulates in the growth medium, if one knows the ratio of cell volume to total culture volume (Materials and Methods). In this way, we deduce that if a concentration of cells representing an optical density at 600 nm (OD₆₀₀) of 0.1 causes H₂O₂ to accumulate in the medium at a rate of 0.24 µM/min H₂O₂ (Fig. 4), then the H₂O₂ is formed inside the cells at a rate of approximately 0.1 mM/s. Our previous studies (16) indicated that *B. thetaiotaomicron* consumes glucose at an intracellular rate of slightly more than 1 mM/s. In contrast, based upon protein content and growth rate, we calculate that the fluxes through amino acid biosynthetic pathways are only 10 µM/s; nucleotide synthesis and lipid synthesis are similar, and other pathways are substantially slower. The implication is that central metabolism stands out as having the only pathways with flux capacities compatible with the observed rate of ROS formation. This logic focused our attention upon fumarate reductase.

In vivo evidence does not indicate that fumarate reductase is a major H₂O₂ source. When *E. coli* moves from anoxic to oxic environments, the primary source of H₂O₂ is fumarate reductase (Frd) (26). This enzyme is synthesized only during anoxia, but the extant enzyme continues to operate—and releases ROS—when oxygen is infused into the medium. Frd provides a minor pathway of carbohydrate fermentation in *E. coli* (31), but it is central to the anaerobic metabolism of many bacteria, including *B. thetaiotaomicron* (Fig. 1B). We reasoned that *B. thetaiotaomicron* might contain especially high titers of Frd and that Frd might therefore be the source of the high H₂O₂ flux after aeration. The most straightforward test of that idea would be to quantify H₂O₂ production from a strain that lacked Frd. However, we were unable to recover any *frd* null mutants, despite several attempts and in contrast to our success with all other genes we attempted. We also observed that *B. thetaiotaomicron* grows extremely poorly when heme is omitted from the medium (data not shown). Heme is the source of the heme moiety in *B. thetaiotaomicron* Frd, and we suspect that the requirement for heme in growth medium reflects the importance of this enzyme to anaerobic metabolism. For these reasons, it was not possible to genetically test the hypothesis that Frd produces most of the intracellular H₂O₂.

As an alternative, we appraised the effect of fumarate supplementation upon H₂O₂ release. In principle, the *B. thetaiotaomicron* Frd is especially vulnerable to autoxidation because the fumarase of this bacterium loses activity upon aeration. The inactivity of fumarase blocks the formation of intracellular fumarate, leaving Frd without a natural electron acceptor (Fig. 1B). When other redox enzymes are deprived of their acceptors, they are much more prone to pass electrons to oxygen (27). Therefore, we provided fumarate as a supplement in the medium. Bacteria that employ fumarate as a terminal electron acceptor are usually able to import it, and indeed the *B. thetaiotaomicron* genome encodes two membrane proteins homologous to the *E. coli* DcuA and DcuB fumarate importers (32). Previous work confirmed that exogenous fumarate is imported and reduced by Frd (16). However, fumarate supplements did not diminish the rate of intracellular H₂O₂ production (see Fig. S1A in the supplemental material). This result did not support the notion that Frd is a major source of H₂O₂ in *B. thetaiotaomicron*.

The *B. thetaiotaomicron* Frd, unlike the *E. coli* enzyme, does not produce ROS in vitro. The previous result surprised us. We then examined the behavior of the Frd enzyme in inverted membrane vesicles. The vesicles were prepared by French press. The physiological reaction of Frd is to transfer electrons from quinones to fumarate (Fig. 1A), thereby generating succinate, but the reaction is reversible both *in vitro* and *in vivo*. We found that the succinate:quinone oxidoreductase activities were quantitatively similar in anoxically grown *E. coli* and *B. thetaiotaomicron*, indicating similar titers of Frd, contrary to our expectation (see Fig. S2 in the supplemental material). Assays of NADH:fumarate reductase activity confirmed that result (not shown).

When *E. coli* vesicles were incubated with succinate in oxic buffer, both profuse H₂O₂ and superoxide were formed (Fig. 5 and Fig. 6; Fig. S1B and C). As was reported before, the formation of this ROS depended upon the presence of Frd (see Table S1). In surprising contrast, despite exhibiting ample succinate:quinone reductase activity, the succinate-treated *B. thetaiotaomicron* vesicles did not generate any significant amount of either oxidant (Fig. 5 and 6). The *E. coli* enzyme characteristically produced less superoxide at higher doses of succinate, since the more-reduced enzyme releases electrons to oxygen in pairs rather than singly; at very high doses, succinate prevents all ROS formation by occluding the flavin. The *B. thetaiotaomicron* enzyme did not release either species at any concentration of succinate (Fig. S1B and C).

The contrasting autoxidation behaviors of the two enzymes were also apparent from measurements of oxygen consumption. When *E. coli* membranes were incubated with succinate, oxygen was consumed even if cytochrome oxidase was blocked by cyanide (Fig. 7). The rate of oxygen consumption matched the rate of H₂O₂ release. Yet the *B. thetaiotaomicron* membranes did not consume oxygen at all (<8% the rate of the *E. coli* membranes).

In a prior publication (26), we proposed that obligate anaerobes might minimize ROS formation by Frd through the actions of cytochrome *bd* oxidase. When oxygen is present, this oxidase competes with Frd for reduced quinones and thereby diminishes ROS production by the *E. coli* enzyme. However, this arrangement does not explain the lack of O₂⁻ and H₂O₂ deriving from the *B. thetaiotaomicron* Frd, as virtually no oxygen consumption occurred even when cyanide was not provided to inhibit the oxidase (Fig. 7). In separate experiments, we confirmed that *B. thetaiotaomicron* consumes oxygen upon aeration, but the rate is undimin-




FIG 5 In membrane vesicles, the *E. coli* Frd produces H_2O_2 , but the *B. thetaiotaomicron* Frd does not. (A) Representative time course during incubation with 0.4 mM succinate. (B) Rates of H_2O_2 production at 0.4 mM succinate. pr., protein. The fumarate reductase activities (succinate:plumbagin oxidoreductase activities) of the membranes were approximately equivalent (Fig. S2).

ished in cytochrome *bd* oxidase mutants. Instead, rubredoxin:oxygen oxidoreductase (Roo), a soluble enzyme, is responsible (see Table S2 in the supplemental material). We infer that under our anoxic culturing conditions, the cytochrome *bd* oxidase was not expressed and therefore had no effect on Frd behavior.

During normal metabolism, electrons flow from NADH through the menaquinone pool to Frd (Fig. 1A). This electron transfer pathway is active in inverted vesicles (see Fig. S3A in the supplemental material). The addition of cyanide to *E. coli* vesicles diminished the rate of NADH oxidation by inhibiting the cytochrome oxidases, but it did not limit NADH oxidation entirely, because adventitious electron transfer from reduced chain components to oxygen persisted. Much of this residual flux was due to Frd autoxidation, as it was blocked by malonate, a potent competitive inhibitor that binds opposite the flavin and impedes the approach of oxygen. With *B. thetaiotaomicron* vesicles, only a modest amount of oxygen consumption was detected, and it was resistant to both cyanide and malonate. This NADH consumption was likely due to the autoxidation of NADH dehydrogenase (4). The data indicate that Frd does not autoxidize even when electrons arrive through the quinone pool. That conclusion was further supported by assays of H_2O_2 production (Fig. S3B). Importantly, these data indicate not only that Frd is not the source of the outsized H_2O_2 formation in *B. thetaiotaomicron*, but also that other components of the electron transport chain are not the source, either.

The previous study of *B. fragilis* showed that a mutation that

knocked out *frdC* diminished the rate of H_2O_2 release from whole cells by about 40%; an inference was drawn that Frd might be the source of the lost H_2O_2 . However, *B. fragilis* *frd* mutants grow at only 30% the rate of their wild-type parent (33), which raised the alternative possibility that slow metabolism was the cause of the drop. To look more directly, we prepared membranes from the same *B. fragilis* strain and tested the ability of its Frd to generate superoxide. Like the *B. thetaiotaomicron* enzyme, with which it shares ca. 90% identity, the *B. fragilis* Frd did not generate any detectable superoxide (Fig. 6; Fig.S1B).

The identity of the iron-sulfur subunit determines the rates of Sdh/Frd autoxidation. A key structural distinction between the Frd enzymes of *E. coli* and *B. thetaiotaomicron* is that the latter enzyme is among the members of the complex II family (34) that have two heme binding sites in the membrane anchor subunit FrdC (Fig. 1; see Fig. S4 in the supplemental material). In that respect it resembles *E. coli* Sdh, which also includes a heme moiety (35). In contrast, the *E. coli* Frd instead employs two smaller trans-membrane subunits that lack heme-binding ligands and do not incorporate heme as a cofactor (36). Redox calculations led Yankovskaya et al. to propose that the heme cofactor of Sdh would diminish electron occupancy of its flavin cofactor, which might diminish ROS formation (35). Indeed, Sdh generates far less H_2O_2 and O_2^- than does Frd (25).

To test the significance of this difference, the *frdCAB* operon of *B. thetaiotaomicron* was expressed in a $\Delta frd \Delta sdh$ mutant strain of *E. coli*. The vesicles exhibited robust succinate:quinone oxi-




FIG 6 The *E. coli* Frd produces O_2^- , but the *B. thetaiotaomicron* Frd does not. (A) Representative time course during incubation of vesicles with 0.4 mM succinate. pr., protein. (B) Rates of O_2^- production at 0.4 mM succinate. Membranes from a *B. thetaiotaomicron* SOD⁻ mutant also produced no O_2^- , confirming that the lack of O_2^- detection was not due to contaminating SOD.




FIG 7 Cyanide-resistant O₂ consumption by inverted vesicles confirms that *E. coli* Frd produces ROS but *B. thetaiootaomicron* Frd does not. Reaction mixtures contained 0.4 mM succinate as the electron donor and 3 mM KCN to block oxygen consumption by cytochrome oxidase. Where indicated, 3 mM malonate was added to block the access of oxygen to the flavin of Frd.

doreductase activity, but again the *B. thetaiootaomicron* Frd did not generate either O₂⁻ or H₂O₂ (Fig. 8A). Moreover, the expression of *E. coli* Frd in Hpx⁻ cells caused substantial excretion of H₂O₂ when the cells were aerated, but expression of the *B. thetaiootaomicron* enzyme did not (Fig. 8B). These results confirmed that its nonautoxidizability is a trait of the *B. thetaiootaomicron* Frd itself rather than of the *B. thetaiootaomicron* lipid or quinone environment. To dissect the enzyme further, we attempted to create chimeric enzymes that contained mixtures of the subunits from the two bacterial Frd enzymes. Only the combination of the *E. coli* FrdA subunit and the *B. thetaiootaomicron* FrdB and -C subunits provided active enzyme (see Fig. S5 in the supplemental material). This hybrid enzyme also did not autoxidize (Fig. 8A). Thus, the suppression of Frd autoxidation behavior is not due to differences in the FrdA subunit, which is the actual site of electron transfer to oxygen, but rather to the effects of the FrdB/C subunits upon it. These data fit the model (25, 35) that electron withdrawal by the iron-sulfur cluster and/or heme moieties can minimize autoxidation from the flavin adenine dinucleotide (FAD) site.

To specifically test the impact of the heme moieties upon Frd autoxidation, it would be ideal to eliminate the heme binding sites of the *B. thetaiootaomicron* C subunit. However, we were unable to recover active enzyme from *E. coli* strains expressing such a mutant construct (Fig. S5). It is likely that the hemeless C subunit was unstable. As an alternative, we tested the impact of the heme upon the autoxidation rate of *E. coli* Sdh. Tran et al. generated such a mutant, and in their own studies, they did not observe any increase in Sdh oxidation during succinate respiration (37, 38). To examine this issue more closely, we expressed both the wild-type and heme-deficient enzymes in a quinoneless strain of *E. coli*, thereby ensuring that the rate of electron flow to the quinone pool did not have any impact upon autoxidation. Quite clearly, the removal of heme did not convert Sdh to high ROS production in the fashion of Frd (Fig. S5E). We think this result can likely be extrapolated to *B. thetaiootaomicron* Frd. Although that enzyme additionally contains a distal heme that Sdh lacks, the distal hemes in Frd di-heme enzymes typically lie at low potentials (34) and should have minimal impact as electron sinks. Collectively, these data indicate that the pace of flavin autoxidation in this enzyme family is determined by the nature of the iron-sulfur subunit rather than by either the structure of the flavoprotein subunit or the presence of heme cofactors.




FIG 8 The absence of ROS production by *B. thetaiootaomicron* Frd does not derive from the lipid composition or the nature of its flavoprotein subunit. (A) Cell membranes were prepared from *E. coli* Hpx⁻ cells, from cells of the *E. coli* Hpx⁻ Δfrd mutant carrying a plasmid expressing *B. thetaiootaomicron* wild-type Frd (*frdCBA*), and from cells of the *E. coli* Hpx⁻ Δfrd mutant expressing the chimeric Frd (*E. coli* *frdA* plus *B. thetaiootaomicron* *frdBC*). The rates of *in vitro* O₂⁻ and H₂O₂ production are normalized to the succinate:plumbagin oxidoreductase activities. Reaction mixtures contained 0.4 mM succinate; 3 mM KCN was included in measurements of O₂⁻ formation. (B) Only *E. coli* Frd generates H₂O₂ *in vivo*. The *E. coli* Hpx⁻ ΔfrdABCD strain LC126 was transformed with empty vector, plasmid pfrd(CAB)_{Bt} expressing *B. thetaiootaomicron* Frd, or plasmid pH3 expressing *E. coli* Frd. Fumarate reductase was induced in anoxic lactose-Casamino Acids medium, and H₂O₂ excretion was then quantified after dilution into oxalic glucose buffer (see Materials and Methods).

What is the primary source of ROS in *B. thetaiootaomicron*?

Thus, *B. thetaiootaomicron* generates high levels of intracellular oxidants when it encounters oxygen, but the source is neither Frd nor any of the other components of the membrane-bound respiratory chain. One alternative was the series of redox enzymes that deliver electrons to Roo, the soluble oxygen reductase. Those electrons flow from NADH through rubredoxin to Roo (17, 39, 40). When rubredoxin was deleted from the non-scavenging strain, those mutants (*rd*) generated 25% less H₂O₂ than did the rubredoxin-proficient strain (see Table S3 in the supplemental material). NADH:rubredoxin oxidoreductase (NROR) is one source of those electrons, but not the sole one. Both NROR- and Roo-deficient mutants exhibited wild-type rates of H₂O₂ formation. Thus, the predominant sources of ROS remain unknown.

DISCUSSION

Rapid ROS formation precludes aerobic growth but is not lethal. Dosimetric studies showed that *E. coli* expresses just enough scavenging activity to avoid being poisoned by its endogenous O₂⁻ and H₂O₂ (12). When the steady-state levels of these oxidants were elevated more than 3-fold, its oxidant-sensitive enzymes lost activity, and growth slowed. Against this backdrop, it is striking that aeration drives *Bacteroides thetaiootaomicron* to generate 10-fold more ROS than does *E. coli*. Further, *B. thetaiootaomicron* does not compensate with a proportionate increase in the titers of its scavenging enzymes; in fact, we determined that its SOD levels (2 U/mg) are much lower than those in either aerobic (16 U/mg) or even anaerobic (5 U/mg) *E. coli* cells (see Fig. S6 in the supplemental material). In conjunction with the higher rate of ROS formation, the implication is that the steady-state level of superoxide in fully aerated *B. thetaiootaomicron* may exceed that of *E. coli* by >25-fold. This situation provides a good explanation for why *B. thetaiootaomicron* aeration is followed by the progressive loss of fumarase activity, the collapse of the succinate/propionate branch of its fermentation, and the cessation of growth (16).

This vulnerability is one factor that constrains *B. thetaiootaomicron* to life in anoxic environments. Nevertheless, the bacterium

tolerates aeration well enough to move between hosts. Furthermore, if the human gut is perforated by physical trauma, *Bacteroides* species are able to spread into the erstwhile oxic peritoneum and form life-threatening abscesses (41). These events manifest the theme that all anaerobes must withstand occasional oxygen exposure. How they manage to do so is a focus of substantial research activity.

Several features sustain *B. thetaiotaomicron* when it enters oxic environments. The fact that ROS formation is proportionate to oxygen level (Fig. 4) has the consequence that stress is lessened in tissues, where oxygen levels are at least 5-fold lower than in air-saturated fluids. Second, while the ROS can inactivate enzymes and impair regular metabolism, they do not generate life-threatening DNA lesions (Fig. 3A), because the OxyR-mediated induction of iron storage proteins defuses the possibility of Fenton chemistry (42). Finally, by using its respiration-linked cytochrome *bd* oxidase and cytoplasmic rubredoxin: oxygen oxidoreductase, *B. thetaiotaomicron* presumably can gradually clear oxygen from the local microenvironment, allowing the bacterium to repair its damaged enzymes and to restore its full metabolic capacity. Nevertheless, ROS production places a limit upon the oxygen level that allows the resumption of metabolism. We were surprised to discover that the fumarate reductase of *B. thetaiotaomicron*, which we and others anticipated would be the major source of ROS, has instead acquired a structure that avoids any detectable ROS formation at all.

Why does not *B. thetaiotaomicron* fumarate reductase generate ROS? The complex II family of enzymes—aspartate oxidase, fumarate reductase, and succinate dehydrogenase—has come under especially close study because the autoxidation of these enzymes has important physiological consequences. Previous studies suggested a model for how their redox structures create differences in their autoxidation behaviors. Aspartate oxidase is the simplest member. It is comprised of a single flavoprotein subunit (43), whose flavin is alternately reduced by aspartate and oxidized (in anoxic cells) by fumarate. Hence it is a fumarate reductase. However, in aerobic cells, the fumarate level falls too low, and oxygen itself receives the electrons with stoichiometric conversion to H_2O_2 (26). Kinetic analysis showed that the oxidation step is not saturated by O_2 , which reveals that the enzyme has no authentic O_2 binding site. Thus, aspartate oxidase really is an aspartate:fumarate oxidoreductase whose turnover is sustained in aerobic cells by the chemical oxidizability of its exposed flavin. This switch in electron acceptor is unique and remarkable.

The flavoprotein subunit of respiratory fumarate reductase is tethered to the cell membrane by the FrdB iron-sulfur wire and the FrdC/D integral membrane subunits (Fig. 1A). The latter also comprise a quinone binding site. In anaerobic cells, electrons arrive from the quinone pool of the respiratory chain, flow through the iron-sulfur centers to the flavin, and again are transferred to fumarate as the physiological acceptor. However, when *E. coli* cells transit to an oxic environment, oxygen can again accept electrons from the reduced flavin. Oxygen is a diradical, so the electron transfer necessarily occurs through consecutive single-electron events (44). When Frd is oxidized, the predominant product is O_2^- rather than H_2O_2 (25), because after transfer of the first electron from FADH_2 , the second electron can be sequestered on a nearby iron-sulfur cluster. The initial O_2^- diffuses out of the active site, and the second electron is ultimately transferred to oxygen in a second oxidation event. The O_2^- that Frd forms is more

hazardous than the H_2O_2 that aspartate oxidase forms, because O_2^- damages vulnerable enzymes more quickly.

The failure of *B. thetaiotaomicron* Frd to autoxidize like *E. coli* Frd is reminiscent of the behavior of *E. coli* Sdh. Sdh shares the same flavin-iron-sulfur linkage as Frd, but its clusters I and III sit at a higher potentials (+10 mV and +65 mV [45, 46]) than those of Frd (−35 mV and −65 mV [47, 48]) and in the divalently reduced enzyme are expected to pull the electron density away from the flavin. A much slower pace of flavin oxidation is the expected result. Yandovskaya et al. suggested that this effect might be further enhanced by the Sdh heme (35), but data from Tran et al. (37, 38) and from Fig. 8 show that this is not necessary.

Against this backdrop, a reasonable explanation for the non-oxidizability of *B. thetaiotaomicron* Frd is that it, like Sdh, may have higher-potential clusters that suppress flavin autoxidation. Measurements will be necessary to confirm the idea; other explanations are possible. However, the fact that *B. thetaiotaomicron* FrdB/C subunits suppressed the autoxidation of *E. coli* FrdA certainly fits the model.

What is new about this observation is that, unlike *E. coli* Sdh, the *B. thetaiotaomicron* Frd retains its role as a fumarate reductase. That is, the suspected shift of electron density away from its flavin does not hinder physiological electron flow to the fumarate site. Although at first blush this outcome might seem surprising, further consideration of redox dynamics would argue that electron distribution on an enzyme does not need to correlate rigidly with the direction of its flow. Electron exchange between the redox moieties in all of these enzymes is certainly extremely fast relative to the lifetime of either enzyme-substrate complex; thus it is unlikely that a shift in electron density will create a bottleneck in the catalytic cycle, which involves the much slower steps of substrate binding and product release (49). The same idea can be invoked to explain why catalysis in either direction would be unimpeded by the requirement that electrons flow across the very-low-potential cluster II (−250 to −300 mV) (50). Yet the redistribution of electron density from one side of the enzyme to the other will strongly affect the pace of electron transfer to oxygen, since the collision of oxygen with the flavin is a momentary elastic event.

There is an interesting parallel between *B. thetaiotaomicron* Frd and class I ribonucleotide reductases, which evolved to reduce ribonucleotides in the aerobic world. Ribonucleotide reductases employ a cysteinyl radical to abstract an electron from ribonucleotides. Such radicals are extremely vulnerable to oxidation by molecular oxygen—but in the class I enzymes, this calamity is avoided because the radical is predominantly localized on an electronically linked tyrosine residue, which is buried in the protein and thereby shielded from oxygen. When substrate binds, resonance relocates the radical to the active-site cysteine at a rate sufficient for robust turnover. Similarly, in *B. thetaiotaomicron* Frd—and *E. coli* Sdh—the localization of the electron pair on iron-sulfur clusters away from the exposed flavin may be an adaptation that circumvents inappropriate autoxidation.

Interestingly, some complex I (NADH dehydrogenase) isozymes exhibit an extra redox moiety that, like the heme of Sdh, had been proposed to suppress ROS formation by electron withdrawal. NADH dehydrogenase uses a long iron-sulfur wire to move electrons from the solvent-exposed flavin to quinone-reduction site, but an additional cluster (N1a) lies adjacent to the flavin but outside of the wire itself. It was hypothesized that during turnover, the extra cluster might momentarily sequester an elec-

tron from the flavin, minimizing its flavosemiquinone content and thereby diminishing ROS production (51, 52). However, conversion of the N1a cluster to a nonreducible low-potential center did not alter the autoxidation rate of the isolated enzyme (53). Thus, for the moment the disparity in autoxidation behaviors of *B. thetaiotaomicron* and *E. coli* Frd enzymes is a singular demonstration that some redox enzymes can acquire structures that suppress their tendency to contribute to oxidative stress. This is a logical complement to the evolutionary appearance in some organisms of oxidant-resistant enzymes, such as special isozymes of fumarase (54), dihydroxyacid dehydratase (55), hydrogenase (56), and pyruvate:ferredoxin oxidoreductase (57).

Finally, we note that the identity of the residual ROS source in *B. thetaiotaomicron* remains unknown. The *in vitro* data indicated that none of the components of the anaerobic respiratory circuit is likely to be responsible, as ROS leakage from the chain *in vitro* was not markedly different from that from the *E. coli* chain. Prior analyses have shown that, aside from Frd, the *E. coli* chain is not a major ROS contributor to that bacterium (58). In *B. thetaiotaomicron*, the soluble rubredoxin-dependent electron chain seems to be a minor source. An attractive candidate for the residual ROS might be the ferredoxin-mediated flow of electrons from pyruvate to hydrogenase, since this high-volume pathway involves a series of metal centers that operate at low potential near enzyme surfaces. It will be important to identify the main *B. thetaiotaomicron* ROS source, since it is likely responsible for the deactivation of fumarase upon aeration and in part for the consignment of the bacterium to hypoxic and anoxic habitats.

MATERIALS AND METHODS

Chemicals. Most chemicals were purchased from Sigma. Amplex Ultra-Red was bought from Invitrogen. Horseradish peroxidase (HP), horse heart cytochrome *c*, and *E. coli* iron-containing superoxide dismutase were all Sigma products.

Cell growth and media. Anaerobic cultures were grown at 37°C in an anaerobic glove box (Coy Laboratory Products) containing 85% N₂, 10% H₂, and 5% CO₂. Defined medium for *E. coli* contained minimal A salts (59), 0.2% glucose, 0.2% Casamino Acids, 0.5 mM tryptophan, and 5 μg/ml thiamine. BHIS medium and defined glucose medium for *B. thetaiotaomicron* were made as described previously (60). *B. fragilis* was also grown in BHIS medium. Media for anaerobic cultures were autoclaved and then moved into the anaerobic chamber and degassed for at least 24 h before use. Plasmids were transformed by the CaCl₂ method into *E. coli* Hpx⁻ (LC106 or LC126) or *ubiA menA* (KM8) strains inside the anaerobic chamber. To ensure that quinone mutants had not reverted during cell growth for the preparation of membranes, some harvested cells were streaked onto aerobic LB medium to confirm that the colonies remained tiny after 2 days of growth.

Cell growth was monitored by measurement of OD₆₀₀. Prior to aeration of previously anaerobic cultures, log-phase cells were grown anoxically from an OD₆₀₀ of 0.005 to an OD₆₀₀ of approximately 0.2. Cells were centrifuged, and the pellets were then resuspended in warm aerobic medium. The viability of aerated *B. thetaiotaomicron* was tracked by transferring cells back to the anaerobic chamber and plating them on anoxic BHIS plates. Colonies were counted after 3 days.

The strains and plasmids used in this study are listed in Table S4 in the supplemental material. The *B. thetaiotaomicron* strains were derived from BT5482 Δtdk , as described previously (61). The deletion of genes was performed by standard methods (61), and deletions were confirmed by sequencing or genome PCR. A low-copy-number plasmid (pWKS30) was used to construct plasmids that express chimeric fumarate reductase enzymes in *E. coli*. Genes encoding the three subunits were cloned behind the *lac* promoter. The ribosome binding site (RBS) of *E. coli gapA* was inserted

upstream of each gene. Construction was confirmed by digestion and/or DNA sequencing.

Plasmid pfrd(CAB)_{B1} was used as the template for site-directed mutagenesis of *B. thetaiotaomicron* frdC (H178L, H178Q, or H178Y). After mutagenesis PCR, DpnI was used to digest the template plasmid for 1 h at 37°C. Oligonucleotides were synthesized by Eurofins MWG Operon (United States). Mutant alleles were confirmed by sequencing (ACGT, Inc., USA).

The plasmids were transformed into the *E. coli* $\Delta sdh \Delta frd$ strain (KM7) or the *E. coli* Hpx⁻ Δfrd strain (LC126). Cells were grown in minimal medium containing lactose as the sole carbon source and inducer before membrane vesicles were prepared, as described below.

Preparation of inverted membrane vesicles. Membrane vesicles were prepared by standard methods (27). Cells were grown under anaerobic conditions from an A₆₀₀ of 0.005 to 0.2 to 0.3 and then were washed in cold potassium phosphate buffer (50 mM [pH 7.8]), suspended in the same buffer at 1% of the original volume, and lysed by French press. Cell debris was removed by centrifugation (4°C, 20,000 × *g* for 15 min), and the supernatant was then diluted 5-fold in buffer and centrifuged in an ultracentrifuge (4°C, 100,000 × *g* for 1.5 h). The supernatant was discarded, and the resulting pellet was resuspended in the same potassium phosphate buffer. The ultracentrifugation step was then repeated, and the vesicles were finally suspended in ~1% of the original culture volume in potassium phosphate buffer. Vesicles were stored on ice. Protein concentrations were measured using the Coomassie blue reagent from Thermo-Fisher.

Measurements of oxygen consumption. Oxygen consumption by respiring vesicles was measured using a Clark oxygen electrode. Vesicles were added to KP₁ buffer that had been prewarmed in a 37°C water bath. Succinate (0.4 mM) and KCN (3 mM) were added to detect oxygen consumption that was independent of electron flow through cytochrome oxidase; this rate represented production of O₂⁻ and H₂O₂ by adventitious electron transfer to oxygen upstream in the respiratory chain. Where indicated, 3 mM malonate was added as a specific inhibitor of fumarate reductase. Measurements were performed at 37°C.

Subunit A of fumarate reductase was detected after electrophoresis of membrane samples containing 275 μg protein on SDS-polyacrylamide gels (62). The gels were exposed to UV transillumination, and the UV fluorescence of the bound flavin was quantified by a Quantity One system (Bio-Rad).

Measurements of H₂O₂ formation. The formation of H₂O₂ by aerated cells was measured. Hpx⁻ cells that had been cultured into log phase in anaerobic rich medium were harvested by centrifugation, washed with phosphate-buffered saline (PBS) buffer (pH 7.2), and resuspended to an OD₆₀₀ of 0.1 in warm PBS buffer containing 0.05% glucose. The OD₆₀₀ was around 0.1. Cells that had been grown in anaerobic basic media were resuspended in minimal A medium with 0.05% glucose and 0.01% Casamino Acids or in defined medium with glucose but without hemin and cysteine. Hemin and cysteine can interfere with H₂O₂ formation. The resuspended cells were shaken at 37°C under room air. At intervals, samples were removed, and their H₂O₂ content was determined by Amplex Red-horseradish peroxidase (HRP) analysis.

The dependence of H₂O₂ formation upon solution oxygen content was determined in a similar way. *B. thetaiotaomicron* Hpx⁻ cells were cultured from an OD₆₀₀ of 0.01 to 0.25 in anoxic BHIS medium. Cells were washed and resuspended to an OD₆₀₀ of 0.1 in anoxic PBS (pH 7.2) buffer containing 0.2% glucose; the medium had been presaturated by a gas mixture of N₂-air and maintained in a 37°C water bath. At intervals, samples were removed, and H₂O₂ was quantified by the Amplex Red-HRP method. The gas mixture was established by mixing gas flow from nitrogen and oxygen cylinders at a Y intersection; the gas stream then was bubbled through a water trap (to ensure hydration) and finally through the cell culture.

The rate of H₂O₂ excretion by cells can be used to derive the rate of intracellular H₂O₂ production. We previously determined that 1 ml of

E. coli cells at an OD₆₀₀ of 1 comprises 0.5 μM cytoplasm (63); therefore, for the cultures at an OD₆₀₀ of 0.1 used in H₂O₂ measurements, the ratio of culture volume to cytoplasmic volume is 20,000:1. Hence an excretion rate of 0.24 μM H₂O₂/min into the medium (Fig. 4) represents an intracellular production rate of 4.8 mM/min, or 80 μM/s. We make the assumption that the relationships between OD₆₀₀ and cell volume are similar between *E. coli* and *B. thetaiotaomicron*; this notion is supported by the fact that we recover similar amounts of protein per OD unit.

The production of H₂O₂ was also measured during the respiration of inverted vesicles that had been prepared from *E. coli* Hpx⁻ or *B. thetaiotaomicron* Hpx⁻ cells. The cells had been grown in anaerobic minimal glucose media. Vesicles were added to 20 ml aerobic potassium phosphate buffer (50 mM [pH 7.8]) containing 0.4 mM succinate or 40 μM NADH. Superoxide dismutase (100 U/ml) was included in the reaction mixture. Where indicated, 3 mM malonate was used as an inhibitor of fumarate reductase. The reaction mixtures were shaken at 37°C, samples were removed at intervals, and their H₂O₂ content was measured by the Amplex Red-HRP method (64).

Measurements of O₂⁻ formation and ferricyanide and plumbagin reduction by inverted membrane vesicles. Superoxide formation by inverted vesicles was measured in 1 ml KP_i buffer (pH 7.8) with succinate or NADH as the electron donor. The reduction of cytochrome *c* was monitored at 550 nm ($\epsilon = 21.0 \text{ mM}^{-1} \text{ cm}^{-1}$), in the presence and absence of SOD (27). The SOD-resistant activity represents direct reduction of cytochrome *c* by membrane enzymes, and this rate was subtracted from the no-SOD rate to derive the O₂⁻ rate. Where indicated, 3 mM KCN was included to block possible electron flux from fumarate reductase through the cytochrome oxidase. Plumbagin reduction leads to immediate superoxide formation, so cytochrome *c* reduction was again monitored, in the presence of 3.2 mM succinate, 3 mM KCN, and 0.4 mM plumbagin (27). Ferricyanide reduction was monitored in the presence of respiratory substrate plus 0.2 mM potassium ferricyanide and 3 mM KCN; the reduction of ferricyanide was tracked by the disappearance of its absorbance at 420 nM (1 mM⁻¹ cm⁻¹).

SOD activities were measured by the xanthine oxidase-cytochrome *c* method (65).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://mbio.asm.org/lookup/suppl/doi:10.1128/mBio.01873-16/-/DCSupplemental>.

Figure S1, TIF file, 2.7 MB.

Figure S2, TIF file, 1.4 MB.

Figure S3, TIF file, 1.7 MB.

Figure S4, TIF file, 2.6 MB.

Figure S5, TIF file, 2.8 MB.

Figure S6, TIF file, 1.9 MB.

Table S1, DOC file, 0.1 MB.

Table S2, DOC file, 0.1 MB.

Table S3, DOC file, 0.1 MB.

Table S4, DOC file, 0.1 MB.

ACKNOWLEDGMENTS

We thank Gary Cecchini and Jeff Smith for generously providing plasmid constructs and strains that were used in this study.

This investigation was funded by grant GM49640 from the National Institutes of Health. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

FUNDING INFORMATION

This work, including the efforts of James Imlay, was funded by HHS | National Institutes of Health (NIH) (GM49640).

REFERENCES

- Anbar AD. 2008. Elements and evolution. *Science* 322:1481–1483. <http://dx.doi.org/10.1126/science.1163100>.
- Wagner AF, Frey M, Neugebauer FA, Schäfer W, Knappe J. 1992. The

- free radical in pyruvate formate-lyase is located on glycine-734. *Proc Natl Acad Sci U S A* 89:996–1000. <http://dx.doi.org/10.1073/pnas.89.3.996>.
- Pieulle L, Magro V, Hatchikian EC. 1997. Isolation and analysis of the gene encoding the pyruvate-ferredoxin oxidoreductase of *Desulfovibrio africanus*, production of the recombinant enzyme in *Escherichia coli*, and effect of carboxy-terminal deletions on its stability. *J Bacteriol* 179:5684–5692. <http://dx.doi.org/10.1128/jb.179.18.5684-5692.1997>.
- Messner KR, Imlay JA. 1999. The identification of primary sites of superoxide and hydrogen peroxide formation in the aerobic respiratory chain and sulfite reductase complex of *Escherichia coli*. *J Biol Chem* 274:10119–10128. <http://dx.doi.org/10.1074/jbc.274.15.10119>.
- Gardner PR, Fridovich I. 1991. Superoxide sensitivity of the *Escherichia coli* 6-phosphogluconate dehydratase. *J Biol Chem* 266:1478–1483.
- Kuo CF, Mashino T, Fridovich I. 1987. α,β -Dihydroxyisovalerate dehydratase: a superoxide-sensitive enzyme. *J Biol Chem* 262:4724–4727.
- Flint DH, Tuminello JF, Emptage MH. 1993. The inactivation of Fe-S cluster containing hydro-lyases by superoxide. *J Biol Chem* 268:22369–22376.
- Jang S, Imlay JA. 2007. Micromolar intracellular hydrogen peroxide disrupts metabolism by damaging iron-sulfur enzymes. *J Biol Chem* 282:929–937. <http://dx.doi.org/10.1074/jbc.M607646200>.
- Sobota JM, Imlay JA. 2011. Iron enzyme ribulose-5-phosphate 3-epimerase in *Escherichia coli* is rapidly damaged by hydrogen peroxide but can be protected by manganese. *Proc Natl Acad Sci U S A* 108:5402–5407. <http://dx.doi.org/10.1073/pnas.1100410108>.
- Anjem A, Imlay JA. 2012. Mononuclear iron enzymes are primary targets of hydrogen peroxide stress. *J Biol Chem* 287:15544–15556. <http://dx.doi.org/10.1074/jbc.M111.330365>.
- Gu M, Imlay JA. 2013. Superoxide poisons mononuclear iron enzymes by causing mismetallation. *Mol Microbiol* 89:123–134. <http://dx.doi.org/10.1111/mmi.12263>.
- Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nat Rev Microbiol* 11:443–454. <http://dx.doi.org/10.1038/nrmicro3032>.
- Carlioz A, Touati D. 1986. Isolation of superoxide dismutase mutants in *Escherichia coli*: is superoxide dismutase necessary for aerobic life? *EMBO J* 5:623–630.
- Sobota JM, Gu M, Imlay JA. 2014. Intracellular hydrogen peroxide and superoxide poison 3-deoxy-D-arabinoheptulosonate 7-phosphate synthase, the first committed enzyme in the aromatic biosynthetic pathway of *Escherichia coli*. *J Bacteriol* 196:1980–1991. <http://dx.doi.org/10.1128/JB.01573-14>.
- Xu J, Bjursell MK, Himrod J, Deng S, Carmichael LK, Chiang HC, Hooper LV, Gordon JI. 2003. A genomic view of the human-*Bacteroides thetaiotaomicron* symbiosis. *Science* 299:2074–2076. <http://dx.doi.org/10.1126/science.1080029>.
- Pan N, Imlay JA. 2001. How does oxygen inhibit central metabolism in the obligate anaerobe *Bacteroides thetaiotaomicron*? *Mol Microbiol* 39:1562–1571. <http://dx.doi.org/10.1046/j.1365-2958.2001.02343.x>.
- Mishra S, Imlay JA. 2013. An anaerobic bacterium, *Bacteroides thetaiotaomicron*, uses a consortium of enzymes to scavenge hydrogen peroxide. *Mol Microbiol* 90:1356–1371. <http://dx.doi.org/10.1111/mmi.12438>.
- Massey V, Strickland S, Mayhew SG, Howell LG, Engel PC, Matthews RG, Schuman M, Sullivan PA. 1969. The production of superoxide anion radicals in the reaction of reduced flavins and flavoproteins with molecular oxygen. *Biochem Biophys Res Commun* 36:891–897. [http://dx.doi.org/10.1016/0006-291X\(69\)90287-3](http://dx.doi.org/10.1016/0006-291X(69)90287-3).
- Fridovich I. 1970. Quantitative aspects of the production of superoxide anion radical by milk xanthine oxidase. *J Biol Chem* 245:4053–4057.
- Ralph SJ, Moreno-Sánchez R, Neuzil J, Rodríguez-Enríquez S. 2011. Inhibitors of succinate:quinone reductase/complex II regulate production of mitochondrial reactive oxygen species and protect normal cells from ischemic damage but induce specific cancer cell death. *Pharm Res* 28:2695–2730. <http://dx.doi.org/10.1007/s11095-011-0566-7>.
- Ishii T, Miyazawa M, Onouchi H, Yasuda K, Hartman PS, Ishii N. 2013. Model animals for the study of oxidative stress from complex II. *Biochim Biophys Acta* 1827:588–597. <http://dx.doi.org/10.1016/j.bbabi.2012.10.016>.
- Quinlan CL, Orr AL, Perevoshchikova IV, Treberg JR, Ackrell BA, Brand MD. 2012. Mitochondrial complex II can generate reactive oxygen species at high rates in both the forward and reverse reactions. *J Biol Chem* 287:27255–27264. <http://dx.doi.org/10.1074/jbc.M112.374629>.

23. Perevoshchikova IV, Quinlan CL, Orr AL, Gerencser AA, Brand MD. 2013. Sites of superoxide and hydrogen peroxide production during fatty acid oxidation in rat skeletal muscle mitochondria. *Free Radic Biol Med* 61:298–309. <http://dx.doi.org/10.1016/j.freeradbiomed.2013.04.006>.
24. Dröse S. 2013. Differential effects of complex II on mitochondrial ROS production and their relation to cardioprotective pre- and postconditioning. *Biochim Biophys Acta* 1827:578–587. <http://dx.doi.org/10.1016/j.bbabi.2013.01.004>.
25. Messner KR, Imlay JA. 2002. Mechanism of superoxide and hydrogen peroxide formation by fumarate reductase, succinate dehydrogenase, and aspartate oxidase. *J Biol Chem* 277:42563–42571. <http://dx.doi.org/10.1074/jbc.M204958200>.
26. Korshunov S, Imlay JA. 2010. Two sources of endogenous hydrogen peroxide in *Escherichia coli*. *Mol Microbiol* 75:1389–1401. <http://dx.doi.org/10.1111/j.1365-2958.2010.07059.x>.
27. Imlay JA. 1995. A metabolic enzyme that rapidly produces superoxide, fumarate reductase of *Escherichia coli*. *J Biol Chem* 270:19767–19777.
28. Meehan BM, Malamy MH. 2012. Fumarate reductase is a major contributor to the generation of reactive oxygen species in the anaerobe *Bacteroides fragilis*. *Microbiology* 158:539–546. <http://dx.doi.org/10.1099/mic.0.054403-0>.
29. Rocha ER, Owens G, Jr, Smith CJ. 2000. The redox-sensitive transcriptional activator OxyR regulates the peroxide response regulon in the obligate anaerobe *Bacteroides fragilis*. *J Bacteriol* 182:5059–5069. <http://dx.doi.org/10.1128/JB.182.18.5059-5069.2000>.
30. Seaver LC, Imlay JA. 2001. Alkyl hydroperoxide reductase is the primary scavenger of endogenous hydrogen peroxide in *Escherichia coli*. *J Bacteriol* 183:7173–7181. <http://dx.doi.org/10.1128/JB.183.24.7173-7181.2001>.
31. Blackwood AC, Ledingham GA, Neish AC. 1956. Dissimilation of glucose at controlled pH values by pigmented and non-pigmented strains of *Escherichia coli*. *J Bacteriol* 72:497–499.
32. Six S, Andrews SC, Uden G, Guest JR. 1994. *Escherichia coli* possesses two homologous anaerobic C4-dicarboxylate membrane transporters (DcuA and DcuB) distinct from the aerobic dicarboxylate transport system (Dct). *J Bacteriol* 176:6470–6478. <http://dx.doi.org/10.1128/jb.176.21.6470-6478.1994>.
33. Baughn AD, Malamy MH. 2003. The essential role of fumarate reductase in haem-dependent growth stimulation of *Bacteroides fragilis*. *Microbiology* 149:1551–1558. <http://dx.doi.org/10.1099/mic.0.26247-0>.
34. Lancaster CRD. 2013. The di-heme family of respiratory complex II enzymes. *Biochim Biophys Acta* 1827:679–687. <http://dx.doi.org/10.1016/j.bbabi.2013.02.012>.
35. Yankovskaya V, Horsefield R, Törnroth S, Luna-Chavez C, Miyoshi H, Léger C, Byrne B, Cecchini G, Iwata S. 2003. Architecture of succinate dehydrogenase and reactive oxygen species generation. *Science* 299:700–704. <http://dx.doi.org/10.1126/science.1079605>.
36. Iverson TM, Luna-Chavez C, Cecchini G, Rees DC. 1999. Structure of the *Escherichia coli* fumarate reductase respiratory complex. *Science* 284:1961–1966. <http://dx.doi.org/10.1126/science.284.5422.1961>.
37. Tran QM, Rothery RA, Maklashina E, Cecchini G, Weiner JH. 2007. *Escherichia coli* succinate dehydrogenase variant lacking the heme b. *Proc Natl Acad Sci U S A* 104:18007–18012. <http://dx.doi.org/10.1073/pnas.0707732104>.
38. Tran QM, Fong C, Rothery RA, Maklashina E, Cecchini G, Weiner JH. 2012. Out of plane distortions of the heme b of *Escherichia coli* succinate dehydrogenase. *PLoS One* 7:e32641. <http://dx.doi.org/10.1371/journal.pone.0032641>.
39. Riebe O, Fischer RJ, Wampler DA, Kurtz DM, Jr, Bahl H. 2009. Pathway for H₂O₂ and O₂ detoxification in *Clostridium acetobutylicum*. *Microbiology* 155:16–24. <http://dx.doi.org/10.1099/mic.0.022756-0>.
40. Kawasaki S, Sakai Y, Takahashi T, Suzuki I, Niimura Y. 2009. O₂ and reactive oxygen species detoxification complex, composed of O₂-responsive NADH:rubredoxin oxidoreductase-flavoprotein A2-desulfoferredoxin operon enzymes, rubperoxin, and rubredoxin, in *Clostridium acetobutylicum*. *Appl Environ Microbiol* 75:1021–1029. <http://dx.doi.org/10.1128/AEM.01425-08>.
41. Wexler HM. 2007. *Bacteroides*: the good, the bad, and the nitty-gritty. *Clin Microbiol Rev* 20:593–621. <http://dx.doi.org/10.1128/CMR.00008-07>.
42. Betteken MI, Rocha ER, Smith CJ. 2015. Dps and DpsL mediate survival *in vitro* and *in vivo* during the prolonged oxidative stress response in *Bacteroides fragilis*. *J Bacteriol* 197:3329–3338. <http://dx.doi.org/10.1128/JB.00342-15>.
43. Mattevi A, Tedeschi G, Bacchella L, Coda A, Negri A, Ronchi S. 1999. Structure of L-aspartate oxidase: implications for the succinate dehydrogenase/fumarate reductase oxidoreductase family. *Structure* 7:745–756. [http://dx.doi.org/10.1016/S0969-2126\(99\)80099-9](http://dx.doi.org/10.1016/S0969-2126(99)80099-9).
44. Naqui A, Chance B, Cadenas E. 1986. Reactive oxygen intermediates in biochemistry. *Annu Rev Biochem* 55:137–166. <http://dx.doi.org/10.1146/annurev.bi.55.070186.001033>.
45. Condon C, Cammack R, Patil DS, Owen P. 1985. The succinate dehydrogenase of *Escherichia coli*. Immunochemical resolution and biophysical characterization of a 4-subunit enzyme complex. *J Biol Chem* 260:9427–9434.
46. Maklashina E, Iverson TM, Sher Y, Kotlyar V, Andréll J, Mirza O, Hudson JM, Armstrong FA, Rothery RA, Weiner JH, Cecchini G. 2006. Fumarate reductase and succinate oxidase activity of *Escherichia coli* complex II homologs are perturbed differently by mutation of the flavin binding domain. *J Biol Chem* 281:11357–11365. <http://dx.doi.org/10.1074/jbc.M512544200>.
47. Léger C, Heffron K, Pershad HR, Maklashina E, Luna-Chavez C, Cecchini G, Ackrell BA, Armstrong FA. 2001. Enzyme electrokinetics: energetics of succinate oxidation by fumarate reductase and succinate dehydrogenase. *Biochemistry* 40:11234–11245. <http://dx.doi.org/10.1021/bi010889b>.
48. Kowal AT, Werth MT, Manodori A, Cecchini G, Schröder I, Gunsalus RP, Johnson MK. 1995. Effect of cysteine to serine mutations on the properties of the [4Fe-4S] center in *Escherichia coli* fumarate reductase. *Biochemistry* 34:12284–12293. <http://dx.doi.org/10.1021/bi00038a024>.
49. Page CC, Moser CC, Chen X, Dutton PL. 1999. Natural engineering principles of electron tunnelling in biological oxidation-reduction. *Nature* 402:47–52. <http://dx.doi.org/10.1038/46972>.
50. Hudson JM, Heffron K, Kotlyar V, Sher Y, Maklashina E, Cecchini G, Armstrong FA. 2005. Electron transfer and catalytic control by the iron-sulfur clusters in a respiratory enzyme, *E. coli* fumarate reductase. *J Am Chem Soc* 127:6977–6989. <http://dx.doi.org/10.1021/ja043404q>.
51. Hincliffe P, Sazanov LA. 2005. Organization of iron-sulfur clusters in respiratory complex I. *Science* 309:771–774. <http://dx.doi.org/10.1126/science.1113988>.
52. Esterházy D, King MS, Yakovlev G, Hirst J. 2008. Production of reactive oxygen species by complex I (NADH:ubiquinone oxidoreductase) from *Escherichia coli* and comparison to the enzyme from mitochondria. *Biochemistry* 47:3964–3971. <http://dx.doi.org/10.1021/bi702243b>.
53. Birrell JA, Morina K, Bridges HR, Friedrich T, Hirst J. 2013. Investigating the function of [2Fe-2S] cluster N1a, the off-pathway cluster in complex I, by manipulating its reduction potential. *Biochem J* 456:139–146. <http://dx.doi.org/10.1042/BJ20130606>.
54. Liochev SI, Fridovich I. 1992. Fumarase C, the stable fumarase of *Escherichia coli*, is controlled by the *soxRS* regulon. *Proc Natl Acad Sci U S A* 89:5892–5896. <http://dx.doi.org/10.1073/pnas.89.13.5892>.
55. Flint DH, Emptage MH. 1988. Dihydroxy acid dehydratase from spinach contains a [2Fe-2S] cluster. *J Biol Chem* 263:3558–3564.
56. Pandelia ME, Nitschke W, Infossi P, Giudici-Orticoni MT, Bill E, Lubitz W. 2011. Characterization of a unique [FeS] cluster in the electron transfer chain of the oxygen tolerant [NiFe] hydrogenase from *Aquifex aeolicus*. *Proc Natl Acad Sci U S A* 108:6097–6102. <http://dx.doi.org/10.1073/pnas.1100610108>.
57. Chabrière E, Charon MH, Volbeda A, Pieulle L, Hatchikian EC, Fontecilla-Camps JC. 1999. Crystal structures of the key anaerobic enzyme pyruvate:ferredoxin oxidoreductase, free and in complex with pyruvate. *Nat Struct Biol* 6:182–190. <http://dx.doi.org/10.1038/5870>.
58. Seaver LC, Imlay JA. 2004. Are respiratory enzymes the primary sources of intracellular hydrogen peroxide? *J Biol Chem* 279:48742–48750. <http://dx.doi.org/10.1074/jbc.M408754200>.
59. Miller JH. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
60. Bacic MK, Smith CJ. 2008. Laboratory maintenance and cultivation of *Bacteroides* species. *Curr Protoc Microbiol* Chapter 13:Unit 13C.1. <http://dx.doi.org/10.1002/9780471729259.mc13c01s9>.
61. Koropatkin NM, Martens EC, Gordon JJ, Smith TJ. 2008. Starch catabolism by a prominent human gut symbiont is directed by the recognition of amylose helices. *Structure* 16:1105–1115. <http://dx.doi.org/10.1016/j.str.2008.03.017>.

62. Cheng VW, Piragasam RS, Rothery RA, Maklashina E, Cecchini G, Weiner JH. 2015. Redox state of flavin adenine dinucleotide drives substrate binding and product release in *Escherichia coli* succinate dehydrogenase. *Biochemistry* **54**:1043–1052. <http://dx.doi.org/10.1021/bi501350j>.
63. Imlay JA, Fridovich I. 1991. Assay of metabolic superoxide production in *Escherichia coli*. *J Biol Chem* **266**:6957–6965.
64. Messner KR, Imlay JA. 2002. In vitro quantitation of biological superoxide and hydrogen peroxide generation. *Methods Enzymol* **349**:354–361. [http://dx.doi.org/10.1016/S0076-6879\(02\)49351-2](http://dx.doi.org/10.1016/S0076-6879(02)49351-2).
65. McCord JM, Fridovich I. 1969. Superoxide dismutase. An enzymic function for erythrocyte hemocuprein (hemocuprein). *J Biol Chem* **244**:6049–6055.
66. Lancaster CRD, Kröger A, Auer M, Michel H. 1999. Structure of fumarate reductase from *Wolinella succinogenes* at 2.2 Å resolution. *Nature* **402**:377–385. <http://dx.doi.org/10.1038/46483>.