



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Letters to the Editor

Main challenges of electrolyte imbalance in older patients with COVID-19 and risk of QT prolongation**Principales desafíos del desequilibrio electrolítico en pacientes geriátricos con COVID-19 y riesgo de prolongación del intervalo QT****To the Editor,**

We read with interest the recent article published by Bonanad et al.¹

The topic of this consensus document is valuable and important; however, we encountered ambiguities and questions in one part of the article.

In table 3 of the article, in the description of the adverse cardiovascular effects of drugs investigated for COVID-19 treatment, the authors state that correction of hyperkalemia and hypermagnesemia is vital, while correction of hypokalemia and hypomagnesemia may be particularly imperative since low serum levels of potassium and magnesium enhances the possibility of QT prolongation.²

Hypokalemia, probably by modification of ion potassium channel function, can prolong the QT interval in a manner that results in heterogeneity and dispersion of repolarization. Similarly, hypomagnesemia is a well-established predisposing risk factor for torsade de pointes.³ In addition, potassium deficiency seems to be common in severe coronavirus disease 2019 (COVID-19).⁴ Several findings indicated that serum potassium should be maintained in the high normal range

(4.5–5.0 mmol/L), although more evidence is needed to support this practice.^{3,5}

Amir Rezazadeh^{a,b} and Niayesh Mohebbi^{a,b,*}

^aDepartment of Clinical Pharmacy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran

^bResearch Center for Rational Use of Drugs, Tehran University of Medical Sciences, Tehran, Iran

* Corresponding author:

E-mail address: nmohebbi@sina.tums.ac.ir (N. Mohebbi).

Available online 4 August 2020

REFERENCES

1. Bonanad C, García-Blas S, Tarazona-Santabalbina FJ, et al. Coronavirus: the geriatric emergency of 2020. Joint document of the Section on Geriatric Cardiology of the Spanish Society of Cardiology and the Spanish Society of Geriatrics and Gerontology. *Rev Esp Cardiol.* 2020;73:569–576.
2. Kallergis EM, Goudis CA, Simantirakis EN, Kochiadakis GE, Vardas PE. Mechanisms, risk factors, and management of acquired long QT syndrome: a comprehensive review. *ScientificWorldJournal.* 2012;2012:212178.
3. Drew BJ, Ackerman MJ, Funk M, et al. Prevention of torsade de pointes in hospital settings: a scientific statement from the American Heart Association and the American College of Cardiology Foundation endorsed by the American Association of Critical-Care Nurses and the International Society for Computerized Electrocardiology. *J Am Coll Cardiol.* 2010;55:934–947.
4. Lippi G, South AM, Henry BM. Electrolyte imbalances in patients with severe coronavirus disease 2019 (COVID-19). *Ann Clin Biochem.* 2020;57:262–265.
5. Thomas SH, Behr ER. Pharmacological treatment of acquired QT prolongation and torsades de pointes. *Br J Clin Pharmacol.* 2016;81:420–427.

<https://doi.org/10.1016/j.rec.2020.07.012>
1885-5857/

© 2020 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2020.05.001>
<https://doi.org/10.1016/j.rec.2020.08.018>

Main challenges of electrolyte imbalance in older patients with COVID-19 and risk of QT prolongation. Response**Principales desafíos del desequilibrio electrolítico en pacientes geriátricos con COVID-19 y riesgo de prolongación del intervalo QT. Respuesta****To the Editor,**

We appreciate the interest and comments from Rezazadeh et al.¹ regarding our article, and we agree on the relevance of electrolyte imbalance in patients with coronavirus 2019 disease (COVID-19), in particular hypokalemia and hypomagnesemia; the reference to hyperkalemia and hypermagnesemia corresponds to a translation error in the document, which has already been corrected. We would

like to point out that hypokalemia, as well as hyponatremia and hypocalcemia, appear to be common in patients with severe clinical forms of COVID-19.^{2–4} Infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) occurs thanks to the binding of the virus to angiotensin-converting enzyme 2, reducing the expression of the enzyme with a consequent increase in circulating angiotensin II, which promotes renal potassium loss.^{2,3,5} In addition, the inflammatory state and gastrointestinal losses can contribute to these imbalances, with potentially significant consequences due to the association with arrhythmic events secondary to QT interval prolongation (especially torsade de pointes), the risk for which increases with the use of certain drugs and also with the inflammatory state itself.^{1,5,6} The infection may also predispose to myocardial damage, especially in patients with existing cardiovascular disease.^{2,3} The pathophysiological changes in aging make older patients particularly vulnerable to electrolyte imbalances, so adequate monitoring and early correction are essential.⁷

Clara Bonanad,^a Pablo Díez-Villanueva,^{b,*} Sergio García-Blas,^a and Manuel Martínez-Sellés^{c,d}

SEE RELATED CONTENT:

<https://doi.org/10.1016/j.rec.2020.07.012>