



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

## Impact of COVID-19 on diagnoses, monitoring, and mortality in people with type 2 diabetes in the UK

The COVID-19 pandemic has had major health and economic effects across the world. At the time of writing (May, 2021), there have been more than 127 000 COVID-related deaths in the UK, with disproportionate effects in people with diabetes. Early in the pandemic, almost a third of all COVID-related deaths were in people with diabetes.<sup>1-3</sup>

The effect of COVID-19 on the National Health Service, and on diabetes services in particular, has been enormous, with frequent interruptions of most routine care. However, the consequences of these service restrictions on diagnosis and monitoring of type 2 diabetes are uncertain. To the best of our knowledge, there is also no available information comparing mortality across the UK nations in people with type 2 diabetes during the pandemic.

We generated a cohort of 25 million patients from 1831 UK general practices registered with the Clinical Practice Research Datalink (CPRD) Aurum (in England) and CPRD GOLD (in Northern Ireland, Scotland, and Wales) databases. The cohort included 14 929 251 patients (median age 41 years [IQR 25-59]; 7 461 565 [50.0%] of whom were women), who were followed between March 1 and Dec 10, 2020. We aimed to compare: (1) the UK-wide incidence of type 2 diabetes; (2) the frequency of HbA<sub>1c</sub> testing; and (3) mortality in people with type 2 diabetes, by nation, before and after the first nationwide COVID-19 lockdown, which began in March 23, 2020 (appendix pp 1-3). There were 790 377 (5.3%) people with type 2 diabetes in the cohort (median age 67 years (IQR 57-76); 348 515 (44.1%) of whom were women).

In April 2020, the rate reduction (RR) of new diagnoses of type 2 diabetes in primary care practices in England was 0.70 (95% CI 0.68-0.71) when compared with 10-year (January, 2010 to February, 2020) historical trends (figure A), with similar reductions in other UK nations (0.68 [0.70-0.66]). Older individuals (ie, those aged 65 years and older), men, and people from deprived areas had the greatest reductions in diagnosis rates (data not shown). The reduced diagnosis rates in April were mirrored by the reduced rates of new metformin prescriptions in general practices in England when compared with 10-year historical trends (0.53 [0.51-0.55]; appendix p 4).

Between May 1 and Dec 10, 2020, the reduced diagnosis rates recovered gradually but remained well below expected levels (figure A). In general practices in England overall, the RR of type 2 diabetes diagnoses was 0.32 (95% CI 0.28-0.35) between March 1 and Dec 10, 2020, with a smaller RR observed in other UK nations (0.21 [0.16-0.26]). Between March 1 and Dec 10, 2020, the rate of new metformin prescriptions in England decreased by 0.20 (0.16-0.23), with a similarly smaller reduction in other UK nations (0.12 [0.06-0.16]).

On the basis of these figures and data from the Office for National Statistics, we estimate that between March and December, 2020, there were approximately 60 000 missed or delayed type 2 diabetes diagnoses across the UK.

Rates of HbA<sub>1c</sub> testing in England (figure B) and other UK nations were greatly reduced in people with type 2 diabetes, with an RR of 0.77 (95% CI 0.76 to 0.78) in England and 0.83 (0.83-0.84) in other UK nations in April, 2020, and 0.31 (0.29-0.33) in England and 0.37 (0.34-0.39) in other UK nations overall between March and December, 2020. The largest reductions in the rates of HbA<sub>1c</sub> testing were observed in older people (data not shown).

In April 2020, mortality rates in people with type 2 diabetes in England were more than two-fold higher when compared with earlier trends between January, 2010, and February, 2020 (mortality rate increase 1.12 [95% CI 1.04-1.20]; figure C), with significantly smaller increases in other UK nations (0.65 [0.58-0.73]; figure D). Peaks in mortality were observed particularly in individuals aged older than 65 years (data not shown). Mortality rates returned to expected levels in people with type 2 diabetes between June and September, 2020, but began to increase again from October to November, 2020 (data for December are underestimated due to recording lags). Overall, between March 1 and Dec 10, 2020, the mortality rate in people with type 2 diabetes in general practices in England increased by 0.19 (95% CI 0.14-0.23) and by 0.13 (0.08-0.16) in all other UK nations (figure C, D).


Previous data on the effect of the COVID-19 pandemic on diagnosis of type 2 diabetes are scarce. A study from Salford, UK, showed that there were 135 fewer diagnoses (a 49% reduction) of type 2 diabetes than expected between March and May, 2020.<sup>4</sup> In the current study, we extend these observations by showing that the reduced diagnosis rate applies to all areas of the UK. To our knowledge, no study has reported the effect of the COVID-19 pandemic on HbA<sub>1c</sub> monitoring or has compared national mortality rates in people with type 2 diabetes across different UK nations.

Type 2 diabetes develops over many years; therefore, it seems unlikely that people's behaviour during the pandemic has reduced the true incidence of this condition. If the true incidence has remained constant, our data suggest that, across the UK, there were nearly 60 000 missed or delayed diagnoses of type 2 diabetes between March and December, 2020. This figure could be an underestimation because of well documented behaviours promoting



Published Online  
May 11, 2021  
[https://doi.org/10.1016/S2213-8587\(21\)00116-9](https://doi.org/10.1016/S2213-8587(21)00116-9)

See Online for appendix



**Figure:** Observed and expected incident diagnoses of type 2 diabetes in England (A), HbA<sub>1c</sub> monitoring in patients with type 2 diabetes in England (B), and monthly mortality rates in people with type 2 diabetes in England (C), and in Northern Ireland, Scotland, and Wales (D). In A–D, purple lines indicate observed monthly rates, and blue shaded regions indicate expected rates with 95% CIs; the vertical grey line at March 1, 2020, separates the rates in primary care before and after the start of the COVID-19 pandemic, and x-axis markers indicate mid-months. Rates were derived by use of data from CPRD Aurum, covering 21 797 864 patients in England, and CPRD GOLD, covering 3 625 958 patients in Northern Ireland, Scotland, and Wales. CPRD=Clinical Practice Research Datalink.

weight gain during lockdown, such as poor diet, reduced physical activity, increased alcohol consumption, poorer mental health, and reduced sleep quality when compared with before lockdown.<sup>5–7</sup> These data are a clinical concern because undiagnosed type 2 diabetes can contribute to serious long-term complications.<sup>8</sup>

The reduction in HbA<sub>1c</sub> testing is another important concern for people with type 2 diabetes, because they, and their treating clinicians, often rely solely on HbA<sub>1c</sub> data to make treatment decisions. We observed reductions in new prescriptions for

insulin, especially in older individuals (data not shown), suggesting a failure to intensify therapy in people with poorly controlled, long-term type 2 diabetes. There are already concerns about clinical inertia in diabetes management, with failures to escalate care when glucose control is poor.<sup>9</sup> These HbA<sub>1c</sub> data indicate potential further delays that are predicted to cause avoidable diabetes-related long-term complications.

The higher COVID-related mortality rate in people with diabetes compared with the general population has been well documented.<sup>1–3</sup> Herein, we add

to these data by showing regional differences in the effect of COVID-19 on mortality rates in people with type 2 diabetes, with higher rates observed in England than in the rest of the UK. Further research is required to understand how population characteristics, including ethnicity, population density, and social deprivation, might explain these differences.

To our knowledge, this is the first UK-wide study reporting the indirect effects of the COVID-19 pandemic on the diagnosis of type 2 diabetes, related prescribing rates, and HbA<sub>1c</sub>

testing in primary care. Our findings in general practices in England were replicated by use of data from other regions of the UK. By combining assessments of diabetes coding and prescribing, our data support the conclusion that reduced rates of type 2 diabetes diagnoses can be genuinely explained by missed diagnoses.

We could not explore ethnicity-related variation in care because ethnicity coding was available in only 40% of records. It is possible that some type 2 diabetes diagnoses could have been made in hospital settings, with delays in updating primary care coding. However, these hospital-based diagnoses would not explain the reductions in metformin prescribing, and it is known that people have tended to avoid attending hospitals during the pandemic. Finally, our findings might not be generalisable to health-care systems outside the UK, due to differences in diabetes care, differing COVID-19 prevalences and effects on health-care systems, and differing COVID-19 measures and policies.

In conclusion, we highlight reductions in the diagnosis and monitoring of type 2 diabetes during the COVID-19 pandemic, which have important clinical and public health implications. Over the coming months, health-care services will need to manage this predicted backlog, and the anticipated deterioration of blood glucose levels and cardiovascular risk factors due to delayed diagnoses and reduced monitoring of patients with established diabetes. Our data showed that older individuals, men, and people

from areas of high deprivation were most adversely affected and could represent specific groups to target for early intervention. Moving forward, during and beyond the pandemic, effective public health communication should ensure that patients remain engaged with diabetes services and make use of HbA<sub>1c</sub> monitoring and remote consultations.<sup>10</sup>

HT reports grants from Dexcom and personal fees from Eli Lilly. NK reports grants and personal fees from Novo Nordisk, Boehringer Ingelheim, Napp Pharmaceuticals, Sanofi, AstraZeneca, Nevro Corporation, and Abbott. NM reports personal fees from Napp Pharmaceuticals, Merck Sharpe & Dohme UK, AstraZeneca, Boehringer Ingelheim, OmniaMed Communications, Takeda UK, Lilly Diabetes, Mylan, MGP, Novo Nordisk, Abbott, Sanofi, Roche, and Nevro; and non-financial support from Napp Pharmaceuticals, Novo Nordisk, and Sanofi. DMA reports grants from AbbVie, Almirall, Celgene, Eli Lilly, Novartis, Janssen, UCB Pharma, and the Leo Foundation. MKR reports personal fees and non-promotional lecture fees from Novo Nordisk. MJC, AKW, and LL declare no competing interests. The study is based on data from the CPRD obtained under license from the UK Medicines and Healthcare products Regulatory Agency. The study and use of CPRD data were approved by the Independent Scientific Advisory Committee for CPRD research (protocol number, 20\_182R).

\**Matthew J Carr*†, *Alison K Wright*†, *Lalantha Leelarathna*, *Hood Thabit*, *Nicola Milne*, *Naresh Kanumilli*, *Professor Darren M Ashcroft*†, *Professor Martin K Rutter*†  
matthew.carr@manchester.ac.uk

†Contributed equally

Centre for Pharmacoepidemiology and Drug Safety, Division of Pharmacy and Optometry, School of Health Sciences (MJC, AKW, DMA), and Division of Diabetes, Endocrinology and Gastroenterology, School of Medical Sciences (AKW, LL, HT, MKR), Faculty of Biology, Medicine and Health, University of Manchester, Manchester M13 9PT, UK; Manchester Academic Health Science Centre, Manchester, UK (MJC, AKW, LL, HT, NM, NK, DMA, MKR); NIHR Greater Manchester Patient Safety

Translational Research Centre, University of Manchester, Manchester, UK (MJC, DMA); Diabetes, Endocrinology and Metabolism Centre, Manchester University NHS Foundation Trust, Manchester, UK (LL, HT, NM, NK, MKR)

- Holman N, Knighton P, Kar P, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol* 2020; **8**: 823–33.
- Barron E, Bakhal C, Kar P, et al. Associations of type 1 and type 2 diabetes with COVID-19-related mortality in England: a whole-population study. *Lancet Diabetes Endocrinol* 2020; **8**: 813–22.
- Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020; **584**: 430–36.
- Williams R, Jenkins DA, Ashcroft DM, et al. Diagnosis of physical and mental health conditions in primary care during the COVID-19 pandemic: a retrospective cohort study. *Lancet Public Health* 2020; **5**: e543–50.
- COVID Symptom Study. Has lockdown influenced our eating habits? The silent pandemic: how lockdown is affecting future health. 2020. <https://covid.joinzoe.com/post/lockdown-weight-gain> (accessed April 3, 2021).
- Robinson E, Gillespie S, and Jones A. Weight-related lifestyle behaviours and the COVID-19 crisis: an online survey study of UK adults during social lockdown. *Obes Sci Pract* 2020; **6**: 735–40.
- Robinson E, Boyland E, Chisholm A, et al. Obesity, eating behavior and physical activity during COVID-19 lockdown: a study of UK adults. *Appetite* 2021; **156**: 104853.
- Simmons RK, Griffin SJ, Lauritzen T, and Sandbæk A. Effect of screening for type 2 diabetes on risk of cardiovascular disease and mortality: a controlled trial among 139 075 individuals diagnosed with diabetes in Denmark between 2001 and 2009. *Diabetologia* 2017; **60**: 2192–99.
- Khunti K, Kosiborod M, and Ray KK. Legacy benefits of blood glucose, blood pressure and lipid control in individuals with diabetes and cardiovascular disease: time to overcome multifactorial therapeutic inertia? *Diabetes Obes Metab* 2018; **20**: 1337–41.
- Wake DJ, Gibb FW, Kar P, et al. Endocrinology in the time of COVID-19: remodelling diabetes services and emerging innovation. *Eur J Endocrinol* 2020; **183**: G67–77.